



Journal Homepage: www.journalijar.com

INTERNATIONAL JOURNAL OF ADVANCED RESEARCH (IJAR)

Article DOI: 10.21474/IJAR01/22363

DOI URL: <http://dx.doi.org/10.21474/IJAR01/22363>



Case Report

A CASE OF REFRACTORY SEIZURES IN JUVENILE CANAVAN DISEASE

Gudluru Saiharsha¹ and P Pravin Prabhu²

1. PG, Department of General Medicine, AVMCH, Puducherry.

2. Assistant Professor, Department of General Medicine, AVMCH, Puducherry.

Manuscript Info

Manuscript History

Received: 6 October 2025

Final Accepted: 8 November 2025

Published: December 2025

Abstract

Introduction: Canavan disease is a rare, progressive neurological disorder classified as a leukodystrophy, caused by mutations in the ASPA gene on chromosome 17, resulting in a deficiency of the enzyme aspartoacylase. This leads to the accumulation of N-acetyl aspartic acid (NAA) in the brain. There are two forms of the disease: a severe infantile/neonatal form and a milder juvenile form, both presenting symptoms such as hypotonia, intellectual disability, feeding difficulties, paralysis, and seizures, with severity varying between forms. Diagnosis is achieved by detecting increased NAA levels in blood, urine, and cerebrospinal fluid, and MRI shows characteristic bilateral white matter changes. Prognosis is poor for infantile form, while juvenile form patients have better outcomes.

"© 2025 by the Author(s). Published by IJAR under CC BY 4.0. Unrestricted use allowed with credit to the author."

Introduction:-

Case Report :

A 22-year-old male presented with a history of seizures, experiencing nine episodes over three days, each lasting 1-2 minutes and involving all four limbs, mouth frothing, upward eye rolling, and involuntary urination (GTCS). He had a normal early childhood but began having seizures at age 10, followed by loss of motor skills and speech difficulties. He experienced seizures approximately every two months, especially when off anti-epileptic medications. His parents were related (second-degree consanguinity), but there's no family history of seizures or intellectual disability. The patient has moderate mental retardation, cannot walk, exhibits spasticity and exaggerated deep tendon reflexes, has a positive bilateral Babinski reflex, and myoclonus in the right lower limb, while the sensory system remains intact. MRI findings showed diffuse white matter leukodystrophy, and MR spectroscopy indicated increased N-acetyl aspartic acid (NAA) peaks.

Conclusion:-

This case highlights the importance of considering rare genetic leukodystrophies such as juvenile Canavan disease in young patients presenting with refractory seizures and intellectual disability. Management remains supportive, focusing on seizure control, rehabilitation, speech therapy, and long-term supportive care.

Corresponding Author:-Gudluru Saiharsha

Address:-PG, Department of General Medicine, AVMCH, Puducherry.