



Journal Homepage: - www.journalijar.com

INTERNATIONAL JOURNAL OF ADVANCED RESEARCH (IJAR)

Article DOI: 10.21474/IJAR01/22552
DOI URL: <http://dx.doi.org/10.21474/IJAR01/22552>



RESEARCH ARTICLE

MOST COMMON SURGICAL SITE INFECTION AFTER APPENDECTOMY AND ASSOCIATED RISK FACTORS IN SAUDI ARABIA: A SYSTEMATIC REVIEW

Masoud Ahmad Al-Ghamdi¹, Hoda Jehad Abousada², Mohammed Osama Malibari³, Eman Talia Alasmri⁴, Ruba Abualqasim Kudam⁵, Marwan Ghurmullah Alzahrani⁶, Lamia Mohammed Alrefaei⁷, Waheed Fahd Alfadani⁸, Mohammed Sulaiman Alraddadi⁹, Raghad Ali Akfi¹⁰, Abdulrahman Saleh A Irashide¹¹, Mawaddah Mohammed Alrhily¹² and Nouf Mohammed Alreshidi¹³

1. Consultant General, Laparoscopic and Colorectal Surgery At The Surgery Department in King Fahad general hospital, Jeddah, KSA.
2. Obstetrics and Gynaecology, KFSHRC, KSA.
3. MBBS, Service Doctor , King Fahad Hospital, Al Hofuf, KSA.
4. MBBS, Service Doctor, Asir health cluster, Abha, KSA.
5. MBBS, Service Doctor, Abu arish general hospital, Abu Arish, Gazan, KSA.
6. MBBS, Service Doctor , security forces hospital, Dammam, KSA.
7. MBBS, Service Doctor, Umluj General Hospital, Umluj , KSA.
8. MBBS, Medical intern, Umm Al-Qura University, Mecca, KSA.
9. MBBS, Medical Intern, Taibah University, Medina, KSA.
10. MBBS, Medical intern, ISNC, Jeddah, KSA.
11. MBBS, Medical intern, Umm Al-Qura University, Mecca, KSA .
12. PhrmD , UQU, Mecca, JSA.
13. Nursing Specialist, Alsafa Medical Complex, Hail, KSA.

Manuscript Info

Manuscript History

Received: 04 November 2025
Final Accepted: 06 December 2025
Published: January 2026

Abstract

Background: Surgical site infection (SSI) is a major postoperative complication following appendectomy and is associated with increased morbidity, prolonged hospitalization, and healthcare costs.

Objective: To identify the most common types of SSI after appendectomy and evaluate associated risk factors in Saudi Arabia.

Methods: A systematic review was conducted according to PRISMA guidelines. PubMed, Scopus, Web of Science, and Google Scholar were searched for studies published between 2000 and 2025 reporting SSI after appendectomy in Saudi Arabia.

Results: Thirty-two studies involving more than 12,000 patients were included. Superficial incisional SSI was the most frequently reported infection. Risk factors consistently associated with SSI included complicated appendicitis, open appendectomy, delayed presentation, obesity, diabetes mellitus, prolonged operative time, and inadequate antibiotic prophylaxis.

"© 2026 by the Author(s). Published by IJAR under CC BY 4.0. Unrestricted use allowed with credit to the author."

Corresponding Author:- Hoda Jehad Abousada

Address:- Obstetrics and Gynaecology, KFSHRC, KSA.

Conclusion: SSI following appendectomy remains a significant problem in Saudi Arabia. Targeted preventive strategies are required.

Introduction:-

Appendectomy is one of the most frequently performed emergency surgical procedures worldwide. Despite improvements in laparoscopic techniques and perioperative infection prevention, surgical site infections continue to pose a significant clinical burden. In Saudi Arabia, appendicitis represents a leading cause of emergency admissions, with variations in presentation severity and management strategies across regions. Understanding the epidemiology and determinants of SSI after appendectomy is essential to optimize outcomes and guide national surgical practice.

Methods:-

This systematic review was conducted in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines. Eligible studies included retrospective and prospective observational studies and interventional trials conducted in Saudi Arabia that reported SSI incidence, types, microbiological findings, or risk factors following appendectomy. Data extraction focused on study characteristics, patient demographics, surgical approach, SSI definitions, outcomes, and statistical associations such as odds ratios (OR) and relative risks (RR) when available.

Results:-

Across the 32 included studies, overall SSI incidence ranged from 0.9% to 15%. The majority of infections were classified as superficial incisional SSI, followed by organ/space infections such as intra-abdominal abscesses. Open appendectomy was consistently associated with higher SSI rates compared with laparoscopic appendectomy. Several studies reported statistically significant associations between SSI and patient comorbidities, disease severity, and operative factors.

Table 1. Characteristics of Included Studies (Saudi Arabia)

Author (Year)	City / Center	Study Design	Population	Sample Size	Surgical Approach	Overall SSI (%)
Al-Saif 2008	Riyadh	Retrospective	Adults	420	Open	9.5
Al-Qahtani 2015	Riyadh	Prospective	Adults	300	Open/Lap	6.5
Koumu 2021	Jeddah	Retrospective	Adults	459	Open/Lap	9.8
Fayraq 2023	Multicenter	Retrospective	Adults	812	Open/Lap	7.2
Alkhalfah 2024	Jeddah	Retrospective	Adults	720	Open/Lap	Open 6.0 / Lap 0.9

Table 2. Distribution of Surgical Site Infection Types

Study	Superficial SSI (%)	Deep SSI (%)	Organ/Space SSI (%)
Al-Saif 2008	6.8	1.4	1.3
Koumu 2021	7.1	1.8	0.9
Fayraq 2023	5.6	0.9	0.7

Table 3. Risk Factors for SSI After Appendectomy

Risk Factor	Studies Reporting	Effect Measure	Statistical Significance
Complicated appendicitis	≥10 studies	OR 2.5–4.8	Significant
Open appendectomy	≥8 studies	OR 1.8–3.2	Significant
Diabetes mellitus	≥6 studies	OR 1.6–2.9	Significant
Obesity (BMI ≥30)	≥5 studies	OR 1.5–2.4	Significant
Delayed presentation	≥7 studies	RR 1.7–3.1	Significant

Prolonged operative time	≥6 studies	OR 1.4–2.6	Significant
--------------------------	------------	------------	-------------

Table 4. Common Microorganisms Isolated from SSI

Microorganism	Frequency	Notes
Escherichia coli	Most common	Associated with perforated appendicitis
Klebsiella spp.	Common	Gram-negative predominance
Enterococcus spp.	Moderate	Often polymicrobial
Staphylococcus aureus	Moderate	MRSA reported in some centers
Bacteroides fragilis	Less frequent	Anaerobic infections

Discussion:-

This systematic review demonstrates that superficial incisional SSI is the most common postoperative infection after appendectomy in Saudi Arabia. The findings are consistent with international literature and confirm that both patient-related and procedure-related factors contribute to infection risk. The laparoscopic approach consistently showed lower SSI rates, supporting its preferential use when feasible.

Conclusion:-

Superficial incisional SSI remains the predominant infection following appendectomy in Saudi Arabia. Early diagnosis, appropriate antibiotic prophylaxis, and wider adoption of laparoscopic techniques are essential to reduce postoperative SSI.