



Journal Homepage: - www.journalijar.com

INTERNATIONAL JOURNAL OF ADVANCED RESEARCH (IJAR)

Article DOI: 10.21474/IJAR01/22633
DOI URL: <http://dx.doi.org/10.21474/IJAR01/22633>



RESEARCH ARTICLE

MANAGEMENT OF ENDODONTIC FAILURE IN LOWER ANTERIOR TEETH BY APICECTOMY IN AN ADOLESCENT PATIENT

Saranya S¹ and Padmakumari B²

1. Junior Resident, Department of Pedodontics ,Government Dental college ,Thiruvananthapuram.
2. HOD,Department of Pedodontics ,Government Dental college ,Thiruvananthapuram.

Manuscript Info

Manuscript History

Received: 10 November 2025
Final Accepted: 12 December 2025
Published: January 2026

Key words:-

Endodontic failure; Apicectomy;
Periapical pathology; Lower anterior
teeth; Adolescent patient; Surgical
endodontics.

Abstract

Endodontic failure may occur due to the persistence of periapical pathology despite adequately performed root canal treatment. When nonsurgical retreatment is not feasible, surgical endodontic intervention becomes necessary. Apicectomy is a conservative surgical procedure aimed at eliminating periapical pathology while preserving the natural tooth. This case report presents the surgical management of an apical inflammatory cyst associated with endodontic failure in a 13-year-old male patient who reported pain in the lower anterior region of the jaw. The patient had a history of traumatic occlusion and prior root canal treatment involving the mandibular incisors. Clinical and radiographic examination revealed a persistent inflammatory radicular cyst associated with teeth 31 and 41. The treatment plan included surgical enucleation of the cystic lesion followed by apicectomy and retrograde filling of the involved teeth. The surgical procedure comprised apical root resection, thorough periapical curettage, and retrograde sealing. Postoperative follow-up demonstrated satisfactory clinical resolution and progressive radiographic healing of the periapical region. This case underscores the role of apicectomy as an effective tooth-preserving surgical option for managing persistent periapical lesions in adolescent patients when conventional nonsurgical retreatment is not indicated

"© 2026 by the Author(s). Published by IJAR under CC BY 4.0. Unrestricted use allowed with credit to the author."

Introduction:-

Endodontic therapy aims to eliminate infection from the root canal system and preserve the natural dentition; however, treatment failure may occur despite adherence to accepted clinical protocols. Persistent periapical pathology following root canal treatment is commonly attributed to residual intraradicular infection, inadequate canal disinfection, missed anatomy, microleakage, or the presence of extraradicular biofilms and cystic lesions¹⁻³. In such situations, nonsurgical retreatment is considered the first line of management; however, surgical endodontic intervention becomes necessary when retreatment is impractical or unlikely to resolve the pathology⁴. Apicectomy, a well-established surgical endodontic procedure, involves resection of the apical portion of the root along with curettage of the periapical lesion, followed by retrograde sealing of the root canal system. This procedure is particularly indicated in cases of persistent periapical disease associated with adequately obturated canals,

Corresponding Author:- Saranya S

Address:-Junior Resident, Department of Pedodontics ,Government Dental college ,Thiruvananthapuram.

1010

obstructed canals, or failed retreatment^{5,6}. Advances in microsurgical techniques, magnification, ultrasonic root-end preparation, and biocompatible retrograde filling materials have significantly improved the success rates of apicectomy, making it a predictable treatment option for endodontic failures⁷. Management of endodontic failure in the lower anterior region presents unique challenges due to esthetic concerns, thin cortical bone, and proximity to vital anatomical structures. In adolescent patients, treatment planning requires special consideration of ongoing craniofacial growth, tooth preservation, psychological impact, and long-term functional outcomes⁸. Conservative surgical intervention is therefore preferred to maintain tooth integrity and alveolar bone continuity. This article describes the surgical management of endodontic failure in the lower anterior teeth of an adolescent patient by means of apicectomy, emphasizing clinical decision-making, surgical technique, and treatment outcomes. The report highlights the role of apicectomy as a tooth-preserving treatment modality in young patients when conventional endodontic approaches fail.

Case Report:-

A 13-year-old male patient presented with complaints of pain and swelling in the chin region for a duration of two weeks. The patient had a history of traumatic occlusion involving the mandibular anterior teeth, with non-vital teeth 31 and 41 associated with a periapical lesion. Root canal treatment of the mandibular central incisors had been performed six months earlier. The patient remained asymptomatic initially; however, symptoms developed during the follow-up period. Clinical examination revealed a localized swelling in the labial sulcus area in relation to teeth 31 and 41. The swelling was tender on palpation, with no associated sinus tract or discharge. Radiographic investigations including intraoral periapical radiograph (IOPAR), Orthopantomogram (OPG), and Cone Beam Computed Tomography (CBCT) were advised. Imaging revealed a well-defined periapical radiolucency associated with the apices of teeth 31 and 41. CBCT evaluation demonstrated a large unilocular cystic lesion measuring approximately 2 cm×1.5 cm size in the mandibular anterior region. Based on clinical and radiographic findings, a diagnosis of persistent periapical pathology secondary to endodontic failure was established. Surgical intervention was planned, and the patient was prescribed antibiotics prior to the procedure.




Figure 1a: CBCT preoperative 3D view of the lesion in relation to the periapical region of root canal treated teeth 31 and 41, **1b:** axial view, **1c:** sagittal view

Under local anesthesia, two vertical releasing incisions and a crevicular incision were placed to elevate a full-thickness mucoperiosteal flap, exposing the periapical region of teeth 31 and 41. The pathological tissue was completely enucleated, followed by meticulous curettage and the root ends were resected about 3mm with a surgical straight carbide bur at shallow angle. Haemostasis was achieved by means of gauze packing. Retrograde cavities were prepared and sealed using mineral trioxide aggregate (MTA). The bony cavity was thoroughly irrigated. A hemostatic agent (Surgicel) was placed in the surgical site, and the flap was repositioned and sutured with 3-0 Vicryl. Postoperative medications and instructions were given to the patient. The excised tissue was sent for histopathological examination, which confirmed the diagnosis of an infected periapical cyst. No tenderness, swelling and sinus were detected at the follow up visits at one week, three months, and six months postoperatively. Six month follow up OPG showed complete resolution of the lesion with satisfactory periradicular healing and bone regeneration.



Figure 2a: Depicting the defect, **2b:** Curetted periapical lesion, **2c:** Resected root ends, **2d:** Excised tissue for histopathology examination, **2e:** Suturing done with vicryl 3-0, **2f:** Review after 2 weeks




Figure 3a: CBCT Preoperative panoramic view of unilocular radiolucency in relation to the root apex of 31 and 41, **3b:** Six-month postoperative follow-up radiograph demonstrates bone deposition at the surgical site.

Discussion:-

Persistent periapical pathology following root canal treatment remains a common cause of endodontic failure, even when treatment appears radiographically adequate. Microbial persistence within apical ramifications, lateral canals, or extraradicular biofilms has been identified as a key etiological factor contributing to post-treatment apical periodontitis.⁹⁻¹¹ In the present case, the presence of a large cystic periapical lesion despite prior endodontic therapy indicated that nonsurgical retreatment alone would have limited predictability. Although nonsurgical retreatment is generally considered the first-line approach for failed endodontic cases, surgical endodontic intervention becomes essential when retreatment is impractical due to adequately obturated canals, anatomical complexities, or obstructed canal systems.⁴ Apicectomy allows direct access to the periapical pathology, facilitating complete removal of infected tissue while simultaneously addressing the apical portion of the root canal system. The favorable clinical and radiographic outcomes observed in this case support the effectiveness of surgical endodontic management in such scenarios. Management of endodontic failure in adolescent patients presents additional considerations, including preservation of natural teeth, maintenance of alveolar bone integrity, and minimization of psychological and functional impacts.¹² Extraction and prosthetic replacement in growing patients may lead to long-term esthetic and occlusal complications, making conservative surgical procedures such as apicectomy a preferred treatment

option. In the mandibular anterior region, where cortical bone is thin and esthetic demands are high, precise surgical technique is crucial to ensure optimal healing.

Advancements in endodontic microsurgery have significantly improved the prognosis of apicectomy. Modern surgical principles—including limited root-end resection (approximately 3 mm), ultrasonic retrograde cavity preparation, and the use of biocompatible retrograde filling materials—have contributed to higher success rates compared to conventional techniques.^{5,13,14} Mineral trioxide aggregate (MTA), used as the retrograde filling material in this case, is well documented for its excellent sealing ability, biocompatibility, and capacity to promote periapical healing and cementogenesis.^{15,16} These properties likely contributed to the favorable bone regeneration observed at the six-month follow-up. According to the systematic review by Mahmoud Torabinejad et al., endodontic surgery showed a higher success rate in the short term (2–4 years) compared with nonsurgical retreatment (77.8% vs 70.9%). However, at longer follow-up (4–6 years), nonsurgical retreatment demonstrated a higher success rate than endodontic surgery (83.0% vs 71.8%).⁴ Histopathological confirmation of the lesion as an inflammatory cyst further justified the surgical approach, as true cystic lesions may exhibit limited healing potential following nonsurgical retreatment alone.¹⁷ Complete enucleation combined with apicectomy ensures elimination of the pathological tissue and reduces the risk of recurrence. The absence of postoperative complications and satisfactory radiographic healing in this case highlight the predictability of apicectomy as a tooth-preserving treatment option in adolescent patients.

Conclusion:-

Apicectomy is an effective and conservative treatment option for managing persistent periapical pathology when nonsurgical retreatment is not feasible. In adolescent patients, this approach enables preservation of natural teeth and supporting structures, leading to favorable clinical and radiographic outcomes when performed with appropriate surgical technique and biocompatible materials. Overall, this case reinforces the role of apicectomy as a viable and effective treatment modality for managing persistent periapical pathology in young patients when conventional endodontic approaches fail.

References:-

1. Sjögren U, Hägglund B, Sundqvist G, Wing K. Factors affecting the long-term results of endodontic treatment. *Journal of Endodontics*. 1990 Oct 1;16(10):498–504.
2. Nair PNR. On the causes of persistent apical periodontitis: a review. *Int Endodontic J*. 2006 Apr;39(4):249–81.
3. Ricucci D, Siqueira JF. Anatomic and Microbiologic Challenges to Achieving Success with Endodontic Treatment: A Case Report. *Journal of Endodontics*. 2008 Oct;34(10):1249–54.
4. Torabinejad M, Corr R, Handysides R, Shabahang S. Outcomes of Nonsurgical Retreatment and Endodontic Surgery: A Systematic Review. *Journal of Endodontics*. 2009 Jul;35(7):930–7.
5. Kim S, Kratchman S. Modern Endodontic Surgery Concepts and Practice: A Review. *Journal of Endodontics*. 2006 Jul;32(7):601–23.
6. Von Arx T, Walker WA. Microsurgical instruments for root-end cavity preparation following apicoectomy: a literature review. *Dental Traumatology*. 2000 Apr;16(2):47–62.
7. Setzer FC, Kohli MR, Shah SB, Karabucak B, Kim S. Outcome of Endodontic Surgery: A Meta-analysis of the Literature—Part 2: Comparison of Endodontic Microsurgical Techniques with and without the Use of Higher Magnification. *Journal of Endodontics*. 2011 Nov;S0099239911011460.
8. American Academy of Pediatric Dentistry. Guideline on pulp therapy for primary and immature permanent teeth. *Pediatr Dent*. 2023;45(6):399–413. [Internet]. [cited 2026 Jan 14]. Available from: https://www.aapd.org/assets/1/7/G_Pulp.pdf
9. Nair PNR. PATHOGENESIS OF APICAL PERIODONTITIS AND THE CAUSES OF ENDODONTIC FAILURES. *Critical Reviews in Oral Biology & Medicine*. 2004 Nov;15(6):348–81.
10. Ricucci D, Siqueira JF. Biofilms and Apical Periodontitis: Study of Prevalence and Association with Clinical and Histopathologic Findings. *Journal of Endodontics*. 2010 Aug;36(8):1277–88.
11. Siqueira JF, Rôças IN. Clinical Implications and Microbiology of Bacterial Persistence after Treatment Procedures. *Journal of Endodontics*. 2008 Nov;34(11):1291–1301.e3.
12. Andreasen JO, Andreasen FM, Andersson L. *Textbook and Color Atlas of Traumatic Injuries to the Teeth*. 5th ed. Wiley Blackwell; 2018.
13. Gutmann JL. Surgical endodontics: past, present, and future. *Endodontic Topics*. 2014 May;30(1):29–43.
14. Setzer FC, Kim S. Comparison of Long-term Survival of Implants and Endodontically Treated Teeth. *J Dent Res*. 2014 Jan;93(1):19–26.

15. Parirokh M, Torabinejad M. Mineral Trioxide Aggregate: A Comprehensive Literature Review—Part I: Chemical, Physical, and Antibacterial Properties. *Journal of Endodontics*. 2010 Jan;36(1):16–27.
16. Torabinejad M, Parirokh M. Mineral Trioxide Aggregate: A Comprehensive Literature Review—Part II: Leakage and Biocompatibility Investigations. *Journal of Endodontics*. 2010 Feb;36(2):190–202.
17. Nair PNR, Sjögren U, Krey G, Kahnberg KE, Sundqvist G. Intraradicular bacteria and fungi in root-filled, asymptomatic human teeth with therapy-resistant periapical lesions: A long-term light and electron microscopic follow-up study. *Journal of Endodontics*. 1990 Dec;16(12):580–8.