



Journal Homepage: [-www.journalijar.com](http://www.journalijar.com)  
**INTERNATIONAL JOURNAL OF  
ADVANCED RESEARCH (IJAR)**

Article DOI: 10.21474/IJAR01/22859  
DOI URL: <http://dx.doi.org/10.21474/IJAR01/22859>



**RESEARCH ARTICLE**

**CEREBRAL VENOUS THROMBOSIS DURING TUBERCULOUS  
MENINGOENCEPHALITIS: A RARE CASE IN THE ICU**

Armelbouchra<sup>1</sup>, Byadimahmoud<sup>2</sup>, Mahasekal<sup>2</sup>, Hamzazerouali<sup>2</sup>, Othmanesebbata<sup>2</sup> and El Moussaoui Rachid<sup>2</sup>

1. Emergency Critical Care Department IBN Sina University Hospital, Rabat.
2. Department of Intensive Care Unit, IBN Sina University Hospital, Rabat.

**Manuscript Info**

**Manuscript History**

Received: 16 December 2025  
Final Accepted: 18 January 2026  
Published: February 2026

**Key words:-**

tuberculosis, altered consciousness,  
cerebral venous thrombosis,  
anticoagulation

**Abstract**

Cerebral venous thrombosis is an uncommon neurovascular complication, marked by significant clinical and etiological heterogeneity. Infectious causes account for fewer than 10% of cases, with tuberculous involvement of the central nervous system representing an exceptional etiology. We report the case of a 25-year-old patient presenting with febrile impaired consciousness and anisocoria, in whom the medical history revealed close contact with a confirmed case of tuberculosis. Initial laboratory investigations and cerebrospinal fluid analysis were highly suggestive of tuberculous infection of the central nervous system, despite the absence of Mycobacterium tuberculosis isolation from CSF. Cerebral MR angiography demonstrated findings consistent with tuberculous meningoencephalitis complicated by cerebral venous thrombosis. Management included initial stabilization, endotracheal intubation followed by tracheostomy, anti-tuberculosis therapy, and full-dose anticoagulation. Clinical evolution was favorable, with progressive neurological recovery. This case highlights the rarity and diagnostic complexity of the coexistence of tuberculous meningoencephalitis and cerebral venous thrombosis, and underscores the need for prompt, comprehensive management combining anti-tuberculosis treatment, corticosteroid therapy, and anticoagulation.

"© 2026 by the Author(s). Published by IJAR under CC BY 4.0. Unrestricted use allowed with credit to the author."

**Introduction:-**

Tuberculosis remains a major public health issue in Morocco, with approximately 30,000 new cases reported annually across all forms, of which tuberculous meningitis (TBM) accounts for about 0.8% (1). Despite its low prevalence, TBM remains one of the most severe forms of the disease, characterized by a frequently delayed diagnosis and high morbidity and mortality (2). Cerebral venous thrombosis (CVT) is a rare vascular complication of the central nervous system, with an incidence estimated at 1.5 cases per 100,000 population. It may be favored by infectious, inflammatory, or immunological disorders (3). The association between TBM and CVT is exceptional and only rarely described in the literature. This coexistence presents a major diagnostic challenge due to the nonspecific nature of clinical manifestations, as well as therapeutic and prognostic complexities. We report the case of a young patient with tuberculous meningoencephalitis complicated by cerebral venous thrombosis, illustrating the diagnostic

and therapeutic challenges and underscoring the importance of appropriate management in a tuberculosis-endemic setting.

### **Observation:-**

A 25-year-old patient, an active smoker (18 pack-years) and cannabis user, with a history of tuberculosis exposure (mother treated for pulmonary and gastrointestinal tuberculosis), was admitted to the intensive care unit for febrile altered consciousness associated with anisocoria. The medical history revealed, over the preceding three weeks, signs suggestive of tuberculosis, including fever, night sweats, and unquantified weight loss, followed by headaches, vomiting, and progressive deterioration of consciousness.

On admission, the patient had a Glasgow Coma Scale (GCS) score of 6, necessitating orotracheal intubation and initial stabilization. Non-contrast brain CT demonstrated active quadriventricular hydrocephalus with several hypodense areas associated with cerebral edema.

### **A lumbar puncture was performed, revealing:**

- CSF protein: 7.62 g/L
- CSF glucose: 0.29 g/L (with concomitant blood glucose of 1.34 g/L)
- White blood cells: 8 cells/mm<sup>3</sup>
- Red bloodcells: 9,600/mm<sup>3</sup>
- Direct examination:negative
- GeneXpert PCR for Mycobacterium tuberculosis: negative.

Despite the absence of bacillus identification, the combination of clinical, radiological, and laboratory findings pointed towards tuberculous meningitis. Standard anti-tuberculosis therapy, combined with corticosteroid treatment, was initiated. Further evaluation with cerebral MR angiography revealed abnormalities consistent with tuberculous meningoencephalitis complicated by cerebral venous thrombosis. Curative anticoagulation was subsequently started.

### **The clinical course was favorable, marked by:**

- the performance of a tracheostomy on day 10.
- Neurological improvement (GCS 9)
- Resolution of the febrile syndrome
- Improvement of hemodynamic parameters and transcranial Doppler finding
- Follow-up CT scan showed stable hydrocephalus without any new visible complications

### **Discussion:-**

The coexistence of tuberculous meningitis (TBM) and cerebral venous thrombosis (CVT) is uncommon and remains only sparsely documented in the literature. This association poses diagnostic challenges, as both conditions may present with overlapping symptoms such as headache, vomiting, altered consciousness, and signs of intracranial hypertension. In regions where tuberculosis is endemic, maintaining a high index of suspicion is crucial to prevent diagnostic delays that could lead to serious clinical consequences. In a recent series by Li et al. involving 28 patients with tuberculous meningitis complicated by cerebral venous thrombosis, the mortality rate reached 17.9%. Among the survivors, most achieved a satisfactory functional recovery following the initiation of antituberculous therapy combined with anticoagulation (4). Several isolated case reports have also described this association, emphasizing its rarity but also its potential severity (Table 1).

From a pathophysiological standpoint, multiple mechanisms have been proposed to explain the development of cerebral venous thrombosis in the context of tuberculous meningitis. These include direct endothelial injury caused by granulomatous inflammation, a prothrombotic state driven by the systemic inflammatory response, and platelet activation that has been documented in severe forms of tuberculosis (5,6). These mechanisms collectively contribute to an increased risk of thrombus formation within the cerebral venous sinuses. The diagnosis may be challenging due to the non-specific nature of clinical signs and the overlap between the manifestations of cerebral venous thrombosis and those of tuberculous meningitis itself. Magnetic resonance venography remains the reference imaging modality, as it can detect CVT even when clinical findings or initial CT imaging do not suggest the diagnosis (7). In our case, tuberculous meningitis was strongly suspected based on clinical, biological, and radiological findings, despite the absence of Mycobacterium tuberculosis isolation in the cerebrospinal fluid—an occurrence that is common in this condition. Management is based on a three- or four-drug antituberculous regimen combined with corticosteroid

therapy, which helps reduce meningeal inflammation and the risk of neurological complications. Anticoagulation is recommended in cases of cerebral venous thrombosis, unless contraindicated, to prevent thrombus extension and improve overall prognosis. In our case, the early initiation of these treatments led to a favorable clinical outcome. This case highlights the importance of early diagnosis and a multidisciplinary approach when managing this rare association, particularly in regions with a high prevalence of tuberculosis.

**Tableau 1 review of the literature for cerebral venous thrombosis during tuberculosis**


article	Number of cases	evolution
Aseptic cerebral venous thrombosis and cerebral tuberculomas complicating miliary tuberculosis.(8)	1 case	Good evolution
Tuberculosis, an uncommon cause of cerebral venous thrombosis(9)	1 case	Good evolution
Septic cavernous sinus thrombosis caused by tuberculosis infection(10)	1 case	Oculomotorpare sis
Superior sagittal sinus thrombosis caused by calvarial tuberculosis (11)	1	Good evolution

**Conclusion:-**

the occurrence of cerebral venous thrombosis during tuberculous meningoencephalitis represents a rare but serious complication, often challenging to recognize due to overlapping clinical manifestations. This case emphasizes the need for heightened vigilance in regions with high tuberculosis prevalence and the early use of vascular imaging when atypical neurological progression is observed. Prompt management combining antituberculous therapy, corticosteroids, and anticoagulation can improve prognosis and minimize neurological sequelae.



**Figure 1 Non-contrast brain CT scan demonstrating quadriventricular hydrocephalus with hypodense regions**



**Figure 2 Cerebral MR angiography demonstrating septic involvement of the telencephalon associated with cerebral venous thrombosis**

**Informed consent :**

Written informed consent was obtained from the patients and their legal guardians for publication of this case report and the accompanying images. A copy of the consent is available for review by the Editor-in-Chief of this journal

**Figure and table:-**

**Table 2 review of the literature for cerebral venous thrombosis during tuberculosis**

Figure 1 Non-contrast brain CT scan demonstrating quadriventricular hydrocephalus with hypodense regions

Figure 2 Cerebral MR angiography demonstrating septic involvement of the telencephalon associated with cerebral venous thrombosis

**Conflict of interest:**

The authors declare that they have no conflict of interest.

**Author Contributions:**

All authors contributed to the realization of this work

**Reference:-**

1. Chahboune M, Barkaoui M, Iderdar Y, Alwachami N, Mourajid Y, Ifleh M, et al. [Epidemiological profile and diagnostic and evolutionary features of TB patients at the Diagnostic Centre for Tuberculosis and Respiratory Diseases in Settati, Morocco]. *Pan Afr Med J.* 2022;42:185.
2. Taous A, Berri MA, Boulahri T, Traibi I, Amrani HN, Rouimi A. [Neuromeningeal tuberculosis: clinical, paraclinical, therapeutic and evolutionary profile of 21 patients]. *Pan Afr Med J.* 2022;43:214.
3. Zhou LW, Yu A Y X, Ngo L, Hill MD, Field TS. Incidence of Cerebral Venous Thrombosis: A Population-Based Study, Systematic Review, and Meta-Analysis. *Stroke.* Janv 2023;54(1):169-77.
4. Li Q, Han J, Wang Y, Song Y. Cerebral venous thrombosis (CVT) complicating tubercular meningitis. *BMC Neurol.* 24 juin 2023;23(1):245.
5. The Haematological and Biochemical Changes in Severe Pulmonary Tuberculosis. *QJM: An International Journal of Medicine* [Internet]. déc 1989 [cité 4 oct 2025]; Disponible sur: <https://academic.oup.com/qjmed/article/73/3/1151/1582749/The-Haematological-and-Biochemical-Changes-in>
6. Baynes RD, Bothwell TH, Flax H, McDonald TP, Atkinson P, Chetty N, et al. Reactive thrombocytosis in pulmonary tuberculosis. *J Clin Pathol.* juin 1987;40(6):676-9.

7. Bonneville F. Imaging of cerebral venous thrombosis. *Diagnostic and Interventional Imaging*. déc 2014;95(12):1145-50.
8. Messouak O, Amara B, Benjelloun FZ, Tizniti S, Benjelloun MC, Belahsen MF. Thrombose veineuse cérébrale aseptique et tuberculomes cérébraux compliquant une miliaire tuberculeuse pulmonaire. *Revue Neurologique*. févr 2007;163(2):238-40.
9. Fiorot Júnior JA, Felício AC, Fukujima MM, Rodrigues CA, Morelli VM, Lourenço DM, et al. Tuberculosis: an uncommon cause of cerebral venous thrombosis? *Arq Neuro-Psiquiatr*. sept 2005;63(3b):852-4.
10. Xia P, Jiao Y. Septic cavernous sinus thrombosis caused by tuberculosis infection. *BMJ Case Reports*. 25 nov 2014;2014:bcr2014206209.
11. Sundaram PK, Sayed F. SUPERIOR SAGITTAL SINUS THROMBOSIS CAUSED BY CALVARIAL TUBERCULOSIS: CASE REPORT. *Neurosurgery*. avr 2007;60(4):E776.