



Journal Homepage: [-www.journalijar.com](http://www.journalijar.com)

## INTERNATIONAL JOURNAL OF ADVANCED RESEARCH (IJAR)

Article DOI: 10.21474/IJAR01/23088  
DOI URL: <http://dx.doi.org/10.21474/IJAR01/23088>



### RESEARCH ARTICLE

## PHYTOCHEMICAL PROPERTIES AND IN VITRO ANTIOXIDANT ACTIVITY OF LAWSONIA INERMIS (L) EXTRACTS

Dembele Bintou, Tounkara Fatoumata, Cisse Salimatou and Badiaga Mamadou

### Manuscript Info

#### Manuscript History

Received: 12 January 2026

Final Accepted: 14 February 2026

Published: March 2026

#### Key words:-

Lawsonia inermis,  
Extracts, Phytochemistry,  
Antioxidant-activities

### Abstract

The aim of this study was to determine the phytochemical composition and antioxidant activity of Lawsonia inermis extracts, a plant traditionally used in Mali. The sample (Leaves, bark roots and trunk bark) were collected in Kayes. The polyphenols, flavonoids, tannins and anthocyanins of the extracts were quantified using spectrophotometry and identified via HPLC. The antioxidant activity was assessed using 2, 2-azinobis-3-ethylbenzothiazoline-6-sulfonic acid (ABTS) and 2, 2-Diphenyl-1-Picrylhydrazyl (DPPH) radicals scavenging capacities. The leaf extracts were mainly in flavonoids; Polyphenols, anthocyanins, however, they showed low tannins compositions. The values were 9.21 mg ECt; 7.77 mg EAG; 5.61 mg ECt; 0.74 mg ECG per 1 g of dry matter (DM) respectively for flavonoids, polyphenols, anthocyanins and tannins. The HPLC analysis of the extracts revealed the presence in Leaf: of Apigenin, Chlorogenic and Lawsone acids were also found, in the bark of Roots: Protocatechic Acid. All extracts showed good ABTS and DPPH radical scavenging capacities. The antioxidant activities of samples correlated with their phytochemical compositions.

"© 2026 by the Author(s). Published by IJAR under CC BY 4.0. Unrestricted use allowed with credit to the author."

### Introduction:-

Lawsonia inermis, or henna, is a thorny shrub 2 to 5 meters tall, native to Arabia and Iran, but also found in Mediterranean and subtropical Africa, Madagascar, Asia, and Australia. The plant has been cultivated in the Sahel and Sudanian regions near dwellings. *L. inermis* prefers sandy soils (Arbonnier, 2002). According to Muhammad HS and Muhammad S in 2005, chloroform and aqueous extracts of the leaves inhibit the growth of bacteria responsible for wound infections. In vivo studies conducted on rats reportedly showed hepatoprotective activity of the ethanolic extract of the leaves (Ahmed et al., 2000). In vitro studies of the methanolic extract of the leaves reportedly showed immunomodulatory activity (Mikhaeil et al., 2004). According to Kaplan and Hammerman in 2006, a deficiency in certain enzymes such as glucose-6-phosphate dehydrogenase causes side effects (hemolysis, anemia, reticulocytosis and hyperbilirubinemia). Lawsonia inermis contains quinonoid derivatives of xanthenes, flavonoids, coumarins, mannites, Gallic and tannic acids as well as lawsone which is thought to be responsible for the plant's colour (Kelmanson et al, 2002; Ong et al, 2009). Siddiqui and Kardar isolated and identified two triterpenes (lawsonic acid and lawsonin) in the plant's leaves in 2001. Chloroform extract revealed the presence of seven phenolic compounds in the leaves: p-coumaric acid, lawsone, apigenin, luteolin, 2-methoxy-3-methyl-1.4-naphthoquinone, cosmosin, and apigin. The ABTS [2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid)] assay of each of these isolated compounds showed antioxidant activity comparable to that of ascorbic acid (Mikhaeil et al., 2004). The role of phenolic

**Corresponding Author:-Tounkara Fatoumata**

Address:-Laboratory of Biosciences and Applications, Faculty of Sciences and Techniques (FST),  
University of Sciences, Techniques and Technologies of Bamako (USTTB), BP: 3206, Bamako, Mali.

compounds in protecting against certain diseases is well established due to their potential interaction with numerous enzymes and their antioxidant properties (Zhang, 2003). Specifically, flavonoids are attributed with a variety of biological properties: antitumor, free radical scavenging, anti-inflammatory, analgesic, anti-allergic, antispasmodic, antibacterial, and anticancer. Several in vitro and in vivo studies have focused on evaluating the antibacterial and antifungal properties of polyphenols. Currently, this effect is demonstrated by numerous experimental studies. Moroh et al., in 2008, demonstrated the antibacterial effect of acetic extracts rich in alkaloids and flavonoids of *Morindamorindoides* on eight strains of *Escherichia coli*, bacteria commonly found in diarrhea in infants and children up to the age of five. The inhibitory activity of flavonoids on bacterial growth was studied by Katarzyna et al in 2007 who showed that many flavonoid compounds have an effect on different gram negative (*Escherichia coli*...) and gram positive (*Staphylococcus aureus*...) bacterial strains. Given the limited number of scientific studies devoted to the plant and its use in traditional medicine, the objective of this study was to identify the polyphenolic compounds present in the leaves, trunk bark, and root bark, and to evaluate their antioxidant effects, which would justify the plant's uses in traditional medicine.

## **Material and Methods:-**

### **Material:-**

The samples (Leave, trunk bark, roots bark) of *Lawsonia inermis* were collected in Kayes (Mali). They have been transported and identified to the Department of Traditional Medicine (DMT) under the number (0903). Folin-Ciocalteu, Gallic, Protocatechic, Chlorogenic, Caffeic acids, Lawsone, Rutin, Apigenin, Quercetin, Kaempferol. ABTS and DPPH were provided by the companies SIGMA-Aldrich (France) and Across organics (Belgium). All other chemicals and solvents used were obtained from a commercial source and were of analytical grade.

### **Methods:-**

#### **Samples Preparation:-**

After the initial cleaning process, samples (the different parts of the plant) were dried in the shade and at room temperature in the Natural Substances laboratory of the FST. After drying, the samples were pounded using a laboratory scale hammer miller and the resulting powder sieved until a fine powder was obtained.

#### **Preparation of Extracts:-**

The preparation of the sample extracts was carried out by the method described by Abu Bakar et al., 2009, with slightly modified. To 5 g of plant powder were added 2 times 20 mL of a 50/50 (V/V) hydromethanolic solution for UV-Visible Spectroscopy and HPLC. The mixtures were stirred for 6 hours, and then filtered through a 0.45 µm millipore membrane. The filtrates collected were centrifuged at 1500 ×g for 20 min. The extracts obtained have been cooled and conserved at (+4°C) in bottles before analysis.

#### **Total polyphenol content (TPC):-**

The total polyphenolic content was determined using the Folin-Ciocalteu method. The plant extract (0.1 ml) was mixed with 0.1 ml of the Folin-Ciocalteu reagent (1 N). The mixture was left at room temperature for 5 minutes. Then, 1 ml of 7% sodium carbonate ( $\text{Na}_2\text{CO}_3$ ) was added to the mixture. After 90 minutes of incubation at 23°C, absorbances were measured at 750 nm against a blank prepared with methanol. Note that a calibration curve was established before using gallic acid under the same conditions as the samples to be analyzed. The results obtained are expressed in mg gallic acid equivalent per g of dry matter (mg EGA/g). All samples were analyzed at least three times, and the mean was calculated.

#### **Total Flavonoid Content (TFC):-**

Total flavonoids were assessed colorimetrically. A 0.25 mL aliquot of the plant extract was taken, and 1 mL of distilled water and 75 µL of 5% sodium nitrate ( $\text{NaNO}_3$ ) were added. After 5 minutes, 75 µL of 10% aluminum trichloride ( $\text{AlCl}_3$ ) were added to the mixture. After 6 minutes, 0.5 mL of 1 M sodium hydroxide ( $\text{NaOH}$ ) and 0.6 mL of distilled water were added to the mixture while stirring. A calibration curve was developed using standard catechin solutions prepared at different concentrations. The absorbance of the mixture was determined by UV spectrometry at 510 nm relative to a blank prepared with water. The total flavonoid content was expressed as mg catechin equivalent per gram of dry matter (mgCE/g). All samples were analyzed at least three times, and the mean was calculated.

**Total Anthocyanin Content (TAC):-**

Total anthocyanin compounds were evaluated colorimetrically using a UV spectroscope. The method used was based on the change in absorbance using two buffers: one containing 0.025 M potassium chloride (KCl) (pH 1) and the other 0.4 M sodium acetate (pH 4.5). 0.2 mL of the extract was mixed with 1.8 mL of each buffer solution. The absorbance of the solution was determined at 510 nm and 700 nm against a blank prepared with methanol. The change in absorbance was calculated using the following formula:

$$\Delta A = [(A_{510} - A_{700}) \times \text{pH}1] - [(A_{510} - A_{700}) \times \text{pH}4] \quad (1)$$

The concentration of anthocyanin pigment in the extract is expressed in mg cyanidin-3-glycoside equivalent per liter of solution or in mg cyanidin-3-glycoside equivalent per gram of dry matter (mg CG/g) by comparison to a calibration curve established with cyanidin-3-glycoside. All samples were analyzed at least three times, and the average was calculated. The anthocyanin concentration was calculated using the following formula:

$$C(\text{mg/L}) = \Delta A \times Mm \times Df \times (100/Ma) \quad (2)$$

With  $\Delta A$ : the change of absorbances, Mm: molecular mass of cyanidin (449.2);  
Df: dilution factor; Ma: molecular absorptivity (26.900).

**Total tannin content (TTC):-**

In a test tube containing 1.5 mL of concentrated sulfuric acid, we added 50  $\mu\text{L}$  of extract and 3 mL of a 4% methanol-vanillin solution. The mixture was left to stand for 15 minutes. Absorbance was measured at 500 nm against a methanol blank. The calibration curve was developed using standard solutions of catechins prepared at different concentrations. The calibration curve has been established with a correlation coefficient  $R^2 = 0.9899$ . The results are expressed in mg equivalent catechins per 1 g of dry matter (mg ECt/1 g DM). All samples were analyzed at least three times.

**HPLC Analysis:-**

The HPLC analysis was carried out according to the method described by Muanda et al., 2011, with a slight modification, using an elution gradient consisting of three phases. Solvent A: 50 mM ammonium phosphate ( $\text{NH}_4\text{H}_2\text{PO}_4$ ) at pH 2.6 (adjusted with ortho phosphoric acid), solvent B: (80/20 (v/v)) acetonitrile/solvent A, and solvent C: 200 Mm ortho phosphoric acid ( $\text{H}_3\text{PO}_4$ ) at pH 1.5 (pH was adjusted with 0.1 M NaOH). After preparation, the solvents were put in an ultrasonic device for 10 min for homogenization. The profile of the gradient used for 60 min is presented in Table 1. The elution flow rate was 1 mL/min and the injection loop capacity 20  $\mu\text{L}$ . Detection was performed at 280 and 320 nm. Standard phenolic compounds (9 standards) were prepared by dissolving 2 mg/mL. In each sample, the phenolic compound was identified by the retention time of the corresponding standard and the concentration of the phenolic compound was calculated by comparing the peak areas. The samples were analyzed at least three times. After each cycle, the system was reconditioned 10 minutes before a new analysis.

**Table 1. Profile of the gradient used for 60 min.**

T (mn)	A%	B%	C%
0-4	100	0	0
4-10	92	8	0
10-22.5	0	14	86
22.5-27.5	0	16	84
27.5-50	0	25	75
50-55	0	20	80
55-60	100	0	0

**Antioxidant Activity Assay:-****Scavenging capacity of ABTS radicals:-**

The method developed by Kim et al., 2002, slightly modified has been used in this experiment. 1.0 mM AAPH was mixed with 2.5 mM ABTS using buffer. The buffer solution consists of 100 mM potassium phosphate (pH 7.4) containing 150 Mm NaCl. The mixture was heated in a water bath at 68°C for 20 minutes until the concentration of

the blue-green ABTS radical complex gives an absorbance of between  $0.65 \pm 0.02$  at 734 nm. To 60  $\mu\text{L}$  of the sample has been added 2.94 mL of the radical blue-green solution of ABTS. The mixture was incubated in a water bath at  $37^\circ\text{C}$  for 20 minutes. The control consists of 60  $\mu\text{L}$  of methanol and 2.94 mL of ABTS and was checked for each series of samples. The absorbance was measured at 734 nm. Stable radical scavenging activity in the ABTS test of phenolic compounds was expressed in mg equivalent vitamin C (mg EVC/100 (DM)). The radical solution was prepared daily. All samples were analyzed at least three times

#### Scavenging capacity of DPPH radicals:-

The DPPH radical scavenging activity was determined using the method developed by Hoste et al., 2006, with some modifications. To 2.90 ml of an aqueous solution of 50% methanol (100 mM of DPPH), 100  $\mu\text{L}$  of plant extract were added. The mixture has been heated in a water bath at  $20^\circ\text{C}$  away from light for 40 min. The blank was prepared with (100  $\mu\text{L}$  of 50% methanol and 2.90 mL of the DPPH solution) and checked for each series of samples. The decrease in absorbance was measured at 517 nm 40 minutes later. The free radical scavenging activity in the DPPH test of total phenolic compounds was expressed in mg equivalents vitamin C (mg EVC/100 g (DM)). The radical solution has been prepared daily. All samples were analyzed at least three times.

#### Statistical Analysis:-

The results were processed with software such as: Excel version 2019 and Minitab 18.1, for analysis of variance (ANOVA) was used to compare the mean values of these varieties with the Fischer test at the probability threshold  $P = 0.05$ .

### Results and Discussion:-

#### Total Polyphenolic Content (TPC):-

Figure 1 shows the results of the phenolic compound composition of *L. inermis* extracts analyzed by UV-visible spectroscopy. Analysis of the results shows that the different extracts are rich in phenolic compounds (flavonoids, anthocyanins, tannins, and polyphenols). The leaf showed the lower in total tannins.




Figure 1: Total Polyphenolic Content (TPC) of the *L. inermis* extracts

#### HPLC Analysis:-

The results of HPLC analyses of *L. inermis* extracts are shown in Figure 2 (chromatograms of leaf and root bark extracts). Analysis of this figure reveals the presence of some polyphenolic compounds in the both extracts. However the results of qualitative and quantitative analyses of the different *L. inermis* extracts are given in Table 2. Analysis of these results shows that Protocatechic acid ( $1503.54 \mu\text{g/ml}$ ) is the predominant compound in the root extracts, while apigenin ( $166.40 \mu\text{g/ml}$ ) is the predominant compound in the leaf extracts. However, the leaf extracts also contain chlorogenic acid ( $60.17 \mu\text{g/ml}$ ) and lawsone ( $9.07 \mu\text{g/ml}$ ). Our results are comparable to


those of Handa et al, in 1997 and Mebe et al, in 1998 on the extract of the leaves of *L.inermis*, in particular by the presence of flavonoids, coumarins, naphthoquinones and triterpenes. In contrast, the instrument did not detect any of our standards in the trunk bark extracts. It should be noted that not all compounds were identified during our analyses due to a lack of suitable standards.

**Table 2. HPLC Analysis results of the *L. inermis* extracts.**


Names of compound	Rt(min)	Leaves $\mu\text{g/mL}$	Trunk Bark $\mu\text{g/mL}$	Bark of Roots $\mu\text{g/MI}$
Gallic Acid	09.65	Nd	Nd	Nd
Protocatechic Acid	11.85	Nd	Nd	1503.54 $\pm$ 40.80
Chlorogenic Acid	16.72	60.17 $\pm$ 0.4.07	Nd	Nd
Caffeic Acid	19.55	Nd	Nd	Nd
Lawsone acid	25.98	09.07 $\pm$ 0.12	Nd	Nd
P-Cumaric Acid	33.90	Nd	Nd	Nd
Rutin	35.50	Nd	Nd	Nd
Quercetin	39.82	Nd	Nd	Nd
Apigenin	41.41	166.40 $\pm$ 0.4. 34	Nd	Nd
Kaempferol	42.52	Nd	Nd	Nd

Results reported are means of triplicate samples  $\pm$  standard deviation. Rt: Retention time, Nd: no detected.

a/



b/






Figure 2: Chromatogram profiles of the extracts of Leave (a and b), of root bark (c).

#### Antioxidant Activity:-

The results of the antioxidant activity analysis of extracts from different parts of *L. inermis* are reported in Table 3. The result showed that the extract of *L. inermis* leaves exhibits an antioxidant activity of 2.49 mg VCE by ABTS, which could explain the presence of apigenin (166.4 µg/g dry matter) in *L. inermis* leaves. Mikhaeilet al. demonstrated in 2004 that the flavonoids contained in *L. inermis* leaves have antiradical activity comparable to that of ascorbic acid; in particular, apigenin-7-apiosyl glucoside showed a high potential for scavenging free radicals. These compounds act as free radical scavengers by preventing the peroxidation of the phospholipid membrane and protecting immunomodulatory cells. This could explain the hepatoprotective effect of henna (Anaad et al., 1992).

Table 3. ABTS and DPPH radical scavenging activities of *L. inermis* extract

Samples	ABTS-Test (g EVC/100 g (DM))	DPPH-Test (g EVC/100 g (DM))
Leaves	2.49±0/06	2.23±0.002
Trunk Bark	1.74±0/004	2.0±0/01
Bark of Roots	0.96±0..01	2.15±0.003

Results reported are means of triplicate samples ± standard deviation

#### Conclusion:-

The results of this study revealed the presence of Apigenin and Protocatechic Acid with considerable quantity in the extracts of our studied plant. These chemical constituents are known to have the ability to scavenge free radicals from ABTS and DPPH. This explains the best antioxidant activities of our plant extracts. In sum, the results of phytochemistry and antioxidant activity analysis of *L. inermis* extracts revealed that it is a medicinal plant.

#### Conflicts of Interest:-

The authors declare no conflicts of interest regarding the publication of this paper.

#### References:-

1. Abu Bakar, M., Mohamed, M., Rahmat, A., Fry J. (2009). Phytochemicals and antioxidant activity of different parts of bambangan (*Mangiferapajang*) and tarap (*Artocarpusodoratissimus*), *Food Chem*, 113(3), 483-479.
2. Ahmed S., Rahman A., Alam A., Saleem M., Athar M., Sultana S. (2000). Evaluation of the efficacy of *Lawsonia Alba* in the alleviation of carbon tetrachloride-induced oxidative stress. *Journal of Ethnopharmacology*, 69(2), 157-164
3. Anaad, K. K., Singh, B., Chand, D., and Chandon, B., K. (1992). An evaluation of *Lawsonia Alba* extract as hepatoprotective agent. *Planta Med.*, 58(1), 22-25.
4. Arbonnier, M., 2002. Arbres, arbustes et lianes des zones sèches d'Afrique de l'ouest. 2nd Edn. CIRADMNHN. 573 pages.
5. Hoste, H., F., Jackson, S., Athanasiadou, S., M., Thamsborg, Hoskin, S., O. (2006). The effects of tannin-rich plants on parasitic nematodes in ruminants. *Trends in Parasit*, 22(6), 253-261.

6. Kaplan, M., Hammerman, C. (2006). Glucose-6-phosphate dehydrogenase-deficient neonates: a potential cause for Concern in North American- Pediatrics, 106(6), 1478-1480
7. Katarzyna, U., Anna, M., Marta, M., Joanna, J.; B., and Grzegorz, W. (2007). Assessment of antibacterial effects of flavonoids by estimation of generation times in liquid bacterial cultures. *Biologia*, 62 (2), 132-135.
8. Kelmanson, J., E., Jager A., K., Staden, J., V. (2002). Zulu medicinal plants with antibacterial activity. *Journal Ethnopharm*, 69(2000), 241-246
9. Kim, D. O., Lee, K., W., Lee, H., J., and Lee, C.,Y. (2002). Vitamin C equivalent antioxidant capacity VCEAC) of phenolic phytochemicals. *J. Agric. Food Chem.*, 50(13), 3713-3717.
10. Kim, D. O., Chun, O., K., Kim, Y., J., Moon, H., Y., and Lee, C., Y. (2003). Quantification of polyphenolics and their antioxidant capacity in fresh plums. *J. Agric. Food Chem.*, 51(22), 6509-6515.
11. Mikhail, B., R., Badria, F., A., Maatooq G.T., and Amer, M., M., A. (2004). Antioxidant and Immunomodulatory Constituents of Henna Leaves. Department of Pharmacology, Faculty of Pharmacy, Mansura University Egypt. 59c: 468- 476 p.
12. Moroh, J., L., A., Bahi, C., Dje, K., Loukou Y., G., and Guede-Guina, F. (2008). Study of the antibacterial activity of the acetate extract (ACE) of *Morindamorindoides* (Baker) *milne-redheat* (Rubiaceae) on the in vitro growth of *Escherichia coli* strains. *Bulletin de la Societe Royale des Sciences de Liège*, 77, 44–61
13. Muanda, F., D. Kone, A. Dicko, R. Soulimani, Younos, C. (2011). Phytochemical composition and antioxidant capacity of three Malian medicinal plant parts. *Evidence-Based Complem. And Altern. Med.*, doi: 10.1093/ecam/nep109
14. Muhammad, H., S., and Muhammad, S. (2005). The use of *Lawsonia inermis* (Henna) in the management of burn wound infections. *African Journal of Biotechnology*, 4 (9), 934-937
15. Ong, C., Y., Ling, S., K., Ali, R., M., Chee, C., F., Samah, Z., A., Siong, A., Ho, H., Teo, S., H., Lee, H., B. (2009). Systematic analysis of in vitro photo- cytotoxic activity in extracts from terrestrial plants in Peninsula Malaysia for Photodynamic Therapy. *Journal of Photochemistry and Photobiology B: Biology*, 96, 216-222
16. Siddiqui, B., S., and Kardar, M., N. (2001). Triterpenoids from *Lawsonia Alba*. *Phytochemistry*, 58: 1195-1198
17. Zhang, L., Henriksson, G., and Gellerstedt, G. (2003). The formation of  $\beta$ -  $\beta$  Structures in lignin biosynthesis – are there two different pathways? *Org. Biomol. Chem.*, 1, 3621-3624