



Journal Homepage: www.journalijar.com
**INTERNATIONAL JOURNAL OF
ADVANCED RESEARCH (IJAR)**

Article DOI: 10.21474/IJAR01/23097
DOI URL: <http://dx.doi.org/10.21474/IJAR01/23097>



RESEARCH ARTICLE

**ATYPICAL MANIFESTATION OF CUTANEOUS INFANTILE HEMANGIOMAS:
TWO RARE CASE REPORTS**

Furquana Niaz¹, Tehreem Aslam², Zainab Zaidi², Aisha Miral², Pankhri Mangi² and Waquar Ahmad³

1. Professor and Head of Dermatology Department Naimat Begum/Hamdard University Hospital, Karachi.
2. MBBS, Hamdard University, Karachi.
3. Consultant Paediatrician at WWL NHS Trust Hospital, UK.

Manuscript Info

Manuscript History

Received: 14 January 2026
Final Accepted: 16 February 2026
Published: March 2026

Key words:-

Hemangioma, strawberry naevus, infantile hemangioma, vascular tumor, vascular malformation, propranolol, superficial hemangioma.

Abstract

Background: Infantile hemangioma (IH), or strawberry naevus, is a benign, bright-red vascular birthmark caused by rapid blood vessel growth, affecting 4-10% of infants. It usually involves the head and neck, rarely the limbs and genitalia, grows rapidly for 12–15 months, and then gradually regresses over years. Treatment includes topical beta-blockers and steroids, oral corticosteroids and beta blockers, laser therapy, or surgical options in certain cases. In this case report, we included atypical presentations of IH in infants.

The aim of study: To assess atypical presentations, changes in size, morphology, progression and treatment response in varieties of hemangioma.

Case Reports: We presented two cases with rare and uncommon, atypical presentations of IH who came to NBH/HUH, Karachi. Case 1 involved a 3-month-old female with a typical presentation characterized by a solitary, asymmetric rapidly enlarging vascular lesion on right deltoid region, treated with 2% topical propranolol and oral steroids resulting in regression with color changes and atrophy of the lesion over a period of one year. Case 2 involved a 3-month-old male with an atypical presentation characterized by multiple, bilateral symmetrical lesions (atypical manifestation) on extensor surfaces of both wrists and the penile area, treated with 2% propranolol but patient was lost to follow up.

Conclusion: Early Recognition of such uncommon manifestations is essential to avoid misdiagnosis, exclude other differentials, anticipate common complications and initiate timely treatment, particularly in high-risk anatomical sites to prevent morbidity and good treatment response. However, the limited number of reported cases necessitates further studies to better characterize atypical presentations of infantile hemangiomas.

"© 2026 by the Author(s). Published by IJAR under CC BY 4.0. Unrestricted use allowed with credit to the author."

Introduction:-

The most prevalent benign vascular tumors of infancy are infantile hemangiomas (IH), which usually manifest in the first few weeks of life and affect 4% to 10% in Caucasian infants, with some variation across populations. [1] IH is more common in female infants (female to male ratio of 1.4:1 to 3.1), twins, premature infants, Caucasians, low birth weight infants, multiple pregnancies, preeclampsia, placental anomalies, and positive family history of IH in a first-degree relative. IHs are clinically characterized by a proliferative phase followed by regression phase, but 10% of cases require early treatment due to its location, size and complications otherwise spontaneous regression do not require any treatment. Clinical features of IH are not present at birth but develop in the first 1-2 weeks after birth, rapidly multiply within 1–3 months of age, stop proliferating until 5 months and then slowly regress into adipose and fibrous tissue until 4 years but up to the age of 10 years in some cases.

Pathologically, IHs are glucose transporter-1 protein (GLUT-1) positive, which distinguishes them from other vascular tumors but its pathogenesis has not been completely explained. There are several key hypotheses explaining IH. The first suggests that increased VEGFR signaling and mutations in hemangioma stem cells (derived from CD34⁺/CD133⁺ endothelial progenitor cells) promote differentiation into GLUT-1 positive endothelial cells under angiogenic factors like VEGF-A and HIF-1 α . The second, the placental embolization theory, proposes that displaced placental cells drive hemangioma formation, mimicking placental growth and regression. The third hypothesis links tissue hypoxia (e.g., prematurity, low birth weight) to HIF-1 α -mediated angiogenesis via VEGF and bFGF. The fourth involves the renin-angiotensin system, where elevated renin and angiotensin activity stimulate proliferation, while Beta-blockers promote regression by inhibiting renin release. [2] Infantile hemangiomas can be broadly categorized as localized, segmental, or multifocal lesions based on their distribution and morphological features, such as superficial, deep, and mixed types, or reticular/abortive/minimal growth types. [3]

Superficial IHs are located in the epidermis and dermis with little or no subcutaneous involvement, appearing bright red and previously termed “strawberry hemangiomas,” while deep IHs lie beneath the skin surface, are more diffuse and ill-defined, and present with normal skin color or a bluish hue, formerly called “cavernous hemangiomas.” Mixed IHs include both superficial and deep components, and deep or mixed types typically involve deeper soft tissues and become noticeable around 1–2 months of age or later. [2] Although the majority of IH manifest as single, localized cutaneous lesions, bilateral or multifocal presentations are uncommon and are rarely reported in the literature, so it is crucial to record and examine such cases. IH can develop anywhere on the body, but they most commonly occur in the head and neck region (60%), followed by the trunk (25%) and the extremities (15%), reflecting their well-documented predilection for the head and neck area [4] Hemangioma of the genitalia is extremely uncommon. Genital hemangioma is self-limited (resolves spontaneously), therefore conservative treatment is usually suggested. [5] The exact cause of glans hemangioma remains unclear; it is variously considered a benign vascular tumor or congenital vascular anomaly, while other theories suggest it may result from herniation of cavernosal tissue or develop secondary to a prior penile hematoma.

Treatment options for small hemangiomas include surgical excision, cryotherapy and electrofulguration. Recent treatment includes sclerotherapy and laser fulguration. Surgical excision of penile hemangioma has an increased risk of bleeding during the excision because of rich blood supply of penis but nocturnal erections may occur post-operatively. Laser treatment with Nd:YAG in hemangioma of glans penis has also been successfully suggested. [6] The majority of lesions are asymptomatic and disappear on their own, but a small percentage can cause ulceration, bleeding, infection, associated structural anomalies, disfigurement and depending on the anatomical location, functional impairment like airway or vision obstruction. [1]

Diagnosis of infantile hemangiomas is uncertain and needs differentiation from other vascular malformations. Imaging modalities such as ultrasonography with Doppler, magnetic resonance angiography (MRA) or magnetic resonance imaging (MRI) may be useful for the diagnosis of IH and monitor their response to treatment. Regarding its treatment, timolol maleate, a topical nonselective beta-blocker, is used at a dose of 1–2 drops of 0.5% gel-forming ophthalmic solution, applied twice daily for the treatment of small, thin and superficial Hemangiomas. Oral standard treatment strategies include corticosteroids and oral propranolol (non-selective beta-adrenergic antagonist). Oral propranolol is the gold standard treatment for high-risk IHs at a dosage of 2–3 mg/kg/day in 2 daily divided doses for at least 6 months. [2]

Propranolol causes early color change in hemangiomas (pink to violaceous) due to vasoconstriction, reduces growth by downregulating proangiogenic factors (e.g., MMP-2, MMP-9, bFGF, VEGF), and promotes endothelial cell apoptosis, leading to lesion regression. Atenolol is a selective beta-1 antagonist which can also be used in a dose of 1 mg/kg/day and has a decreased risk of bronchospasm and hypoglycemia. Oral steroids were previously used for treatment of hemangiomas in a dose of 2–3 mg/kg/day for 9 to 12 months but their use was gradually reduced since beta blockers introduced. Combined treatment of propranolol and corticosteroids can be used in complicated cases of hemangiomas. Intra-lesional steroid injections can be helpful for small and initial proliferating stage of IHs. [7] Pulse dye laser therapy (PDL) or surgery is recommended for the treatment of residual skin changes after IH regression but may be used earlier to treat selected cases of IHs. [2]

The natural history and outcomes of IH highlight the significance of early detection: although many hemangiomas regress with passage of time, complex or high-risk lesions can cause serious morbidity or irreversible cosmetic changes if left untreated. [8] This case of bilateral infantile hemangioma advances knowledge of the clinical range of this prevalent pediatric vascular tumor because bilateral involvement is very uncommon and may have consequences for treatment and results.

Case Presentation:-

CASE 1:-

After taking verbal consent from patient's attendant, we presented a case of a 3-month-old female infant who came to the skin outpatient department at Naimat Begum Hospital/Hamdard University Hospital Karachi, with a rapidly enlarging lesion over the right deltoid region. She was born at full term with a birth weight of 2.3 kg via spontaneous vaginal delivery. The lesion was first noted at approximately 1 month of age and showed progressive enlargement. Prenatal, natal, and post-natal history was not significant. Vaccination history was up-to-date.

On gross examination, the child looked healthy and comfortable. Skin examination showed an oval, erythematous, edematous and palpable boggy mass, 10 × 7 cm in size, present on the right upper deltoid region. (Figure 1A) It was soft, normothermic, non-tender and compressible, with no evidence of ulceration or spontaneous bleeding while rest of the skin, hairs and nails were normal. There was normal muscle structure and function and no muscle atrophy/dystrophy or joint deformity was noted in the right limb. Pulse was normal, sensation and joint reflexes were intact in the involved limb. No pressure symptoms were noted. Rest of the systemic examination like CVS, CNS, respiratory, abdominal, urogenital and locomotory was unremarkable.

The differential diagnoses considered included infantile fibrosarcoma, non-involuting congenital hemangioma (NICH), kaposiform hemangioendothelioma, tufted angioma, and pyogenic granuloma, all of which can present as vascular or soft-tissue lesions in infancy. However, the postnatal onset, rapid growth during early infancy, and classic clinical morphology favored a diagnosis of infantile hemangioma. No imaging studies were performed due to the typical clinical presentation and absence of red-flag features. Topical 2% propranolol therapy was initiated on the day of presentation. Subsequently, oral corticosteroids in a dose of 0.5 mg/kg body weight were added to enhance regression. At follow-up visits till age of 1 year, marked regression in lesion size, thickness with few areas of atrophy, and color changes (from erythematous to violaceous) was observed. (Figure 1B) The treatment was well tolerated, with no adverse events noted. Verbal informed parental consent was obtained, and serial clinical photographs documenting pre- and post-treatment changes (over one year of age) were recorded.

Case 2:-

After taking verbal consent from the patient's attendant, we presented a case of a 3-month-old male infant who came for the first time to the skin OPD at Naimat Begum Hospital/Hamdard University Hospital Karachi, with bilateral symmetrical erythematous lesions involving both wrists and the penile region. According to mother, child was hospitalized just after birth due to chest congestion for one week, and the lesions have developed at the site of cannulization on both wrists. He was born full term with a birth weight of 2.8 kg. The lesions were first noticed at 2 months of age and progressively increased in size. There was no history of pain/difficulty in joint movement of both upper limbs or during micturition or defecation. Prenatal, natal, and post-natal history was not significant. Vaccination history is up-to-date. On gross examination, the child looked healthy and comfortable. Skin examination showed multiple well-demarcated, erythematous soft tissue swellings, present over both dorsal surface of wrists and the base of penis. (Figure 2) It was soft, normothermic, non-tender, and compressible while rest of the skin, hairs and nails were normal. There was normal muscle structure and function and no muscle atrophy/dystrophy

or joint deformity was noted in both upper limbs and wrist. Pulse was normal, no radio-radial or radio-femoral delay, sensation and joint reflexes were intact in both upper limbs and wrist. No pressure symptoms were noted. There was no ulceration, telangiectasia, significant bleeding, or associated anorectal or urogenital anomaly on clinical assessment. Rest of the systemic examination like CVS, CNS, respiratory, abdominal, urogenital and locomotory systems were unremarkable.

Topical 2% propranolol and topical steroid were initiated on the day of presentation. The patient was advised regular follow-up for clinical monitoring but patient did not return for follow-up, and therefore treatment response and long-term outcomes could not be assessed. Verbal informed parental consent was obtained at initial presentation, and clinical photographs were taken for documentation.

Characteristics	Case 1	Case 2
Age at presentation	3 months	3 months
Sex	Female	Male
Birth weight	2.3 kg	2.8 kg
Lesion site	Deltoid region	Bilateral wrists + genital region
Lesion size	Single (10 × 7 cm)	Multiple superficial
Differential diagnosis	Ruled out clinically	IH diagnosed clinically
Treatment	Topical 2% propranolol Topical steroid Oral steroids	Topical 2% propranolol Topical steroid
Outcome	Regression observed	Lost to follow-up

Table 1: summary table of case 1 and case 2




Figure 1: Case 1 (A) oval lesion, 10 × 7 cm in size, erythematous in color, edematous, palpable boggy mass, presented on right upper deltoid region (B) change in color from erythematous to violaceous with few areas of atrophy and decrease in thickness of the lesion on follow-up.



Figure 2: Case 2 (A, B) Multiple well-demarcated, erythematous soft tissue swellings, present over the dorsum of the hand, wrist and left thumb (C) Multiple well-demarcated, erythematous soft tissue swelling over the dorsal aspect of medial side of right wrist and radial aspect of right forearm (D) well-demarcated, erythematous soft tissue swellings present at the base of the penis.

Discussion:-

Infantile Hemangiomas (IH) are hamartomatous, benign vascular tumors, arising from endothelial proliferation forming multiple blood-filled cavities. Infantile Hemangioma can be classified into superficial cutaneous IH, with papillary and reticular dermis involvement and deep cutaneous IH, with reticular dermis and subcutaneous tissue infiltration. Mixed IHs include both superficial and deep components, and deep or mixed types typically involve deeper soft tissues and become noticeable around 1–2 months of age or later [2], the onset of infantile hemangiomas at two months of age in both cases aligns with the existing literature. [9] Among all soft-tissue tumors, Hemangiomas contribute to 7%, whereas intramuscular hemangiomas are comprised of 0.8% of all hemangiomas. [10] The most frequently affected location of these lesions are the head and neck followed by the trunk. [4] Limbs and perineum are among the least commonly involved sites, however, involvement of these locations (limbs, wrist and genitalia) was observed in both cases, which is very rare and contradicts previously reported findings in literature. Of these lesions, 45% are found in lower extremity, [10] however, case 1 presented with upper extremity lesion, which is discordant with existing literature. Even though our patient (case 2) was a male, hemangiomas are more frequently presented in females with female to male ratio being 3–5:1, which correlates with case 1 but not consistent with case 2. Diagnosis of IH is clinical, however when diagnosis is uncertain, imaging modalities like ultrasonography and MRI can be used. Imaging typically demonstrates a solid, well-circumscribed mass with high-flow vascularity on Doppler ultrasonography. On MRI, it usually appears as a well-defined mass that is isointense to muscle on T1, hyperintense on T2, with flow voids. [2] The diagnosis was made clinically in our patients, and imaging studies were not performed because of the non-affordability and loss to follow-up of the patient in case 2.

The patient in case 1 presented with an IH over the deltoid region. Although IH involving limbs is not entirely uncommon, this particular presenting location is atypical especially with concomitant swelling of the surrounding area. It is crucial to differentiate infantile hemangioma from an infantile fibrosarcoma. Although similar in

appearance, key exam findings like firmness, spherical/protruding and grotesque appearance can point towards an alternate diagnosis of infantile fibrosarcoma, [10] which was incongruent with our patient's clinical findings. Another differential diagnosis is a NICH (Non-involuting congenital hemangioma) which grows in utero and is present at birth [8] and can be differentiated from IH on the basis of a careful history and examination as well. Other possible differentials include infection, trauma, kaposiform hemangioendothelioma, tufted angioma and pyogenic granuloma. [11] Approximately 27% of intramuscular hemangiomas occur in the upper extremity, most commonly involving the triceps, biceps, FDS, and FDP muscles; however, in our case, the lesion is located in the deltoid muscle, which is atypical compared to the usual distribution. Only one case report by Shah et al has been published in Pakistan, of a 40-year-old male presenting with an upper extremity intramuscular hemangioma in the triceps muscle. [12] No literature has been identified reporting upper limb IH over the deltoid region in Pakistan as well as internationally.

The patient in case 2 presented with bilateral symmetrical superficial IH involving both wrists without restriction of wrist and thumb movements along with the dorsal aspect of left hand and base of the penis, which is an extremely rare anatomical location for IH. There have been no reported local or international cases of bilateral symmetrical IH of the limbs, with very few cases of hemangiomas involving the wrist area, and only one reported case of isolated deep cutaneous hemangioma of the wrist. [13] The differential diagnosis of wrist hemangiomas includes lipomas, ganglion cysts, vascular malformations, glomus tumors, and other soft tissue tumors. Soft tissue masses of the wrist account for approximately 97% of cases, with ganglion cysts being the most common (54.5%), followed by hemangiomas (9.9%). [14] This report describes a rare case of bilateral wrist hemangiomas in a male child, highlighting the diagnostic challenges associated with atypical presentations. It underscores the importance of including hemangiomas in the differential diagnosis of wrist swellings, particularly when clinical features are unusual or atypical. Accurate diagnosis relies on thorough clinical evaluation and imaging; however, imaging could not be performed in this case due to loss to follow-up.

Penile hemangiomas is a vascular tumor, benign in nature, characterized by painful red to brown lump/nodule, and account for only about 1% of all hemangioma cases, [14] and to date, just 28 cases of penile epithelioid hemangiomas have been reported in the literature, which may be associated with anorectal, urogenital and spinal anomalies but no wrist joint, finger joint and genital abnormality was noted in our patient in case 2. [15,16] Penile epithelioid hemangiomas should be differentiated from several conditions, including Peyronie's disease, epithelioid hemangioendothelioma, epithelioid hemangiosarcoma, Kimura's disease, bacillary angiomatosis, and penile carcinoma. Epithelioid hemangioendothelioma is a borderline malignant tumor with metastatic potential, while epithelioid hemangiosarcoma is an aggressive malignant vascular neoplasm. Bacillary angiomatosis, caused by *Bartonella henselae*, typically occurs in immunocompromised patients. Kimura's disease presents with subcutaneous nodules, lymphadenopathy, and eosinophilia. Penile cancer is a painless ulcerative or cauliflower lesion, therefore clinical and radiological distinction is important. [15,17] Hemangioma of the genitalia is very rare, as found in our patient and usually spontaneously regresses so conservative treatment is generally advised.

Hemangioma of the genital/perineal region is associated with PELVIS syndrome (perineal hemangioma, external genitalia malformations, lipomyelomeningocele, vesicorenal abnormalities, imperforate anus, and skin tag). No such abnormalities were observed in case 2. On immunohistochemical staining, epithelioid hemangiomas are positive for CD31, CD34 and factor VIII-related antigen and negative for keratin and epithelial membrane antigens. [15] The involvement of the genital area is rare and particularly critical to recognize because of the risk of infection and ulceration, requiring early intervention with propranolol. [16,18] The combination of symmetrical IH and genital involvement has not been described well in previous published literature, as presented in our patient (Case 2), since IHs often present as solitary/segmental lesions. A case of multiple hemangiomas of the scrotum, perineum and pelvis has been reported by Nouira Y et. al. [19] In contrast, our case demonstrates a hemangioma at the base of the penis, which is an extremely unusual presentation. Abdominal ultrasonography is advised for infants <6 months with ≥ 5 cutaneous hemangiomas to rule out hepatic hemangiomas. [20] However, neither patient fit the criteria for screening for hepatic hemangioma, hence abdominal ultrasonography was not performed in our patients.

The hemangioma in patient 1 regressed with treatment with combination of oral steroids and topical propranolol which is consistent with known response to combination therapy for IH. [21] Patient 2 was also prescribed the same treatment however no treatment outcome data is available for comparison due being lost to follow up. The risk factors for developing IH include female sex, low birth weight, progesterone intake, miscarriage history, anemia in pregnancy, PPRM (preterm premature rupture of membranes), placenta previa, PROM (premature rupture of

membranes) and abnormal amniotic fluid volume. [22,23] There were risk factors like female gender and low birth weight (2.3 kg weight at time of birth) present in patient 1, with no significant risk factors in patients 2 but unusual presentation in male as IH are more common in females as compare to males and no literature support for this presentation.

IH is associated with serum markers like GLUT-1, IGF-2, VEGF-A with a positive correlation between disease severity and serum marker levels. [24] There are a few suggested hypotheses for the pathogenesis of IH. Hypoxia induced expression of hypoxia-inducible factor-2 alpha (HIF-2 alpha) and suppression of aldehyde dehydrogenase 1 (ALDH1A1), migration of placental origin IH stem cells supported by presence of circulating levels of Chromosome 19 miRNA cluster (C19MC) microRNA, as well as elevated levels of ACE (angiotensin converting enzyme) and AGTR1 (angiotensin II receptor type 1) are all pathogenic mechanisms hypothesized for this disease. [25] The bilaterality of the IH presenting in patient 2 suggests developmental pathology, which is consistent with current literature. However, we believe this presentation could possibly act as a phenotypic timestamp, suggesting that the pathological mechanism occurred early in development, before progenitor cells are spatially restricted. Solitary lesions, by contrast, likely result from later or localized alterations in endothelial progenitors.

Conclusion:-

Infantile hemangiomas can present with atypical patterns and locations that expand the spectrum of known disease manifestations. Recognition of such uncommon manifestations is essential to avoid misdiagnosis, exclude important mimics, anticipate common complications and initiate timely treatment, particularly in anatomical sites with high risk of complications. These observations support a possible developmental basis for symmetric lesions and underscore the need for further reporting and study of rare infantile hemangioma phenotypic presentations.

Acknowledgements:-

I acknowledge all the work done by all authors.

Conflict of Interest:-

There is no conflict of interest among all authors.

Funding Source:-

None

Ethical Considerations:-

The research adhered to the ethical framework. Participant was comprehensively informed about the study's objective and the researcher's role, ensuring transparency. IRB approval was obtained and informed consent was obtained from the patient's attendants. Anonymity and confidentiality were upheld.

Abbreviations:-

IH = Infantile hemangioma
GLUT-1 = Glucose Transporter 1
VEGFR = Vascular Endothelial Growth Factor Receptors
VEGF-A = Vascular Endothelial Growth Factor A
IGF-2 = Insulin-like Growth Factor 2
HIF-1 α = Hypoxia-inducible factor 1-alpha (HIF-1 α)
bFGF = Basic fibroblast growth factor
CVS = Cardiovascular System
CNS = Central Nervous System
Nd:YAG = Neodymium-doped Yttrium Aluminum Garnet
MMP = Matrix Metalloproteinase
FDS = Flexor Digitorum Superficialis
FDP = Flexor Digitorum Profundus

References:-

1. Sandru F, Petca A, Radu AM, Preda AG, Turenschi A, Constantin AT, Miulescu RG. Infantile Hemangioma: Risk Factors and Management in a Preterm Patient-A Case Report. Reports (MDPI). 2024 Jan 1;7(1):3. doi: 10.3390/reports7010003.
2. Jung HL. Update on infantile hemangioma. Clin Exp Pediatr. 2021 Nov;64(11):559-572. doi: 10.3345/cep.2020.02061.
3. Jaiswal P, Singh J, Dua A, Sirohi A, Ashraf F. Unveiling Infantile Hemangiomas: A Comprehensive Study of Patterns, Presentations, and Complications. Dermatol Pract Concept. 2025;15(4):5515. doi:10.582b/dpc.1504a5515
4. Sun Y, Zhao J, Meng Y, Luo X, Jiang C, Deng G, Lei S. The prevalence, complications, and risk factors for infantile hemangioma: a systematic review and meta-analysis. Int J Dermatol. 2024 Jun;63(b):737-74b. doi: 10.1111/ijd.170b2.
5. Alter GJ, Trengove-Jones G, Horton CE Jr. Hemangioma of penis and scrotum. Urology. 1993 Aug;42(2):205-8. doi: 10.1016/0090-4295(93)90649-u.
6. Aydur E, Erol B, Tahmaz L, Irkilata HC, Eken C, Peker AF. Coagulation of a giant hemangioma in glans penis with holmium laser. Asian J Androl. 2008 Sep;10(5):819-21. doi: 10.1111/j.1745-7262.2008.00359.x. Epub 2007 Dec 20.
7. Merlino L, Volpicelli AI, Anglana F, D'Ovidio G, Dominoni M, Pasquali MF, Gardella B, Inghirami P, Lippa P, Senatori R. Pediatric Hemangiomas in the Female Genital Tract: A Literature Review. Diseases. 2024 Feb 28;12(3):48. doi: 10.3390/diseases12030048.
8. Leung AKC, Lam JM, Leong KF, Hon KL. Infantile Hemangioma: An Updated Review. Curr Pediatr Rev. 2021;17(1):55-b9. doi:10.2174/1573396316666200508100038.
9. Holm A, Mulliken JB, Bischoff J. Infantile hemangioma: the common and enigmatic vascular tumor. J Clin Invest. 2024 Apr 15;134(8):e172836. doi: 10.1172/JCI172836.
10. Hafedh Daly, Faiez Boughanmi, Mohamed Zayati, Leith Limayem, Mohamed Ali Chaouch, Fethi Jebali, Infantile fibrosarcoma of the left upper limb mimicking a hemangioma: A case report, International Journal of Surgery Case Reports, Volume 131, 2025, 111417, ISSN 2210-2612, <https://doi.org/10.1016/j.ijscr.2025.111417>.
11. Elluru RG. Cutaneous vascular lesions. Facial Plast Surg Clin North Am. 2013 Feb;21(1):111-26. doi: 10.1016/j.fsc.2012.11.001.
12. Shah SA, Hassan F. Intramuscular hemangioma of the triceps muscle. Journal of Surgery Pakistan (International). 2012 Jan;17(1):40-2.
13. Ranjan R, Kumar R, Jeyaraman M, Biswas S, Kumar A, Agarwal G. Hemangioma of First Extensor Compartment of the Wrist- A Rare Case Report. J Orthop Case Rep. 2021 Jun;11(6):14-18. doi: 10.13107/jocr.2021.v11.i06.2238.
14. Zahid rasheed, Nayab Hafeez, Iqra Iftikhar , et al. Swelling to Surgery: Hemangioma on the dorsal aspect of wrist joint causing impaired movements in a Young Adult ; A Rare Case Report. Authorea. January 30, 2026. DOI: 10.22541/au.176976927.78044144/v1
15. Rajender, Archana; Phillips, Elizabeth; Munarriz, Ricardo. Penile Epithelioid Hemangioma: A Case Report. Journal of Integrative Nephrology and Andrology 2(1):p 38-40, Jan-Mar 2015. | DOI: 10.4103/2225-1243.150011
16. Tran C, Tamburro J, Rhee A, Golden A. Propranolol for Treatment of Genital Infantile Hemangioma. J Urol. 2016 Mar;195(3):731-7. doi: 10.1016/j.juro.2015.09.069.
17. Ismail M, Damato S, Freeman A, Nigam R. Epithelioid hemangioma of the penis: case report and review of literature. J Med Case Rep. 2011 Jun 30;5:260. doi: 10.1186/1752-1947-5-260.
18. Hwang HW, Lee SB, Shin J, Choi GS, Byun JW. A Case of Perineal Hemangioma, External Genitalia Malformations, Lipomyelomeningocele, Vesicorenal Abnormalities, Imperforate Anus, and Skin Tag (PELVIS) Syndrome with Extensive Perineal Infantile Hemangioma. Ann Dermatol. 2023 May;35(Suppl 1):S88-S92. doi: 10.5021/ad.21.183.
19. Nouira Y, Kbaier I, Attyaoui F, Menif E, Horchani A. Megapenis associated to corpus spongiosum agenesis with scrotal and pelvic hemangiomas. Eur Urol. 2001 Nov;40(5):571-4. doi: 10.1159/000049838.
20. Demie S, Bossak M. Classical Findings of Infantile Hepatic Hemangiomas. HCA Healthc J Med. 2021 Aug 29;2(4):267-272. doi: 10.36518/2689-0216.1250.
21. Aly MM, Hamza AF, Abdel Kader HM, Saafan HA, Ghazy MS, Ragab IA. Therapeutic superiority of combined propranolol with short steroids course over propranolol monotherapy in infantile hemangioma. Eur J Pediatr. 2015 Nov;174(11):1503-9. doi: 10.1007/s00431-015-2561-1.

22. Chehad AS, Hamza O, Mansoul T. Clinical and epidemiological risk factors for infantile hemangioma: A case-control study. *Pediatr Dermatol.* 2023 Jul-Aug;40(4):647-650. doi: 10.1111/pde.15363.
23. Gong X, Qiu T, Feng L, Yang K, Dai S, Zhou J, Zhang X, Chen S, Ji Y. Maternal and Perinatal Risk Factors for Infantile Hemangioma: A Matched Case-Control Study with a Large Sample Size. *Dermatol Ther (Heidelb).* 2022 Jul;12(7):1659-1670. doi: 10.1007/s13555-022-00756-4.
24. Aydın Köker S, Kömüroğlu AU, Köksoy AY, Şiraz ÜG, Tekin E, Köker A. Evaluation of GLUT1, IGF-2, VEGF, FGF 1, and angiopoietin 2 in infantile hemangioma. *Arch Pediatr.* 2021 May;28(4):296-300. doi: 10.1016/j.arcped.2021.02.009
25. Kowalska M, Dębek W, Matuszczak E. Infantile Hemangiomas: An Update on Pathogenesis and Treatment. *J Clin Med.* 2021 Oct 9;10(20):4631. doi: 10.3390/jcm10204631.