

 <p>ISSN (O): 2320-5407 ISSN (P): 3107-4928</p>	<p>Journal Homepage: - www.journalijar.com</p> <h2>INTERNATIONAL JOURNAL OF ADVANCED RESEARCH (IJAR)</h2> <p>Article DOI: 10.21474/IJAR01/23210 DOI URL: http://dx.doi.org/10.21474/IJAR01/23210</p>	
--	---	---

RESEARCH ARTICLE

EFFECTS OF ARBUSCULAR MYCORRHIZAL FUNGAL STRAINS ON THE GROWTH OF COWPEA [VIGNA UNGUICULATA (L.) WALP.] IN THE PRESENCE OF TILEMSI NATURAL PHOSPHATE (TNP)

Souleymane Kone¹, Souleymane Bathie Kone¹, Rokiadou Fane¹ and Vital Traore²

1. Laboratory for Research in Microbiology and Microbial Biotechnology (Labo REM-Biotech) of the Faculty of Sciences and Techniques (FST) of the University of Sciences, Techniques and Technologies of Bamako (USTT-B), BP E 3206, Mali.
2. Laboratory of Analytical Chemistry, Department of Chemistry, Faculty of Sciences and Techniques (FST) of the University of Sciences, Techniques and Technologies of Bamako (USTT-B), BP E 3206, Mali.

Manuscript Info

Manuscript History

Received: 8 February 2026
Final Accepted: 10 March 2026
Published: April 2026

Key words:-

Mycorrhization, Fertilization, Cowpea.

Abstract

The growth of cowpea [*Vigna unguiculata* (L.) Walp.] was evaluated in response to the inoculation of seeds at sowing with an arbuscular mycorrhizal fungus strain (*Glomus aggregatum*) in the presence of Tilemsi natural phosphate (TNP). The soil collected from the large cultivation plot of IPR/IFRA in Katibougou served as the substrate after sterilization. Sowing was done with 3 cowpea seeds per pot, and after one week, two plantlets were kept in each pot. The pots were placed on metallic tables in the greenhouse of the Soil Microbiology Unit (LMS) of LaboREM Biotech for the two months of the experiment. The results showed that, despite a low rate of root colonization by the mycorrhizal fungi (F: 80.75%; I: 7.31), the average growth of inoculated plants was more significant compared to non-inoculated plants. The results also indicate that the fungus + natural phosphate treatment significantly increased the average number of inflorescences, aerial, and root biomasses. The treatment with the fungus alone was more significant in terms of the frequency and intensity of mycorrhization. However, as biomass predicts yield, this study suggests that the mycorrhizal biofertilizer provides important benefits to the plant by solubilizing the TNP. Mycorrhizal fungi can thus be used to improve the growth and yield of cowpea in several agricultural perimeters after TNP application.

"© 2026 by the Author(s). Published by IJAR under CC BY 4.0. Unrestricted use allowed with credit to the author."

Introduction:-

The constantly increasing African population forces farmers to make lands more and more productive. Sub-Saharan Africa alone accounts for nearly 200 million people (FAO, 2010). This results in undernourishment combined with limited and degraded land resources (Le Cacheux, 2011; Fitter, 2012).

Corresponding Author:-Souleymane Kone

Address:-Laboratory for Research in Microbiology and Microbial Biotechnology (LaboREM-Biotech) of the Faculty of Sciences and Techniques (FST) of the University of Sciences, Techniques and Technologies of Bamako (USTT-B), BP E 3206, Mali.

In reality, agricultural production on soils poor in nutrients and deficient in water yields very low returns. Currently, the high productivity of intensive agriculture is greatly dependent on chemical inputs, particularly mineral phosphate fertilizers, which enrich the soils with the necessary nutrients for plant growth (Childers et al., 2011; Cordell et al., 2009; Gilbert, 2009; Roy-Bolduc and Hijri, 2010). Agriculture's dependence on these chemical inputs leads to negative environmental impacts. To avoid a food crisis and achieve sustainable growth, more appropriate and environmentally friendly high-yield technologies have been recommended (Bationo et al., 1998; Scherr and McNeely, 2008; Brussaard et al., 2010). These fall within the framework of efforts undertaken by researchers and extension workers to improve global food security. Cowpea (*Vigna unguiculata* (L) Walp) is one of the main legumes cultivated worldwide. Its cultivation area is located between latitude 35°N et 30°S of the equator (Singh et al., 1997). It is an annual plant with an upright or creeping habit. The leaves, as well as the green pods and seeds of cowpea, are a good source of protein, vitamins, and minerals for human and animal consumption (Cisseand Hall, 2003). Furthermore, during the dry season, in some regions of West and Central Africa, the monetary value of stored cowpea haulms becomes very high (Cisseand Hall, 2003). Its cultivation, like that of other legumes, can help restore soil fertility thanks to its high potential for biological nitrogen fixation (Bado, 2002; Graham and Vance, 2003).

Despite the numerous studies conducted on this species, the average yield per hectare of cowpea cultivation remains relatively unstable. It varies between 50 to 550 kg/ha in Africa and depends on climatic conditions, varieties, cropping systems, and the level of use of inputs and pesticides (Cisseand Hall, 2003). The low yields observed are due, among other things, to low rainfall, soil deficiency in nitrogen and phosphorus, which are essential for the plant's nutrition and production, as well as low soil fertility and extensive agriculture without soil fertility restoration (FAO, 2003 ; Bationo et al., 1998). Mineral phosphate fertilizers are very expensive and inaccessible to farmers because of their low purchasing power. However, several African countries have significant deposits of natural phosphates (NP), a potential, cheaper source of phosphorus for farmers, which could reduce dependence on chemical fertilizers, reduce pollution risks, and thus enhance local fertilizing resources such as Tilemsi natural phosphates in Mali, Matam in Senegal, and Kodjari in Burkina Faso (Koneet al. 2021, 2019, 2018 ; Babana and Antoun, 2006, 2005 ; Dianou and Bâ, 1999 ; Asimi and Kambou, 2000).

Nevertheless, the utilization of natural phosphates (NP) by plants is not always easy because of their low solubility (Bagayoko and Coulibaly, 1995). To improve NP solubility, several strategies are used, including inoculation with arbuscular mycorrhizal fungi (AMF) (Strullu, 1991 ; Babana and Antoun, 2003). These microorganisms, naturally present in soils, form symbioses with the majority of terrestrial plants, including cultivated ones (Smith and Read, 2008). They play an important role in water and mineral nutrition (Smith and Read, 2008), protecting host plants against pathogens (Sikes et al., 2009) and against drought (Augeet al., 2001). In addition, they improve ecosystem sustainability by enhancing soil structure (Wilson et al., 2009) and reducing nutrient losses that can be caused by runoff (Van der Heijden, 2010). This study is part of the effort to improve cowpea growth. For this reason, the use of a selected arbuscular mycorrhizal fungus strain, *Glomus aggregatum*, as a biological fertilizer, coupled with mineral fertilization based on Tilemsi natural phosphate, could ensure sustainable management of this soil's fertility and increase the growth of cowpea, an important plant in human diet.

Materials and Methods :-

Study Site and Soil Sampling :-

The study was conducted at the Soil Microbiology Unit (LMS) of LaboREM-Biotech at the Faculty of Sciences and Techniques (FST) of the University of Sciences, Techniques and Technologies of Bamako (USTT-B). The soil was sampled from the 0-30 cm horizon in Katibougou (Koulikoro/Mali), 60 km from Bamako. After sterilization in an autoclave for 20 minutes at 120° C (2 times), it was distributed into 5 kg plastic pots with holes in the bottom, at a rate of 3.75 kg per pot. The characteristics of this soil are mentioned in Table 1.

Table 1: Some physicochemical characteristics of the soil used.

	Clay (%)	Sand (%)	C (%)	N (%)	C/N	pH	Ca (mg)	Mg (mg)	P (ppm)	K (mg)
Value	10.4	57.6	0.2	0.03	3.9	5.8	10	1.8	2.48	9.3

Source: SACKO/AUF 2014 project data

Plant Material :-

The cowpea variety used belongs to the collection of the Rural Economy Institute (IER) of Mali. It is KOROBALLEN, a short-cycle variety (65 to 70 days) cultivated in the 300 to 800mm isohyet.

Fungal and Fertilization Materials:-

The inoculum used is the *Glomus aggregatum* (AMF) strain. This inoculum comes from the collection of the Soil Microbiology Unit of the Laboratory of Research in Microbiology and Microbial Biotechnology/LaboREM-Biotech. This mycorrhizal fungus strain was isolated from an *Acacia mangium* plantation and has proven effective in plants and crops (Sacko et al., 2012, 2014). It was multiplied on millet in a culture chamber according to the method described by Bâ et al. (1996). The inoculum consists of a mixture of spores, hyphae, millet root fragments colonized by *Glomus aggregatum* (Ga), and sand. At the time of sowing, 20g of this mixture was added per pot, as well as 20g of TNP 30% (P_2O_5). The TNP used comes from Bourem in the Tilemsi valley in Northern Mali. It is used in its powder form at 30% (P_2O_5), 10% of elements smaller than 40 μ m and 60% of elements smaller than 60 μ m. It exhibits a solubility of 0.007% in water.

Culture Conditions:-

In this study, two factors were examined:

- **Inoculation with 2 modalities:**
 - Control (without inoculant).
 - Inoculation with *Glomus aggregatum* (Ga) at a rate of 20g of inoculum per pot.
- **Fertilization with TNP at 2 variations levels:**
 - Control (without fertilizer).
 - Fertilization with PNT at 20g per pot.

The experimental design was a Fischer block with four treatments of 10 repetitions. Three cowpea seeds were sown directly per pot. Thinning occurred on the 10th day after sowing, leaving 2 plants per pot. Watering was done according to demand, and the experiment lasted 2 months.

Measured Parameters :-

To evaluate biomasses and mycorrhization parameters, a sampling of five homogeneous plants per treatment was conducted on the 60th day after sowing. The plants were uprooted, and the aerial and root parts of each plant were separated. They were then placed in labeled envelopes, dried in an oven at 70°C for 72 hours, and then weighed. The roots of the plants from each treatment were stained according to the method described by Phillips and Hayman (1970), and the slides were then read using the scale described by Trouvelot et al. (1986).

Statistical Analyses :-

The results were analyzed using ANOVA in R2.101 softwares, and the NEWMAN-KEULS test, at a 5% threshold, was used to compare the means.

Results :-

Influence of Endomycorrhizal Inoculation on Growth Parameters :-

The effect of inoculation on aerial and root biomasses was determined on the 60th day after sowing (DAS). The results obtained show that all inoculated plants allowed an increase in biomasses on the 60th DAS compared to the controls. The results of the variance analysis show significant effects of the *Glomus aggregatum* x Tilemsi Natural Phosphate interaction on the production of cowpea plant biomasses at 2 months (Table 2).

Table 2: Analysis of variance of dry aerial and root biomasses of cowpea plants after 2 months of culture.

Treatments	Dry Aerial Biomass	Dry Root Biomass
Ga + TNP	6.04 a	2.10 a
Ga	2.82 c	1.64 b
TNP	4.00 b	1.90 ab
Control	2.68 c	0.70 c

Legend : Values followed by the same letter are not statistically different from each other at the 5% threshold according to the Tukey Contrasts Test. Ga = *Glomus aggregatum*; TNP = Tilemsi Natural Phosphate.

Mycorrhization Parameters of Cowpea Plants :-

At 60 days of culture, the intensities and frequencies of mycorrhization were determined. For all treatments, the results showed low mycorrhization intensities (less than 10%) with very high frequencies (>70%). The variance analysis between the inoculation and phosphate fertilization factors showed significant effects of the *Glomus aggregatum* x Tilemsi Natural Phosphate interaction on the mycorrhization parameters of cowpea plants (Table 3).

Table 3: Mean values of Mycorrhization Frequencies and Intensities of cowpea plants per pot after 2 months of culture.

Treatments	Mycorrhization Frequency (%)	Mycorrhization Intensity (%)
Ga + TNP	76.0 a	6.68 a
Ga	85.5 b	7.94 b

Discussion:-

This work evaluated the response of cowpea to endomycorrhizal inoculation with *G. aggregatum* in the presence of TNP in the greenhouse. The results show that inoculation with *G. aggregatum* and TNP application significantly improves cowpea biomasses in the greenhouse (+81%). Thus, significant effects of the *Glomus aggregatum* x Tilemsi Natural Phosphate interaction on the biomass production of cowpea plants at 2 months were observed. Similar results have been observed on cowpea under semi-controlled conditions by (Haro et al., 2012) and also on other plant species (Bâ et al., 2001 ; Avio et al., 2006 ; Koneet al. 2021, 2019, 2018). This positive effect of inoculation in the presence of TNP on the growth of cowpea plants is largely due to a better improvement in phosphate nutrition. Indeed, arbuscular mycorrhizal fungi are likely to increase the growth and production of cultivated plants by improving their water and mineral nutrition and protecting them against pathogens (Dommergues et al., 1999; Smith and Read, 2008; Sikes et al., 2009). These results on cowpea are consistent with those of (Smith et al., 2000) and (Avio et al., 2006) on *Medicago sativa*.

They also align with those of (Sampath Kumar et al, 2002; Prakash et al, 2004; Boureima et al, 2007) who observed an improvement in sesame biomass in the greenhouse following mycorrhizal inoculation. They also corroborate those of (Sampath Kumar et al., 2002; Koneet al. 2021, 2019, 2018), which showed that the association with different genera of arbuscular mycorrhizal fungi (*Acaulospora*, *Gigaspora*, *Glomus*, *Sclerocystis*, and *Scutellospora*) allowed for greater biomass at both the root and aerial parts of local varieties of sesame, maize, and sorghum. Our results are identical to those of (Prakash et al., 2004; Koneet al. 2021, 2019, 2018) who obtained good assimilation of natural phosphate and indirectly an intense absorption of nitrogen nutrients with an improvement in the biomass of mycorrhized sesame, sorghum, and maize plants coupled with natural phosphate application in the greenhouse. Indeed, the results allowed for a total biomass in all mycorrhized plants superior to the controls. Despite the lower mycorrhization parameters in inoculated plants, a significant positive effect on cowpea plant growth was observed. Smith et al. (2004) concluded that mycorrhization intensities are not always correlated with plant growth. Similarly, significant effects of the *Glomus aggregatum* x Tilemsi Natural Phosphate interaction on the mycorrhization parameters of cowpea plants were observed.

Indeed, the plant's response to mycorrhization depends on the mycorrhizal fungus species (Plenchette et al., 1982; Krishna and Dart, 1984) but also on the host plant species (Plenchette et al., 1983; Krishna et al., 1985). Our results are consistent with those of (Subba Rao et al., 1985; Plenchette et al., 2000; Koneet al. 2018, 2019, 2021), who observed very significant growth stimulations on millet, sorghum, and maize. Also (Sampath Kumar et al., 2002; Koneet al. 2021, 2019, 2018) obtained an infectivity of (97%) with the *Glomus* genus on local varieties of sesame, maize, and sorghum. Inasmuch as the symbiosis improves water and mineral nutrition, the plant will not find it necessary to form this symbiosis if nutrients are available in the environment. Thus, according to (Gianinazzi-Pearson and Gianinazzi, 1986; Plenchette, 1982; Strullu, 1991), this symbiosis will gradually establish and develop with the depletion of nutrients directly accessible to the plant roots in the soil. Corroborating our experimentation, (Dianou and Bâ, 1999; Asimi and Kambou, 2000) with Burkina phosphate, as well as (Babana and Antoun, 2006, 2005; Koneet al. 2021, 2019, 2018) with Tilemsi natural phosphate, observed a positive influence of these natural phosphates associated with mycorrhizal fungi on the growth of cowpea, wheat, maize, and sorghum.

Conclusion:-

This work allowed for the study of the greenhouse effect of inoculating a Malian variety of *Vigna unguiculata* with a fungal strain *G. aggregatum* in the presence of Tilemsi natural phosphate. Inoculation with the endomycorrhizal fungus, *Glomus aggregatum*, improves the growth of cowpea cultivated on Katibougou soil thanks to better phosphate nutrition. This suggests a great ability of the introduced symbiont. Hence, this biofertilizer could be recommended to farmers to improve the phosphate nutrition of phosphorus-deficient soils. Thus, in tropical countries where soil acidity is a limiting factor for plant growth, the inoculation of plants with mycorrhizal fungus strains appears as an alternative to the use of chemical fertilizers for sustainable agriculture and long-term preservation of soil fertility.

Bibliographical References :-

1. FAO. 2010. The State of Food Insecurity in the World : Addressing Food Insecurity in Protracted Crises. Rome, Italy, 63p.
2. Fitter A. 2012. Why plant science matters ? *New Phytologist* 193 : 1-2.
3. Le Cacheux, J. 2011. World and European agriculture, challenges of the 21st century. *Revue de l'observatoire français des conjonctures économiques*, vol. 120, p. 198-234.
4. Childers, D. L., Corman, J., Edwards, M. and Elser, J. J., 2011. Sustainability Challenges of Phosphorus and Food : Solutions from Closing the Human Phosphorus Cycle. *BioScience*, vol. 61, n°2, p. 117-124.
5. Cordell, D., Drangert, J.-O. and White, S., 2009. The story of phosphorus : Global food security and food thought. *Global Environmental Change*, vol. 19, p. 292-305.
6. Gilbert, N., 2009. The disappearing nutriment. *Nature*, vol. 461, n°8, p. 716-718.
7. Roy-Bolduc, A. and Hijri, M., 2010. The Use of Mycorrhizae to Enhance Phosphorus Uptake : A Way Out the Phosphorus Crisis. *Journal of biofertilizers and biopesticides*, vol. 2, n°1, p. doi : 10.4172/2155-6202.1000104.
8. Bationo et al., 1998; Bationo A, Koala S, Ayuk E. 1998. Soil fertility for cereal production in the Sahelo-Sudanian zone and valorization of natural phosphates. *Cahiers Agricultures* 7 : 365-371.
9. Scherr SJ, McNeely JA. 2008. Biodiversity conservation and agricultural sustainability : towards a new paradigm of 'ecoagriculture' landscapes. *Phil. Trans. R. Soc. B* 363 : 477-494.
10. Brussaard L, Caron P, Campbell B, Lipper L, Mainka S, Rabbinge R, Babin D, Pulleman M. 2010. Reconciling biodiversity conservation and food security : scientific challenges for a new agriculture. *Current Opinion in Environmental Sustainability* 2 : 34-42.
11. Singh BB, Mohan Raj DR, Dashiell KE, Jackai LEN. 1997. Advances in cowpea research. co publication of International Institute of Tropical Agriculture (IITA) and Japan International Research Center for Agricultural Sciences (JIRCAS). IITA, Abadan, Nigeria, 315 p.
12. Cisse N, Hall AE. 2003. Traditional Cowpea in Senegal, case study. www.fao.org/ag/AGP/AGPC/doc/Publicat/cowpea_cisse/cowpea_cisse_e.htm.
13. Bado BV. 2002. Role of legumes on the fertility of tropical ferruginous soils in the Guinean and Sudanian zones of Burkina Faso. Doctoral thesis, Laval University Quebec, 184 p.
14. Graham & Vance, 2003). Graham PH, Vance CP. 2003. Legumes : importance and constraints to greater use. *Plant Physiol* 131(3) : 872-877.
15. FAO. 2003. Soil fertility management for food security in sub-Saharan Africa. Rome, Italy, 55p.
16. Kone S., Kante F., Diop I., Ndoye F., Diedhiou A G., Kane A., Yattara I I., and Sacko O., 2021. Effect of Rock Phosphate and Arbuscular Mycorrhizal Fungi on Maize (*Zea Mays L.*) Growth in Greenhouse Conditions. *European Journal of Scientific Research*. ISSN 1450-216X / 1450-202X Vol. 160 No 3, pp.126 – 134. <http://www.europeanjournalofscientificresearch.com>.
17. Souleymane Koné, Fallaye Kanté, Ibou Diop, Fatou Ndoye, Abdalla Gamby Dieudhiou, Aboubacry Kane, Ousmane Sacko and Inamoud Ibn Yattara, 2019. Effect of Arbuscular Mycorrhizal Fungal Inoculation on Sorghum bicolor Growth at Different Phosphate Levels : A Greenhouse Study. *Journal of Agricultural Science and Technology B* 9, 234-240, doi : 10.17265/2161-6264/2019.04.002.
18. KONE Souleymane, KANTE Fallaye, DIOP Ibou, NDOYE Fatou, DIEUDHIU Abdalla Gamby, KANE Aboubacry, YATTARA Inamoud, SACKO Ousmane, 2018. Effect of fertilization with three strains of arbuscular mycorrhizal fungi and Tilemsi natural phosphate on maize (*zea mays l.*) growth, 10th edition of the MSAS Conference, July 22-27, Bamako. http://msas.ml/bko2018/ACTES_2018/msas2018_n065p129-136.pdf.
19. Babana A.H. et Antoun H. (2006) - Effect of Tilemsi phosphate rock solubilizing microorganisms on phosphorus uptake and yield of field grown wheat in Mali. *Plant and Soil*, 287 : 51-58.

20. Babana A.H. et Antoun H. (2005) - Biological system for improving the availability of Tilemsi phosphate rock for wheat (*Triticum aestivum* L.) cultivated in Mali. 2005. *Nutrient Cycling in Agroecosystems*, 72 : 147-152.
21. Dianou D. et Bâ A.M. (1999) - Influence of three rhizobacteria on the growth and nitrogen fixation of cowpea in the presence of natural phosphate. *Sciences et Techniques, Sciences Naturelles et Agronomie*, 23 (2) : 7-15.
22. Asimi S. et Kambou G. (2000) - Effects of native bacteria on the solubilization of natural phosphate from Burkina and maize growth. *Sciences et Techniques, Sciences Naturelles et Agronomie*, January-June 2000, 24 (2) : 7-16.
23. Bagayoko M. and Coulibaly B.S. (1995). Promotion and evaluation of Tilemsi phosphate rock in Mali agriculture. In H. Gerner & A.U. Mokwunye, eds. *Use of phosphate rock for sustainable agriculture in West Africa*, pp.77-83. *Miscellaneous Fertilizer studies N° 11 Muscle Shoals, USA, IFDC*.
24. Strullu DG. (1991). *Mycorrhizae of Trees and Cultivated Plants*. Edition Lavoisier : Paris, France ; p. 250.
25. Babana A. H., 2003. Effect of Tilemsi phosphate rock-solubilizing microorganisms on phosphorus uptake and yield of field-grown wheat (*Triticum aestivum* L) in Mali. *Plant and soil* :1-8.
26. Smith SE, Read DJ. 2008. *Mycorrhizal symbiosis*, 3rd edn. Academic Press Inc., London, UK.
27. Sikes BA, Cottenie C, Klironomos JN. 2009. Plant and fungal identity determine pathogen protection of plant roots by arbuscular mycorrhizas. *Journal of Ecology* 97 : 274–1280.
28. AugeRM, Stodola AJW, Tims JE, Saxton AM. 2001. Moisture retention properties of a mycorrhizal soil. *Plant and Soil* 230 : 87–97.
29. Wilson GWT, Rice CW, Rillig MC, Springer A, C. HD. 2009. Soil aggregation and carbon sequestration are tightly correlated with the abundance of arbuscular mycorrhizal fungi : results from long-term field experiments. *Ecology Letters* 12 : 452–461.
30. Van der Heijden MGA. 2010. Mycorrhizal fungi reduce nutrient loss from model grassland ecosystems. *Ecology* 91 : 1163–1171.
31. Sacko O., 2014 - Valorization of Tilemsi Natural Phosphate (TNP) by endomycorrhizal Fungi. MSAS Conferences – Bamako, August 03-08, 2014.
32. Sacko O, Yattara I, Messaoud L, Neyra M, 2012. Effects of a rock phosphate on indigenous rhizobia associated with *Sesbania sesban*. *Journal of Environmental Management* 95 (2012) : 265-268.
33. Bâ A. M., BazieM. et Guissou T. (1996) - Effect of natural phosphate on young *Acacia albida* Del. with or without mycorrhizae. In : *Les Parc à Faidherbia, Cahiers Scientifiques du CIRAD-Forêts des Tropiques*, 12 : 237-244.
34. Phillips & Hayman (1970) Phillips J. M. & Hayman D. S., 1970. Improved procedures for clearing roots and staining parasitic and vesicular-arbuscular mycorrhizal fungi for rapid assessment of infection. *Trans. Br. Mycol. Soc.* 5 : 158-161.
35. Trouvelot A., Kouch J. et Gianinazzi-Pearson V., 1986. Measurement of the VA mycorrhization rate of a root system Research for estimation methods having a functional significance, in : Gianinazzi S. (Ed.), *Les mycorrhizes : Physiologie et Génétique*, 1st European Seminar on Mycorrhizae, Dijon, INRA, Paris : 217 – 221.
36. Haro H., Sanon B.K., Diop I., Kane A., Dianda M., Houngnandan P., Neyra Marc, Traore A., 2012. Response to mycorrhizal inoculation of four cowpea varieties [*Vigna unguiculata* (L.) Walp.] cultivated in Burkina Faso and Senegal. *Int. J. Biol. Chem. Sci.* 6(5) : 2097-2112.
37. Bâ AM, Guissou T, Duponnois R, Plenchette C, Sacko O, SidibeD, Sylla K. 2001. Controlled mycorrhization and phosphate fertilization : application to the domestication of the jujube tree. *Fruits* 56 261-269.
38. Avio L, Pellegrino E, Bonari E, Giovannetti M. 2006. Functional diversity of arbuscular mycorrhizal fungal isolates in relation to extraradical mycelial networks. *New Phytologist* 172 : 347–357.
39. Dommergues Y, Duhoux E, Diem HG. 1999. Nitrogen-fixing trees: Fundamental characteristics and role in the management of Mediterranean and tropical ecosystems with particular reference to sub-humid and arid zones. Ed. CIRAD, FAO, IRD, Paris, France, 499p.
40. Smith FA, Jacobsen I, Smith SE. 2000. Spatial differences in acquisition of soil phosphate between two arbuscular mycorrhizal fungi in symbiosis with *Medicago truncatula*. *New Phytologist* 147 : 357-366.
41. Sampath Kumar G, Muruges S, Rajendran A, Madhumathi B, Ganesh Kumar A. 2002. Association of VAM fungi with sesame and its influence on growth. *Sesame and Safflower Newsletter* No. 17.
42. Prakash A, Tandon V, Sharma NC. 2004. Effect of rock phosphate and VAM inoculation on growth and nutrient uptake in *Sesamum indicum* L. *Physiol. Mol. Biol. Plants*, 10(1) : 137-141.
43. Boureima S, Diouf M, Diop TA, Diatta M, Léye EM, Ndiaye F, Seck D. 2007. Effects of arbuscular mycorrhizal inoculation on growth and development of sesame (*Sesamum indicum* L.). *Afr. J. Agric. Res.*, 3(3): 234-238.

44. Smith SE, Smith FA, Jakobsen AI. 2004. Functional diversity in arbuscular mycorrhizal (AM) symbioses: the contribution of the mycorrhizal P uptake pathway is not correlated with mycorrhizal responses in growth or total P uptake. *New Phytologist* 162: 511-524.
45. Plenchette C., Fortin J. A. et Furlan V., 1982 – Effects of different endomycorrhizal fungi on five host plants grown on calcined montmorillonite clay. *J. Amer. Hort Sci.*, 107, pp. 535-538.
46. Krishna. K. R., Dart P. J., 1984 – Effect of mycorrhizal inoculation and soluble phosphorus fertilizer on growth and phosphorus uptake of pearl millet. *Plant Soil*, 81, pp. 247-256.
47. Plenchette C., Fortin J. A. et Furlan V., 1983 – Growth response of several plant species to mycorrhizae in a soil of moderate P – fertility I. Mycorrhizal dependency under field conditions. *Plant and Soil*, 70, pp 199-209.
48. Krishna. K. R., Shetty K. G., Dart F. J. et Andrews D. J., 1985 – Genotype dependent variation in mycorrhizal colonization and response to inoculation of pearl millet. *Plant Soil*, 86 : 113-125.
49. Subba Rao N. S., Tilak K. B. R. et Singh C. S., 1985 – Effect of combined inoculation of *Azospirillum brasilense* and vesicular mycorrhiza on pearl millet. *Plant Soil*, 84, pp. 283-286.
50. Plenchette C., Bois J-F., Duponnois R., Cadet P. (2000). Mycorrhization (*Glomus aggregatum*) of millet (*Pennisetum glaucum*). *Etude et Gestion des sols*, 7, 4, 2000 – special issue – pages 379 to 384.
51. Gianinazzi-Pearson V, Gianinazzi S. 1986. The physiology of improved phosphate nutrition in mycorrhizal plants. In *Les Mycorrhizes, Physiologie et Génétique*. INRA: Paris; 101-109.