

Journal Homepage: www.journalijar.com

INTERNATIONAL JOURNAL OF ADVANCED RESEARCH (IJAR)

Article DOI:10.21474/IJAR01/23149
DOI URL: <http://dx.doi.org/10.21474/IJAR01/23149>



RESEARCH ARTICLE

DIVERSITY AND DISTRIBUTION OF BIRDLIFE ACCORDING TO VEGETATION TYPE AND SEASONS IN TWO DIFFERENT AREAS IN THE EQUATORIAL CLIMATE OF THE CONGO BASIN REGION: THE MOUANKO MANGROVE AND THE EDEA II EVERGREEN FOREST IN CAMEROON (LITTORAL, REGION)

Anthony Gilchrist Kendeg Kendeg¹, Billy Nguembock^{1,2}, Esther Diane Olivia Azang¹, Sali Mahamat³, Agathe Mvondo Abassa¹, François Okali Mvilongo¹, Marcelle Grace Ngo Bitjaga¹, Loic Franc Litila¹, Joseph Ngom Songo¹, Caliel Michel Nyina Mvolo¹, Arthur Jacques Makendi Makendi¹ and Joseph Lebel Tamesse⁴

1. Department of Animal Biology and Physiology, University of Yaoundé I, Cameroon.
2. Department of Biological Sciences, University of Montreal, Canada.
3. Laboratory of Zoology, Department of Biological Sciences of Living Organisms, University of Garoua, P. O. Box 346 Garoua, Cameroon.
4. Higher Teacher Training College, Department of Biological Sciences, University of Yaoundé I, Cameroon.

Manuscript Info

Manuscript History

Received: 12 February 2026
Final Accepted: 15 March 2026
Published: April 2026

Key words:-

birdlife, mangroves, evergreen forest, distribution, diversity

Abstract

In Cameroon, bird studies are increasingly well-known and often quite long-established. In recent years, several studies have been conducted in various areas across the country. To further update knowledge about Cameroonian avifauna, a study was carried out in the evergreen forest of Edea II and the mangrove of Mouanko, in the Sanaga-Maritime department, Littoral Region. The Japanese mist netting method was used to capture birds between December 2022 and December 2024. PAST (v5.2.1), R Studio (v4.1.2), and Excel (2016) software were used to perform statistical analyses on the distribution and diversity of avifauna in these two areas within the department. A total of 643 individuals were captured, belonging to 10 orders, 28 families, 59 genera, and 92 species. Our results showed that passerines were more abundant (83.2%) than non-passerines (16.8%). The most abundant and diverse family was the Pycnonotidae, with 7 genera and 12 species. The calculated diversity indices indicated that the avifauna of the overall study area is quite diverse ($H' = 3.81$), and our results show no dominance of any single species ($1 - \lambda = 0.95$), with an equal distribution of individuals within each species ($J' = 0.83$). It should be noted that the evergreen forest was more abundant and diverse than that of the mangrove ($S=87$ and $H'=3.70$ vs $S=47$ and $H'=3.64$).

"© 2026 by the Author(s). Published by IJAR under CC BY 4.0. Unrestricted use allowed with credit to the author."

Similarly, the avifauna of the evergreen forest was more abundant and diverse in the first year than in the second, while the opposite was true for the mangrove. Furthermore, the analysis of avifauna distribution according to

Corresponding Author:-Anthony Gilchrist Kendeg Kendeg

Address:-Department of Animal Biology and Physiology, University of Yaoundé I, Cameroon.

seasons showed no significant difference ($p=0.14$), whereas vegetation showed a significant difference ($p=0.0019$), indicating that the habitats are not similar.

Introduction:-

Cameroon is located in the Congo Basin region in which the climate is a mixture of equatorial and tropical climate, with the northern part under the tropical climate while the southern part is under the equatorial climate (Suchel, 1988; Nguembock, 2008). Cameroon is also one of the countries in the world which presents a very great diversity of vegetation, at least nine different types of vegetation (equatorial forest, Guinean savannah, savannah forest mosaic, Sudanese savannah, mandara, sahel, floodplain, and mountainous region), and often this vegetation appears more abundant in mountain than in plain regions (Letouzey, 1985; Stuart, 1986; Nguembock, 2008; Nguembock et al., 2020). Only contrary to mountains, the plain is a biogeographical domain without relief and we mainly find in Cameroon the northern lowlands, some parts of the southern Cameroonian, and Adamawa plateau; the Mouanko and Edea II plains belongs to the coastal plains and contrary to the northern lowlands, it is covered in places with rich forest vegetation (Letouzey, 1985; Stuart, 1986; Nguembock et al., 2020).

Birds play a variety of roles: pollination, seed dispersal, regulation of insect populations, biological indicators of habitat health, and more. In tropical regions, these roles are even more pronounced due to species richness and the complexity of ecological interactions (Green et al., 2002; Green and Elmerberg, 2014; Amano et al., 2018; BirdLife International, 2022; Kafando et al., 2023). The spatial distribution of bird species, their reproductive success, and their survival are influenced by the availability of key resources, including food, water, vegetation cover, and weather conditions (Lee et al., 2012; Adhikari et al., 2020). It is also worth noting that the seasonality is an important factor influencing avian diversity and the availability of key resources (Katuwal et al., 2016).

Cameroon has a great diversity of birds with 974 species distributed among 27 orders and 105 families (Lepage, 2024). Furthermore, in Cameroon, it is documented that there are also 11 endemic bird species (*Ploceus bannermani* Chapin, 1932; *Tauraco bannermani* Bates, 1923; *Telophorus kupeensis* Serle, 1951; *Apalis Bamendae* Bannerman, 1922; *Platysteira laticincta* Bates, 1926; *Ploceus batesi* Sharpe, 1908; *Turdoides gilberti* Serle, 1949; *Melichneutes robustus* Bates, 1909; *Laniarius atroflavus* Shelley, 1887; *Pternistis camerunensis* Alexander, 1909; *Zosterops melanocephalus*, Gray, 1862) (Fotso et al., 2001; Lepage, 2024). In the Sanaga-Maritime department where the areas in which our study was conducted, this geographical area presents a rich diversity with 339 species of birds (resident and migratory) distributed in 22 orders and 74 families (Lepage, 2024).

Furthermore, several studies have been conducted on birds in Cameroon (Louette, 1981; Nguembock et al., 2017, 2020; Mahamat, 2023; Nguembock et al., 2023; Azang et al., 2024). However, very few studies have been carried out in the coastal plain, particularly in the Sanaga-Maritime department (Louette, 1981; Fotso et al., 2001; Tchana, 2020). Yet, this ecosystem has ecological, biological, and economic importance (MINEPDED, 2017). The impact of human activities on the floristic structure of mangroves in the Cameroonian estuary is reflected in variations in density, tree height, and individual diameter depending on the degree of disturbance (MINEPDED, 2017).

More generally, the diversity, abundance and distribution of birds are also influenced by foraging opportunities at suitable nesting sites within various land cover types, including forests, shrublands, grasslands, wetlands, agricultural land, and urban areas (Price et al. 2014; Hu et al. 2017). However, in different regions of Cameroon, data are virtually non-existent for several ecosystems. This is the case for the Mouanko mangrove area and the Edea II evergreen forest, where several gaps remain, notably the lack of accurate and up-to-date data, particularly, the species present, their abundance, their migratory regime, and their sensitivity to human disturbances. In this study, we primarily pursued two objectives: firstly, we studied in depth the avifauna of these different areas, the diversity and composition of bird populations in the Mouanko and Edea II areas. Secondly, we analysed the distribution of birds according to seasons and habitats (evergreen forest and mangrove), comparing the different areas and thereby highlighting revealing trends and patterns.

Materials and Methods:-

Description of the study sites:-

Several study areas were investigated: Yoyo I (Os2) (N 03°64.889' E 09° 64.761') and Yoyo II (Os1) (N 03°66.635' E 09° 64.183') in the commune of Mouanko. In the commune of Edea II, the study areas were Ekite (Os3) (N 03° 80.356', E 10° 08.678') and Malimba (Os4) (N 03°84.920', E 10° 10.724') (Figure 1). These communes are located in the Sanaga-Maritime department of the Littoral Region. The climate of the Sanaga-Maritime department is

equatorial, with four seasons (two dry seasons and two rainy seasons). Average annual rainfall varies between 3,000 and 4,000 mm (CWCS, 2004), and average monthly temperatures range from 24.6°C to 28.7°C (Suchel, 1988; Fometé and Tchanou, 1998). The terrain is relatively flat, characterized by coastal plains, hills, and riverine areas, with elevations ranging from 0 to 120 m, the highest point being at Olombe (Nlomo, 2000, cited in Kouam 2012). The vegetation found is Atlantic evergreen rainforest, which occupies a coastal strip extending up to 200 km inland (Figure 2). It includes firstly the coastal forest on sandy plains, with locally particularly dense mangrove forest and tall stands (Figure 2). This area is heavily affected by forestry operations, but also by industrial plantations (oil palm, rubber) and food crops (Letouzey, 1985; Oliry 1986; Ajonina, 2010).

Investigation of the Mouanko and Edea II avifauna:-

During the bird survey in the Mouanko and Edea II sites, we used the mist-netting method, employing dark-coloured nylon nets with three vertical pockets running horizontally along the length of the net. This method is suitable for capturing small and medium-sized birds such as passerines. Our mist nets were fixed with the mounting poles, which had been chosen carefully, and the choice of an appropriate mist-netting site was important for the capture success. In order to ensure the capture success, we mainly identified their preferred flight paths, feeding areas, roosting and shaded sites. Generally, we started our capture very early in the morning (5:00 a.m.) and we finished very late in the evening (sometimes 6:30 p.m.). In order to avoid a skew in our bird survey, we used the same eight (8) mist nets in our different field mission and we captured in the same transects across the different seasons. Furthermore, we conducted eight (8) field missions (of four days each for a total of 64 field surveys) over 24 months (February 2022 to December 2024). Thus, eight linear transects, 1000 m long and 100 m wide, were used, comprising four in Mouanko and four in Edea II (Figure 1). To avoid cases of recapture, no transect was made twice in the same season. The identification manuals used were those of Borrow and Demey (2004, 2015). The captured and identified birds were then released at or near the capture sites.

Analysis of diversity and composition of the avifauna in the Mouanko and Edea II:-

Method for the calculation of abundance and occurrence of avifauna of Mouanko and Edea II:-

We used Excel program in order to calculate the relative abundance and the occurrence. We entered data and ran the software until the results were obtained.

Estimation of specific richness and calculation of the sampling effort:-

The estimate of species richness was calculated from estimators including Jackknife 1 and Jackknife 2, Chao 1, Ace and Bootstrap. These estimates took into account the number of species and their abundance by habitat and, by sector of the study area and throughout the study area. This made it possible to determine the estimated average number of species in the study area and to calculate the sampling effort.

Sampling effort (E) was made to assess the percentage of species sampled:

$$E = \left(\frac{\text{Number of observed species}}{\text{Number of estimated species}} \right) 100$$

Diversity indexes:-

Shannon index (H'): Shannon's diversity index represents the measure of the sum of degree of the uncertainty when it suggests prediction to which species would belong to an individual taken by chance in a collection of S species and N individuals. $H' = 0$ if the community has only one species; H' takes the maximum value $\log_2 S$ only when all species are represented by the same number of individuals. This index is determined by the following relationship:

$$H' = - \sum_{i=1}^s P_i \log_2 (P_i)$$

Where p_i = proportion of individuals of the species "i" ($P_i = n_i/N$); S = total number of species of the sample. n_i = number of individuals of the species "i"; N = total number of individuals of the sample. The Shannon index (H') increases when the number of the species of the community grows and, theoretically, it can reach elevated values. The value of H' varies from 1 to $\log_2 S$. In our study, the Shannon index was calculated with the PAST v5.2.1 software program.

Simpson index (λ): The Simpson index represents the proportion of abundance of the species "i". This index measures the degree of concentration when individuals are classified into types. It is determined by the following relationship:

$$\lambda = \sum_{i=1}^s \frac{n_i(n_i - 1)}{n(n - 1)}$$

Where n_i = number of individuals of the species “i”; n = total number of individuals of the sample. Nevertheless, the most popular of such indexes have been the inverse Simpson index ($1/\lambda$) and the Gini-Simpson index ($1 - \lambda$); both have also been called the Simpson index in the ecological literature. In our study, the Simpson index was calculated with the PAST v 5.2.1 software.

Analysis of distribution of the avifauna in the Mouanko and Edea II:-

Equity index: the Equitability index measures the distribution of individuals within species independently of specific richness. Its value varies from 0 (dominance of one species) to 1 (equal distribution of individuals within species).

Thus, the Equitability index of Pielou (J') is determined by the following formula:

$$J = \frac{H'}{H'_{\max}} = \frac{-\sum_{i=1}^S P_i \log_2(P_i)}{\log_2(S)}$$

H' = Shannon index; $H'_{\max} = \log_2 S$ (S = the total number of species). In our study, the Equitability index was calculated with the PAST v5.2.1 software. All these indexes have been obtained with a confidence threshold of 95%.

Comparison of the two sites with Permanova and pairwise Mann-Whitney tests:-

Shapiro-Wilk test:-

Before choosing the appropriate tests to compare the distribution of taxa within the habitats investigated, the Shapiro-Wilk test

- if $p \geq 0.05$: the distribution is considered normal;
- if $p < 0.05$: the distribution is considered non-Gaussian.

Mann-Whitney U test:-

Since the distribution did not follow a normal distribution and only two habitat types were observed (mangrove and evergreen forest), the Mann-Whitney U test was used to compare the distribution of taxa according to two environmental parameters (vegetation and seasons) between the two habitat types, under the following hypotheses:

H_0 : the distributions are identical;

H_1 : the distributions are different.

Permanova test:-

This is a non-parametric multivariate test used to compare biological communities between different sites, seasons, or treatments. It tests whether the differences between groups are significantly greater than those within groups. It is based on the distances between samples (Anderson, 2001) and relies on the dissimilarity matrix. A multivariate permutation analysis of variance (Permanova) was used to test differences in composition per survey between climatic seasons. A Permanova was also used to test differences in avian composition per survey based on vegetation type (evergreen forest and mangrove). If the communities within the groups are different, F-value is high and $P < 0.05$. If the communities are very similar, F-value is low and $P > 0.05$. In our study, these three tests have been done with the PAST v5.2.1 software; we entered data as described in the user guide and ran software until the results were obtained.

Results:-

Specific Richness of the avifauna of the Mouanko and the Edea II:-

Research work in the forest (Edea II) and mangrove (Mouanko) areas helped to obtain 643 individuals captured with Japanese nets. These 643 individuals were distributed among 10 orders, 28 families, 59 genera and 92 species (Table I). Of the taxa inventoried, 83.20% belong to the order Passeriformes and 16.8% to non-Passeriformes (Accipitriformes, Charadriiformes, Coliiformes, Columbiformes, Coraciiformes, Cuculiformes, Passeriformes, Piciformes, Strigiformes, Suliformes) (Table I).

The Table II shows that the occurrence frequencies of the species captured in the study area varied from 1.6% to 68.8%. Only one regular species was recorded: *Eurillas virens* (68.8%). four (04) species were considered incidental: *Eurillas latirostris* (42.2%), *Ploceus nigricollis* (48.4%), *Cyanomitra olivacea* (26.6%) and *Pycnonotus barbatus* (28.1%). The accidental species (27 species) were *Buccanodon duchaillui* (7.8%), *Chlorocichla falkensteini* (7.8%), *Cinnyris chloropygius* (9.4%), *Camaroptera brachyura* (12.5%), *Cossypha niveicapilla*

(14.1%), and *Chlorocichla simplex* (15.6%), etc. The rare species were the most numerous totalling 60 species (*Accipiter castanilius*, *Acrocephalus arundinaceus*, *Alethe castanea*, *Chrysococcyx klaas*, *Pogoniulus atroflavus*, *Eurillas gracilis*, *Hylia prasina*, *Cisticola bulliens*, *Sylvietta denti*, *Alcedo quadribachys*, *Psalidoprocne nitens*, *Gymnobucco bonapartei*, *Turtur afer*, *Hirundorustica*, *Phyllastrephus icterinus*, *Tringa stagnatilis*, *Fraseria cinerascens*, etc.) (Table II).

The capture of taxa using mist nets allowed for the inventory of 643 individuals belonging to 92 species across different sites. The Edea II forest area yielded 272 individuals grouped into 80 species (2023) and 170 individuals grouped into 39 species (2024). Meanwhile, the Mouanko mangrove area yielded 121 individuals belonging to 44 species in 2023 and 80 individuals grouped into 41 species in 2024. The most abundant species were *Eurillas virens* with 108 individuals (16.80%), *Eurillas latirostris* (6.38%), *Cyanomitra olivacea* (4.67%). The least abundant species were *Accipiter castanilius*, *Alethe diademata*, and *Cisticola anonymus*, each at 0.16% (Table II). Of the 92 species captured using Japanese nets, four (04) are Palearctic migrants (*Acrocephalus arundinaceus*, *Hirundo rustica*, *Saxicola rubetra*, and *Tringa stagnatilis*), whereas seven (07) are intra-African migrants (*Cecropis semirufa*, *Cecropis abyssinica*, *Halcyon leucocephala*, *Ispidina picta*, *Merops albicollis*, *Psalidoprocne pristoptera*, and *Terpsiphone viridis*) (Table II).

Sampling effort:-

Taking into account the specific richness at the different sectors of the study area, the estimators: Chao1 (105.92), Jackknife1 (117.5), Jackknife2 (127.83), Ace (110.11) and Bootstrap (104.09) gave an average of 113.09 species, which corresponds to approximately 113 species. The number of species observed being 92, the estimators therefore show that 21 species (113 estimated -92 species observed) were not encountered in this study area. Considering the number of estimated species and those sampled, the sampling effort for our study was 81.41%.

Diversity indices:-

Bird diversity across different study areas:

The Table III, the data shows the diversity indices for the different sites within the area. The Simpson index is 0.94 in the Edea II forest area; 0.96 in the Mouanko area; and 0.95 across the entire study area. The Shannon index values are 3.7, 3.64, and 3.81 in the Edea II, Mouanko, and overall study areas, respectively. The Pielou evenness index is 0.81 in Edea II; 0.92 in Mouanko; and 0.83 for the entire site. The species richness of the different sites is 80, 39, and 87 in the forest environments (Edea II 2023, 2024, and overall forest), 44 and 41 in the mangrove (2023 and 2024), and 92 in the study area.

Bird diversity indices by season:-

Regarding the seasons, the short dry season was the most diverse species (forest 2023, 2024 and mangrove 2023) and the most abundant in individuals (forest 2023 and 2024). The lowest diversity was observed in the forest in 2024 during the long dry season (LDS) ($H=2.02$), followed by the short rainy season (SRS) in the 2024 forest ($H'=2.27$). Conversely, in the mangrove, the lowest diversity was observed during the short rainy season in 2024 ($H=1.63$), followed by the short rainy season in 2023 (SRS) ($H'=2.29$).

Analysis of distribution of the avifauna in the Mouanko and Edea II:-

Regarding the overall Evenness index, in 2023 the Equitability index was 0.86 in forest while it was 0.90 in mangrove; however, in 2024 it was 0.80 in forest while it was 0.95 in mangrove (Table III).

Comparison of the two sites with the Shapiro-Wilk, Mann-Whitney U and Permanova tests:-

By applying the Shapiro-Wilk test to our data, we obtained for each habitat type, a p-value below the 5% threshold, confirming that the data collected did not follow a normal distribution; thus, the null hypothesis is rejected. The Mann-Whitney U test confirms the existence of a significant difference between the abundance distributions of the two bird samples from the two investigated habitats (Edea II evergreen forest and Mouanko mangrove) because p is less than 0.05 (0.021) (Table IV).

Furthermore, the Permanova test shows that the analysis of the avifauna distribution according to seasons showed no significant difference ($p=0.14$) nor the interaction between these two environmental parameters ($p=0.73$), whereas the vegetation showed a significant difference ($p=0.0019$) between these two habitat types (Table V).

Discussion:-

The study conducted in the study area showed a high representation of passerines (83.20%) compared to non-passerines (16.80%) (Table I). This result is similar to previous research conducted in Cameroon across the various

habitats investigated, including forest, savanna, and mountain areas (Louette, 1981; Fotso et al., 2001; Nguembock et al., 2017, 2019a, 2019b; Tchana, 2020; Nguembock et al., 2020; Kendeg, 2021; Mahamat, 2023; Odoukpe et al., 2023; Azang et al., 2024). This high abundance of passerine birds can be explained primarily by their short incubation period, allowing them to multiply rapidly during the breeding season, especially when food resources are readily available. Secondly, by their strong representation on the globe (Noske and Franklin, 1999; Auer et al., 2007).

The largest families were the Pycnonotidae, followed by the Nectariniidae and Alcedinidae, which were the most prevalent. This result corroborates the findings of several authors (Fatcheu, 2018; Kendeg, 2021; Azang et al., 2024) whose work has shown that forests are a favorable environment for birds. Furthermore, families such as the Pycnonotidae and Nectariniidae prefer open areas and secondary forests, primarily due to the availability of their food resources (Akinsola and Oluseye, 2004; Okosodo et al., 2016). Furthermore, of the 28 families identified, the most diverse were the Pycnonotidae (7 genera and 12 taxa), followed by the Lybiidae (6 genera and 9 taxa), Muscicapidae (6 genera and 8 taxa), Cisticolidae, and Estrildidae (4 genera and 5 taxa). This result is similar to those of Tchana (2020), Kendeg (2021), and Azang et al. (2024), which demonstrate that rainforests are highly diverse due to the presence of numerous resources such as food availability, varied microhabitats (canopy, lianas, epiphytes, etc.), understory, forest soil and others they contain.

Specific presentation of the birdlife of the study area:-

The Edea II site (2023) was the most abundant and diverse. However, in 2024, Edea II was less diverse (39) and less abundant (170). Meanwhile, the Mouanko area in 2023 was less abundant (121) than Edea II but more diverse (44) than Edea II (2024). The Mouanko area in 2024 was also more diverse (41) than the forest in 2024 but the least abundant (80). Overall, the evergreen forest area of Edea II was more diverse and more abundant than the mangrove area of Mouanko. This result corroborates the work of Essayas and Bekele (2011), who state that the abundance of birds in the forest is due to the diverse habitats it offers to different species. This could be attributed to the greater and more varied presence of food, nesting, and play resources that evergreen forests provide compared to mangroves. Regarding landscape composition, the forest ecosystem, particularly evergreen forest, with its shrubs and undergrowth, has a positive impact on birdlife (Wang, 2024). The results of our analyses showed *Eurillas virens* (68.8%) as a regular species, and several accessory species including *Eurillas latirostris* (42.2%), *Ploceus nigricollis* (48.4%); *Cyanomitra olivacea* (26.6%) and *Pycnonotus barbatus* (28.1%). This result is similar to that of Azang et al. (2024), who conducted an inventory in the rainforests of the upper Nyong River valley. This could be explained by the presence of diverse habitats (evergreen rainforests, agricultural areas, etc.) that provide sufficient resources for all taxa (Azang et al., 2024).

Diversity indices and distribution of the study area:-

The diversity indices obtained indicated that the study area is diverse. Generally, a Simpson index value of 0.95 indicates an absence of any one species dominating the others. This result is similar to that of Bunza et al. (2021), who conducted a study in swamp forests and mangroves with varied habitats in Nigeria. This finding corroborates the work of Nguembock et al. (2023) and Azang et al. (2024), who carried out their studies in the Central and Central and Eastern Regions of Cameroon, respectively. Similarly, MacArthur and MacArthur (1961) emphasize the importance of forest habitats for preserving birdlife, as they provide complex community structures, abundant food resources, and suitable nesting sites, while mitigating anthropogenic disturbances. A value of 3.81 for the Shannon index confirms the diversity of the study area (Table III). This result is similar to that of several authors (Tchana, 2020; Abassa, 2021; Nguembock et al., 2023; Mahamat, 2023; Nubuya et al., 2023; Azang et al., 2024). This diversity could be explained by low competition between taxa due to the presence of varied habitats and diverse and abundant resources in the forest and mangrove zone. The Pielou Evenness Index value of 0.83 indicates an even distribution of individuals among the taxa. This result corroborates those of several authors (Abassa, 2021; Bunza, et al., 2021; Nguembock et al., 2023; Mahamat, 2023; Nubuya et al., 2023; Azang et al., 2024), and could explain the diversity of habitats and resource abundance, such as the presence of readily available water (rivers, streams, and the sea) and food resources, are important factors.

Chace and Walsh (2006) showed that birds respond to changes in vegetation composition and structure, which influence their food resources. The habitats were ranked as follows: forest (2023) was the most diverse ($H = 3.9$; $J = 0.86$), followed by mangrove (2024) ($H = 3.77$; $J = 0.95$), followed by mangrove (2023) ($H = 3.6$; $J = 0.90$), and forest (2024) was the least diverse ($H = 3.03$; $J = 0.79$). Overall, forest ($H = 3.70$; $J = 0.81$) was more diverse than mangrove ($H = 3.64$; $J = 0.92$), but there was a slightly more even distribution of birds in forest compared to mangrove. These results are similar to those of several authors (Abassa, 2021; Bunza, et al., 2021; Nguembock et

al., 2023; Mahamat, 2023; Nubuya et al., 2023; Azang et al., 2024). Louette (1981), and Earnst and Holmes (2012) state that the distribution of birds is linked to vegetation (Table III).

Comparison of the two sites with Permanova and pairwise Mann-Whitney tests:-

The results of the analysis performed with the test Mann-Whitney U tests in the different site comparisons (evergreen forest vs. mangrove) showed a significant difference (Table IV). Similarly, analyses performed with two-way PERMANOVA revealed a significant difference between the habitats (evergreen forest vs. mangrove). This demonstrates a slight but significant variation in community structure. These results indicate overall heterogeneity across the entire study site (Table V). This could be due to the fact that the birds share the same biotopes and move throughout the study area, as well as the presence of year-round available resources and the absence of intraspecific competition between taxa. It should also be noted that the presence of evergreen vegetation throughout the year would provide permanent refuge for the different birds. Thus, Telleria and Santos (1994) emphasized that habitat structure influences species distribution. Furthermore, habitat size (Willis, 1979), feeding patterns (Marone, 1991) and floristic composition (Wiens and Rotenberry, 1981) also have an impact on species distribution.

Similarly, analyses carried out with Permanova, taking into account seasonal variations across different sites, showed no significant differences in the forest and mangrove areas (Table V). This could be due to the availability of resources (food, nesting sites, shelter, etc.) throughout the year. The equatorial climate with four seasons is favourable to the flourishing of birds year-round. This contrasts with findings from subtropical monsoon climates, which force some species to migrate, particularly in winter, or reduce their movements in summer (Nepali et al., 2021). Likewise, our results differ from those of Zean (2022) conducted in Ivory Coast, where the dry season appears to be the least favourable. Indeed, the dry season also coincides with the drying out of annual plants bordering bodies of water and the occurrence of bushfires (Poilecot and Loua, 2009); seasonal bird populations vary very little within each site. Similarly, several authors have demonstrated that seasonal variations influence bird abundance and distribution (Katuwal et al., 2016; Nepali et al., 2021; Zean, 2022). Furthermore, the results of our work show that the seasons do not correlate with the distribution of birdlife in the evergreen forest and the mangrove. These results are similar to those of several authors (Lefebvre, 1994; Lefebvre and Poulin, 1997; Mohd-Azlan, 2014) who show that few resources change drastically from one season to the next, that populations are stable, and that mangroves are used.





Fig 1: Geographical location of the study site.



(a)



(b)

Fig 2: Different types of habitats (a)=Evergreen forest(Edéa II) and (b)=mangrove(Mouanko)

Table I: orders, families, genera, species and absolute abundance of the study area

Orders	Families	Genres	Species	Abundances
Accipitriformes	1	1	1	1
Charadriiformes	1	1	1	2
Coliiformes	1	1	1	2
Columbiformes	1	1	2	9
Coraciiformes	2	4	7	64
Cuculiformes	1	1	1	2
Passeriformes	17	41	67	535
Piciformes	2	7	10	25
Strigiformes	1	1	1	1
Suliformes	1	1	1	2
Total	28	59	92	643

Table II: Frequency of occurrences and biogeographical statuses

No.	Species	Occurrence forest	Occurrence mangrove	Frequency of occurrence %	Absolute abundance forest	Absolute abundance mangrove	Total Absolute abundance	Total relative abundance % (RA%)	biogeographical status (Borrow and Demey, 2015)	IUCN status (2025)	MINF OF status (2020)
1	<i>Accipiter castanilius</i> Bonaparte, 1853	1	0	1.6%	1	0	1	0.16%	S	LC	B
2	<i>Acrocephalus arundinaceus</i> Linnaeus, 1758	1	2	4.7%	1	3	4	0.62%	P	LC	C
3	<i>Acrocephalus rufescens</i> Sharpe and Bouvier, 1877	1	2	4.7%	1	3	4	0.62%	S	LC	C
4	<i>Alcedo quadribachys</i> Bonaparte, 1850	1	1	3.1%	1	1	2	0.31%	S	LC	C
5	<i>Alethe castanea</i> Cassin, 1856	2	0	3.1%	2	0	2	0.31%	S	LC	C
6	<i>Alethe diademata</i> Bonaparte, 1850	1	0	1.6%	1	0	1	0.16%	S	LC	C
7	<i>Bleda notatus</i> Cassin, 1856	7	1	12.5%	30	1	31	4.82%	S	LC	C
8	<i>Buccanodon duchaillui</i> Cassin, 1855	2	3	7.8%	2	3	5	0.78%	S	LC	C
9	<i>Butorides striata</i> Linnaeus, 1758	0	2	3.1%	0	2	2	0.31%	S	LC	C
10	<i>Camaroptera brachyura</i> Vieillot, 1821	5	3	12.5%	8	5	13	2.02%	S	LC	C
11	<i>Cecropis abyssinica</i> Guérin-Méneville, 1843	1	0	1.6%	1	0	1	0.16%	M	LC	C
12	<i>Cecropis semirufa</i> Sundevall, 1850	2	0	3.1%	2	0	2	0.31%	S/M	LC	C
13	<i>Chamaetylas poliocephala</i> Bonaparte, 1850	1	0	1.6%	1	0	1	0.16%	S	LC	C
14	<i>Chlorocichla falkensteini</i> Reichenow, 1874	2	3	7.8%	2	4	6	0.93%	S	LC	C
15	<i>Chlorocichla flaviventris</i> Smith, 1834	2	2	6.3%	3	2	5	0.78%	S	LC	C
16	<i>Chlorocichla simplex</i> Hartlaub, 1855	6	4	15.6%	8	6	14	2.18%	S	LC	C
17	<i>Chrysococcyx klaas</i> Stephens, 1815	2	0	3.1%	2	0	2	0.31%	S	LC	C
18	<i>Cinnyris batesi</i> Ogilvie-Grant, 1908	6	5	17.2%	11	7	18	2.80%	S	LC	C
19	<i>Cinnyris chloropygius</i> Jardine, 1842	3	3	9.4%	5	4	9	1.40%	S	LC	C
20	<i>Cinnyris cupreus</i> Shaw, 1812	3	2	7.8%	5	2	7	1.09%	S	LC	C

21	<i>Cinnyris superbus</i> Shaw, 1812	1	0	1.6%	2	0	2	0.31%	S	LC	C
22	<i>Cisticola anonymus</i> Müller, 1855	1	0	1.6%	1	0	1	0.16%	S	LC	C
23	<i>Cisticola bulliens</i> Lynes, 1930	2	0	3.1%	2	0	2	0.31%	S	LC	C
24	<i>Clytospiza monteiri</i> Hartlaub, 1860	1	0	1.6%	1	0	1	0.16%	S	LC	C
25	<i>Colius striatus</i> Gmelin, 1789	2	0	3.1%	2	0	2	0.31%	S	LC	C
26	<i>Cossypha niveicapilla</i> Lafresnaye, 1838	6	3	14.1%	8	4	12	1.87%	S	LC	C
27	<i>Criniger calurus</i> Temminck, 1820	1	0	1.6%	1	0	1	0.16%	S	LC	C
28	<i>Cyanomitra cyanolaema</i> Jardine and Fraser, 1852	3	0	4.7%	4	0	4	0.62%	S	LC	C
29	<i>Cyanomitra olivacea</i> Smith, 1840	12	5	26.6%	22	8	30	4.67%	S	LC	C
30	<i>Cyanomitra verticalis</i> Latham, 1790	1	2	4.7%	1	2	3	0.47%	S	LC	C
31	<i>Elminia longicauda</i> Swainson, 1838	1	0	1.6%	2	0	2	0.31%	S	LC	C
32	<i>Estrilda melpoda</i> Vieillot, 1817	2	0	3.1%	4	0	4	0.62%	S	LC	C
33	<i>Estrilda nonnula</i> Hartlaub, 1883	2	1	4.7%	3	1	4	0.62%	S	LC	C
34	<i>Euplectes macroura</i> Gmelin, 1789	1	2	4.7%	1	2	3	0.47%	S	LC	C
35	<i>Eurillas gracilis</i> Cabanis, 1880	2	0	3.1%	2	0	2	0.31%	S	LC	C
36	<i>Eurillas latirostris</i> Strickland, 1844	19	8	42.2%	33	8	41	6.38%	S	LC	C
37	<i>Eurillas virens</i> Cassin, 1857	28	16	68.8%	83	25	108	16.80%	S	LC	C
38	<i>Fraseria cinerascens</i> Hartlaub, 1857	1	0	1.6%	1	0	1	0.16%	S	LC	C
39	<i>Gymnobucco bonapartei</i> Hartlaub, 1854	1	0	1.6%	1	0	1	0.16%	S	LC	C
40	<i>Halcyon leucocephala</i> Müller, 1776	2	2	6.3%	4	2	6	0.93%	M	LC	C
41	<i>Halcyon malimbica</i> Shaw, 1812	5	4	14.1%	6	4	10	1.56%	S	LC	C
42	<i>Halcyon senegalensis</i> Linnaeus, 1766	2	5	10.9%	3	7	10	1.56%	S/M	LC	C
43	<i>Hirundo rustica</i> Linnaeus, 1758	0	1	1.6%	0	1	1	0.16%	P	LC	C
44	<i>Hylia prasina</i> Cassin, 1855	2	0	3.1%	2	0	2	0.31%	S	LC	C
45	<i>Ispidina picta</i> Boddaert, 1783	7	4	17.2%	14	5	19	2.95%	M	LC	C
46	<i>Ketupa poensis</i> Fraser, 1854	1	0	1.6%	1	0	1	0.16%	S	LC	C
47	<i>Laniarius aethiopicus</i> Gmelin, 1789	1	0	1.6%	1	0	1	0.16%	S	LC	C
48	<i>Lybius vieilloti</i> Leach, 1815	1	0	1.6%	2	0	2	0.31%	S	LC	C
49	<i>Malaconotus cruentus</i> Lesson, 1831	1	2	4.7%	1	2	3	0.47%	S	LC	C
50	<i>Merops albicollis</i> Vieillot, 1817	1	4	7.8%	1	6	7	1.09%	M	LC	C
51	<i>Merops pusillus</i> Müller, 1776	3	3	9.4%	7	3	10	1.56%	S	LC	C
52	<i>Muscicapa sethsmithi</i> Someren, 1922	1	0	1.6%	1	0	1	0.16%	S	LC	C
53	<i>Muscicapa striata</i> Pallas, 1764	2	0	3.1%	2	0	2	0.31%	M	LC	C
54	<i>Neocossyphus poensis</i> Strickland, 1844	8	2	15.6%	10	2	12	1.87%	S	LC	C
55	<i>Nicator chloris</i> Valenciennes, 1826	1	0	1.6%	1	0	1	0.16%	S	LC	C
56	<i>Nicator vireo</i> Cabanis, 1876	0	3	4.7%	0	3	3	0.47%	S	LC	C
57	<i>Pardipicus caroli</i> Malherbe, 1852	1	0	1.6%	2	0	2	0.31%	S	LC	C
58	<i>Passer griseus</i> Vieillot, 1817	1	2	4.7%	1	2	3	0.47%	S	LC	C
59	<i>Phyllastrephus icterinus</i> Bonaparte, 1850	2	0	3.1%	2	0	2	0.31%	S	LC	C
60	<i>Phyllastrephus xavieri</i> Oustalet, 1892	1	4	7.8%	1	4	5	0.78%	S	LC	C
61	<i>Platysteira castanea</i> Fraser, 1843	4	0	6.3%	5	0	5	0.78%	S	LC	C

62	<i>Platysteira chalybea</i> Reichenow, 1897	1	0	1.6%	1	0	1	0.16%	S	LC	C
63	<i>Platysteira cyanea</i> Müller, 1776	4	4	12.5%	6	5	11	1.71%	S	LC	C
64	<i>Ploceus cucullatus</i> Müller, 1776	12	1	20.3%	18	2	20	3.11%	S	LC	C
65	<i>Ploceus nigerrimus</i> Vieillot, 1819	12	3	23.4%	15	4	19	2.95%	S	LC	C
66	<i>Ploceus nigricollis</i> Vieillot, 1805	3	28	48.4%	8	11	19	2.95%	S	LC	C
67	<i>Pogoniulus atroflavus</i> Sparrman, 1798	1	2	4.7%	2	2	4	0.62%	S	LC	C
68	<i>Pogoniulus bilineatus</i> Sundevall, 1850	2	0	3.1%	2	0	2	0.31%	S	LC	C
69	<i>Pogoniulus scolopaceus</i> Bonaparte, 1850	0	3	4.7%	0	6	6	0.93%	S	LC	C
70	<i>Pogoniulus subsulphureus</i> Fraser, 1843	1	0	1.6%	1	0	1	0.16%	S	LC	C
71	<i>Pogonornis bidentatus</i> Shaw, 1799	1	0	1.6%	1	0	1	0.16%	S	LC	C
72	<i>Prinia subflava</i> Gmelin, 1789	1	0	1.6%	1	0	1	0.16%	S	LC	C
73	<i>Psaldiprocne nitens</i> Cassin, 1857	1	0	1.6%	1	0	1	0.16%	S	LC	C
74	<i>Psaldiprocne pristoptera</i> Rüppell, 1840	1	2	4.7%	1	2	3	0.47%	S/M	LC	C
75	<i>Pycnonotus barbatus</i> Desfontaines, 1789	8	10	28.1%	12	17	29	4.51%	S	LC	C
76	<i>Saxicola rubetra</i> Linnaeus, 1758	2	3	7.8%	3	4	7	1.09%	P	LC	C
77	<i>Schistolais leucopogon</i> Cabanis, 1875	1	0	1.6%	1	0	1	0.16%	S	LC	C
78	<i>Spermestes bicolor</i> Fraser, 1843	6	1	10.9%	12	1	13	2.02%	S	LC	C
79	<i>Spermophaga haematina</i> Vieillot, 1807	3	0	4.7%	5	0	5	0.78%	S	LC	C
80	<i>Sylvietta denti</i> Ogilvie-Grant, 1906	1	0	1.6%	1	0	1	0.16%	S	LC	C
81	<i>Sylvietta virens</i> Cassin, 1859	1	2	4.7%	1	2	3	0.47%	S	LC	C
82	<i>Tchagra australis</i> Smith, 1836	2	0	3.1%	2	0	2	0.31%	S	LC	C
83	<i>Tchagra senegalus</i> Linnaeus, 1766	1	0	1.6%	1	0	1	0.16%	S	LC	C
84	<i>Terpsiphone batesi</i> Chapin, 1825	3	0	4.7%	3	0	3	0.47%	S	LC	C
85	<i>Terpsiphone rufocinerea</i> Cabanis, 1875	2	0	3.1%	2	0	2	0.31%	S	LC	C
86	<i>Terpsiphone viridis</i> Müller, 1776	1	5	9.4%	1	5	6	0.93%	S/M	LC	C
87	<i>Thescelocichla leucopleura</i> Cassin, 1855	1	2	4.7%	1	2	3	0.47%	S	LC	C
88	<i>Tricholaema hirsuta</i> Swainson, 1821	1	0	1.6%	1	0	1	0.16%	S	LC	C
89	<i>Tringa stagnatilis</i> Bechstein, 1803	0	2	3.1%	0	2	2	0.31%	P	LC	C
90	<i>Turdus pelios</i> Bonaparte, 1850	2	2	6.3%	2	2	4	0.62%	S	LC	C
91	<i>Turtur afer</i> Linnaeus, 1766	2	0	3.1%	4	0	4	0.62%	S	LC	C
92	<i>Turtur tympanistria</i> Temminck, 1809	4	0	6.3%	5	0	5	0.78%	S	LC	C
Total		/	/	100%	442	201	643	100%	/	/	/

LC: Least concern

B: Specie under partial protection

M: Intra- African migrant C: Common specie

P: Palaearctic migrant S: Resident

Table III: Bird diversity indices of Edea II and Mouanko

Habitats	seasons	years	Species	Individuals	Simpson_1-D	Shannon_H	Equity_J
Forest	LRS	2023	27	55	0.94	3.17	0.89
Forest	LDS	2023	25	49	0.96	3.29	0.95

Forest	SRS	2023	36	91	0.93	3.23	0.85
Forest	SDS	2023	47	77	0.98	3.91	0.94
Forest		2023	80	272	0.96	3.90	0.86
Forest	LRS	2024	18	50	0.91	2.68	0.87
Forest	LDS	2024	7	17	0.88	2.02	0.95
Forest	SRS	2024	12	46	0.87	2.27	0.86
Forest	SDS	2024	21	57	0.91	2.79	0.86
Forest		2024	39	170	0.91	3.03	0.80
Forest		Total forest	87	442	0.94	3.70	0.81
Mangrove	LRS	2023	13	26	0.89	2.48	0.88
Mangrove	LDS	2023	23	46	0.96	3.18	0.94
Mangrove	SRS	2023	9	15	0.90	2.29	0.92
Mangrove	SDS	2023	23	34	0.97	3.33	0.96
Mangrove		2023	44	121	0.96	3.60	0.90
Mangrove	LRS	2024	18	26	0.96	3.09	0.95
Mangrove	LDS	2024	19	28	0.96	3.12	0.95
Mangrove	SRS	2024	4	5	0.90	1.63	0.96
Mangrove	SDS	2024	18	21	0.99	3.25	0.98
Mangrove		2024	41	80	0.98	3.77	0.95
Mangrove		Total mangrove	47	201	0.96	3.64	0.92
Grand Total		Grand Total	92	643	0.95	3.81	0.83

LRS: Long rainy season

SRS: Short rainy season

LDS: Long dry season

SDS: Short dry season

Mann-Whitney test

Table IV: comparison of the Mann-Whitney test

	Forest evergreen	mangrove
Forest evergreen		0.021
mangrove	0.021	

Table V: Analysis of the structure of bird communities according to seasons and habitat using the Permanova test with 9999 permutations

Two-way PERMANOVA					
Source	Sum squares	Df	Mean square	F	p
habitats	0.36	1	0.36	9.70	0.0019
seasons	0.22	3	0.073	1.98	0.14
Interaction	0.065	3	0.021	0.58	0.73
Residues	0.29	8	0.037		
Total	0.94	15			

Conclusion:-

This study allowed us to identify a total of 92 captured species. The families were highly representative of passerines (83.2%). It revealed that the avifauna of the Edea II and Mouanko area is very diverse and that there is a significant difference in the distribution of relative abundances of taxa across the different study sites. Regarding the species composition of the bird populations, our study revealed that the investigated habitats are not similar. This difference within the sites appears to be directly related to vegetation. Furthermore, no significant difference in the distribution of relative abundances of species according to the seasons was observed. This homogeneity is linked to the year-round availability of resources and a four-season equatorial climate that is very favourable to the flourishing of the taxa. The information can then be used to manage and improve bird conservation at local, regional and international levels, given that a large part of Mouanko and Edea II's avian fauna is common to much of West, Central and East Africa. Many migratory and Palearctic bird species are also found there.

Acknowledgments:-

We would like to thank the Idea Wild Foundation for awarding us a grant to purchase equipment for the bird inventory. We also thank the non-governmental organizations, and we would like to thank the government (NGO) YOUTH4SDG, the village chiefs, and the guides for their support during data collection and for the essential documents, they provided for this report. Finally, we also thank the entire team at the Ornithology Unit of the Zoology Laboratory at the University of Yaoundé I for their contribution to this project.

Conflicts of interest:-

The authors declare that there are no conflicts of interest.

References:-

1. Abassa, M.A. (2021): Analysis of the distribution of forest avifauna in the vicinity of the Nkometou quarry and Introduction to the phylogenetic analysis of the family Indicatoridae Wainson, 1820 (birds, Piciformes). Thesis in Animal Biology and Physiology, Faculty of Science, University of Yaoundé I.
2. Adhikari, S., Sharma, H.P., Gautam, R., Basaula, R. (2020): Effects of weather conditions on the breeding success of the gray-headed prinia (*Prinia socialis*) in Manigram, Tiltottama Municipality, Nepal. Nepal. J. Environ. Sci. 8:11-16.
3. Ajonina, G. (2010): Final report on the completion of the mandate. Consultation Project GEF PPG.
4. Akinsola, I. A., Oluseye, A.O. (2004): A twelve-month field study of the West African Thrush *Turdus pelios* (Passeriformes: Muscicapidae). Part I: food and feeding ecology. Revista de Biología Tropical. 52: 1001- 1007.
5. Amano, T., Székely, T., Sandel, B., Nagy, S., Mundkur, T., Langendoen, T., Blanco, D., Soykan, C.U., Sutherland, W.J. (2018): Successful conservation of global waterbird populations depends on effective governance. Nature. 553: 199-202.
6. Anderson, M.J. (2001). A new method for non-parametric multivariate analysis of variance. Austral Ecology. 26:32-46.
7. Auer, S.K., Bassar, R.D., Fontaine, J.J., Martin, T.E. (2007): Breeding Biology of Passerines in a Subtropical Montane Forest in Northwestern Argentina. Condor. 109: 321-333.
8. Azang, E.D.O., Nguembock, B., Mahamat, S., Tadjoung, P., Kendeg, K.G.A., Tamesse, J.L. (2024): Diversity and distribution of birds in a Congo Basin wetland: The upstream part of the Nyong River valley in Cameroon along two regions (Centre and East). International Journal of Avian and Wildlife Biology. 8: 59-70.
9. Birdlife International, (2022). State of the World's Bird Populations: Prospects and Solutions for the Biodiversity Crisis. Cambridge, UK: BirdLife International.
10. Borrow, N., Demey, R. (2015): Birds of West Africa. Delachaux and Niestlé, Paris.
11. Borrow, N., Demey, R. (2004): Field guide to the Birds of Western Africa.
12. Bunza, O.D., Efenakpo, C., Egbochie, A., Uwatt, U.A., Shittu, Adeyanju, R.B., Mukhtar, I.C., Akupueet, B.B., Abubakar, (2021): Avian diversity and abundance at Finima Nature Park, Bonny Island, Rivers State, Nigeria ms. Journal of Agriculture and Environment. 17: 185-195.
13. Clarke, K.R. (1993): Non-parametric multivariate analysis of changes in community structure. Australian Journal of Ecology. 18:117-143.
14. Earnst, S., Holmes, A. (2012): Bird-habitat relationships in the inland Columbia basin, shrub steppe. Condor. 114: 15-29.
15. Essayas, K., Bekele, A. (2011): Species composition, relative abundance and distribution of the avifauna of the Entoto Nature Park and Escarpment, Addis Ababa. Journal Science. 34 :113-122.
16. Fomete, T., Tchanou, Z. (1998): The management of forest ecosystems in Cameroon at the dawn of the year 2000. Monographs of critical sites and annexes. IUCN, Yaoundé, Cameroon. 61-67.
17. Fotso, R., Dowsett-Lemaire, F., Dowsett, R.J., Scholte, P., Languy, M., Bowden, C. (2001): Cameroon in important bird areas of Africa and associated islands: priority sites for conservation. Birdlife Conservation Series. Pisces Publications and Birdlife International. 11: 133-159.
18. Gill, F., Donsker, D., Rasmussen, P. (2024): IOC World Bird List (v 14.1). [https://doi.org/ 10.14344/Ioc.ml.14.1](https://doi.org/10.14344/Ioc.ml.14.1).
19. Green, A.J., Elmberg J. (2014): Ecosystem Services Provided by Waterbirds: Ecosystem Services Provided by Waterbirds. Biological Reviews. 89: 105-22. <https://doi.org/10.1111/brv.12045>.

20. Green, A.J., El Hamzaoui, M., El Agbani, A.M., Franchimont, J. (2002): The Conservation Status of Moroccan Wetlands with Particular Reference to Waterbirds and to Changes since 1978. *Biological Conservation*. 104: 71-82. [https://doi.org/10.1016/s0006-3207\(01\)00155-0](https://doi.org/10.1016/s0006-3207(01)00155-0).
21. Hammer, O., Harper, D.A.T., Ryan, P.D. (2001): PAST: paleontological statistics software package for education and data analysis. *Palaeontologia Electronica*. 4(1):9.
22. Hu Y., Jin K., Huang, Z., Ding, Z., Liang, J., Pan, X., Hu, H., Jiang, Z. (2017): Elevational patterns of non-volant small mammal species richness in Gyirong Valley, Central Himalaya: Evaluating multiple spatial and environmental drivers. *Journal of Biogeography*. 44: 2764-2777. <https://doi.org/10.1111/jbi.13102>.
23. Kafando, W.A., Dibloni, O.T., Sanogo, S., Hien, M. (2023): Diversity of birdlife in urban landscapes of Africa: literature review and perspectives for Bobo-Dioulasso (Burkina Faso). *Journal of applied Biosciences*. 190: 20085-20100. <https://doi.org/10.35759/JABs.190.5>.
24. Katuwal, H.B., Basnet, K., Khanal, B., Devkota, S., Rai, S.K., Gajurel, J.P., Scheidegger, C., Nobis, M.P. (2016): Seasonal changes in bird species and trophic guilds along altitudinal gradients in the central Himalayas, Nepal. *PLoS ONE*. 11: e0158362. <https://doi.org/10.1371/journal.pone.0158362>.
25. Kemege, H.A.F. (2016): The Sanaga River and its impact on socio-economic development in the Sanaga-Maritime region (1960-2012). Thesis, University of Yaoundé I, Cameroon.
26. Kendeg, K.G.A. (2021): Assessment of the diversity and distribution of the avifauna of the village Oveng (Mbankomo) and phylogenetic analysis of the genus *Terpsiphone* Gloger, 1827 (Birds, Monarchidae). Master memory. University of Yaounde I.
27. Kouadio, K.P. (2015): Bird diversity, ecology of Pycnonotidae and Turdidae of Banco National Park (Ivory Coast). Doctoral thesis, Félix Houphouët-Boigny University, Ivory Coast.
28. Kouam, S. (2012): Inventory and evaluation of the stock of wood intended for smoking fish in the Douala-Edéa mangrove area. Report of the Bachelor's cycle, Institute of Fisheries Sciences of Yabassi, Douala, Cameroon.
29. Kougoum, P.G.N., Tamungang, S.A., Wansang, D., Tonleu, J., Soufo, L., Bilal, M., Taku, A., Tegua, A. (2024): Bird diversity in the Kalfou forest reserve and its peripheral zone, Far North of Cameroon. *Open Journal of Animal Sciences*. 14: 137-155. <https://doi.org/10.4236/ojas.2024.142010>
30. Lee, J.H., Park, D., Sung, H.C. (2012): Large-scale modeling of habitat associations of the endangered Korean ratsnake (*Elaphe schrenckii*). *Zoological Science*. 29: 281-285. <https://doi.org/10.2108/zsj.29.281>.
31. Lefebvre, G. (1994): Temporal dynamics of mangrove bird communities in Venezuela with special reference to migrant warblers. *The Auk*. 111: 405-415. <https://doi.org/10.2307/4088604>.
32. Lefebvre, G., Poulin, B. (1997): Bird communities in Panamanian black mangroves: Potential effects of physical and biotic factors. *Journal of Tropical Ecology*. 13: 97-113. Cambridge University Press. <https://www.jstor.org/stable/2559897>.
33. Lepage, D. (2024): Global bird lists. Avibase - The world ornithological base. Avibase - Bird Checklists of the World.
34. Letouzey, R. (1985): Phytogeographic map of Cameroon at 1/500000 with explanatory note. Institute of the International Plant Map. Toulouse, France. 142p.
35. Louette, M. (1981): The birds of Cameroon. An annotated checklist. *Verhandelingen. Koninklijke. Academie voor Wetenschappen. Letteren en Schone Kunsten van België*. 43:1-295.
36. Mahamat, S. (2023): Distribution, phylogeny and biogeography of the avifauna of the forest/savannah transition zone of the Adamaoua plateau (Adamaoua Region, Cameroon). Doctoral thesis PhD. University of Yaounde I.
37. Marone, L. (1991). Habitat characteristics influencing the spatial distribution of birds in the montane desert of Argentina. *Est. J. Ecol*. 1:77-86.
38. McPherson, M. J., Jetz, W. (2007): Type and spatial structure of distribution data and perceived determinants of geographic gradients in ecology: species richness of African birds. *World Ecology Biogeography*. 16:657-667. <https://doi.org/10.1111/j.1466-8238.2007.00318.x>.
39. Microsoft Excel. Microsoft® Office Excel. 2016.
40. MINEPDED. (2017): The mangroves of Cameroon: state of affairs and management. Report, 190 p. <https://ins-cameroun.cm/format/publication>.
41. Mohd-Azlan, J., Noske, R.A. and Lawes, M.J. (2014): Resource Partitioning by Mangrove Bird Communities in North Australia. *Biotropica*. 46: 331-340. <https://doi.org/10.1111/btp.12108>
42. Nepali, A., Khanal, S., Sapkota, S. Nanda, B.S. (2021): Seasonal variation in bird diversity in the Dhaneshwor Baikiwa community forest, Kavrepalanchowk district, Nepal. *Journal of Biodiversity Management and Forestry*. 10: 3.
43. Nguembock, B., Azang, E.D.O., Mahamat, S. (2019): Bird survey in a forest mountain of Congo basin (Elounden, Cameroon) and diversity analysis confirm high diversity of world second reserve and correlation between species diversity and vegetation. *International Journal of Avian & Wildlife Biology*. 4:118-127. <https://doi.org/10.15406/ijawb.2019.04.00163>.
44. Nguembock, B., Djopmou, H.N., Mahamat, S., Azang, E.D.O. (2019): study on bird diversity and abundance in a lowland of the central region of Cameroon (Ekoko II village) confirms high diversity in the Congo basin forest and a greater dominance in the distribution of species in the Lowlands. *International Journal of Animal Research*. 4:31. <https://doi.org/10.28933/ijar-2019-10-0905>.
45. Nguembock, B., Guehoda, Y., Mahamat, S. Azang, E.D.O. (2020): The birds of Cameroon: a bird survey from the Febe Mountain (Centre region, Cameroon) coupled with a diversity analysis confirm a higher bird diversity in mountains compared to lowlands in this region of the Congo Basin Forest. *Biodiversity International Journal*. 4: 140 -149. <https://doi.org/10.15406/bij.2020.04.00176>.

46. Nguembock, B., Mahamat, S., Kemtchouang, W., Azang, E.D.O. (2017): Bird survey of Abobo-Etetak hill (Yaounde, Cameroon) and a glance of the avifaunistic diversity of this hill (Abobo-Etetak). *The Open Ornithology Journal*.10:31-41. <https://doi.org/10.2174/1874453201710010031>.
47. Nguembock, B. (2008): Phylogenetic and biogeographical study of the forest avifauna of the mountains of Western Cameroon. France: Doctoral Thesis: University of Paris VI.
48. Nguembock, Kendeg, G.A., Mahamat, S., Azang, E.D.O. Abassa, M.A., Bitjaga, N.G.M., Okali, M.F. (2023): The birds of Cameroon: bird survey in a forest plain of the Congo basin forest (Oveng, Cameroon) confirms the great diversity of birds linked to the vegetation of the second largest biodiversity reserve in the world as well as the distribution pattern of taxa. *International Journal of Avian and Wildlife Biology*. 7: 111-1119. <https://doi.org/10.15406/ijawb.2023.07.00199>.
49. Noske, R.A. Franklin, D.C. (1999): Breeding seasons of land birds in the Australian monsoonal tropics: diverse responses to a highly seasonal environment. *Australian Journal of Biological Sciences*. 12:72-90.
50. Nubuya, S.G., David, D.L., Hammajoda, S.A., Yani, J.P., Barau, B.W. (2023): Avifauna diversity according to habitat types in Lau and Zing Local Government Areas, Taraba State, Nigeria. *International Journal of Multidisciplinary Research and Growth Assessment*. 04: 529-534.
51. Odoukpe, S.G.K., Assa, S.E. Gueye, F.M. (2023): State of the avifauna of some classified forests in central Ivory Coast: Species diversity, population structure and threats. *International Journal of Innovation and Applied Studies*. 39: 87-99.
52. Oindo, B.O., De By, R. A., Skidmore, A.K. (2001): Environmental factors influencing bird species in Kenya. *Afr. J. Ecol*. 39:295-302. <https://doi.org/10.1046/j.1365-2028.2001.00322.x>.
53. Okosodo, E.F., Orimaye, J.O., Obasogie, F.O. (2016): Food and Feeding Ecology of Common Bulbul (*Pycnonotus barbatus*) in Leventis Foundation Agricultural School Ilesa South Western Nigeria. *Greener Journal of Agricultural Sciences*. 6: 10 - 16. <https://doi.org/10.15580/GJAS.2016.1.121615169>.
54. Olivry, J.C. (1986): Rivers and streams of Cameroon. *Hydrological monographs: Mesres/Orstom*, 9: 745p.
55. Poilecot, P., Loua, N.S. (2009): Fires in the savannas of the Nimba mountains, Guinea. *Bois et Forêts des Tropiques*. 301: 51-66.
56. Price, T.D., Hooper, D.M., Buchanan, C.D., Johansson, U.S., Tietze, D.T., Alström, P., Ghosh-Harihar, M., Ishtiaq, F., Gupta, S.K., Martens, J., Harr, B., Singh, P., Mohan, D. (2014): Niche filling slows the diversification of Himalayan songbirds. *Nature*. 509: 222-225. <https://doi.org/10.1038/nature13272>.
57. Stuart, S.N. (1986): Conservation of Cameroon montane forests. *International Council for Bird Preservation (ICBP)*. Cambridge, UK. 250–263.
58. Suchel, J.B. (1988): *The Climates of Cameroon*. D'Etat, Saint-Etienne: University of St Etienne. 4:1188p.
59. Tchana, N.C. (2020): Investigation of spatial niche overlap between avifauna and chiropteroфаuna in the southern part of the Lake Ossa Wildlife Reserve (Littoral, Cameroon). Master's thesis. University of Yaoundé I.
60. Telleria, J.L. Santos, T. (1994): Factors involved in the distribution of forest birds of the Iberian Peninsula. *Bird Study* 41:161-169. <https://doi.org/10.1080/00063659409477216>.
61. Wiens, J.A. Rotenberry, J.J. (1981): Habitat Association and community structure of birds in a shrub steppe environment. *Ecological Monograph*. 51:21–42. <https://doi.org/10.2307/2937305>.
62. Zean, G.M. (2022): Avifauna of some forest relics in the Haut-Sassandra and Marahoué regions (central-west of Ivory Coast): diversity, abundance and effects of the area of forest fragments on stands. Doctoral thesis, Jean Lorougnon Guédé University, Ivory Coast.