



Journal Homepage: [-www.journalijar.com](http://www.journalijar.com)

INTERNATIONAL JOURNAL OF ADVANCED RESEARCH (IJAR)

Article DOI:10.21474/IJAR01/23514
DOI URL: <http://dx.doi.org/10.21474/IJAR01/23514>



RESEARCH ARTICLE

NUTRITIONAL QUALITY OF FOODS SOLD IN PRIMARY EDUCATION ESTABLISHMENTS IN THE DEPARTMENT OF TIIASSALE, CÔTE D'IVOIRE

Kakou Kouamé René Sylvestre¹, Meite Alassane², Monney Yapo Urbain³, Zahe Kollet Sylvere² and Kati-Coulibaly Séraphin²

1. Food Security and Nutritional Security Research Team, Doctoral School of Biology-Environment and Health, Felix HOUPHOUET-BOIGNY University, Abidjan, Côte d'Ivoire.
2. Laboratory of Biology and Health, UFR Biosciences, Felix HOUPHOUET-BOIGNY University, Abidjan, Côte d'Ivoire.
3. Laboratory of Environmental Sciences, UFR Governance and Sustainable Development, University of Bondoukou, BP V 81 Bondoukou, Côte d'Ivoire.

Manuscript Info

Manuscript History

Received: 12 March 2026

Final Accepted: 14 April 2026

Published: May 2026

Key words:-

Food, nutritional quality, macronutrients, students, Tiassalé.

Abstract

Access to food of adequate nutritional quality is essential for students' well-being and academic performance. Therefore, assessing the nutritional quality of foods sold in schools is important. This cross-sectional study aimed to evaluate the nutritional potential of foods sold to and consumed by pupils in primary schools in the Tiassalé Department, located in southern Côte d'Ivoire. A food consumption survey was conducted from October to November 2024 and enabled the collection of twenty-seven (27) food samples from four (4) primary schools in the Tiassalé Department. Physicochemical analysis was performed to determine the macronutrient composition and energy value of the foods. Analysis of variance (ANOVA) followed by Tukey's HSD tests was used to compare mean values. The survey revealed a variety of foods, including prepared meals, bread, fritters, juices, and snacks. The physicochemical analysis covered twenty-seven (27) samples of prepared meals, bread, and fritters. The results showed high fat and energy contents, with mean values ranging from 9.85 ± 3.41 g/100 g DM and 30.30 ± 7.74 g/100 g DM to 174.05 ± 26.80 kcal/100 g DM and 512.90 ± 52.77 kcal/100 g DM, respectively. In contrast, protein content was generally low (0.83 ± 0.04 g/100 g DM to 15.69 ± 0.03 g/100 g DM). Some foods had a high fiber content (15.62 ± 5.05 g/100g DM and 21.12 ± 3.40 g/100g DM) and carbohydrate content (24.65 ± 0.89 g/100g DM and 54.18 ± 5.63 g/100g DM). Students consume a variety of foods that differ from one school to another. These foods are high in energy and more concentrated in fat than in protein and fiber. Excessive and prolonged consumption of some of these foods can lead to nutritional disorders in children.

"© 2026 by the Author(s). Published by IJAR under CC BY 4.0. Unrestricted use allowed with credit to the author."

Corresponding Author:-Kakou Kouamé René Sylvestre

Address:-Food Security and Nutritional Security Research Team, Doctoral School of Biology-Environment and Health, Felix HOUPHOUET-BOIGNY University, Abidjan, Côte d'Ivoire.

Introduction:-

The quality of food products has become a major concern for populations today. The consumption of low-quality foods and the adoption of unhealthy eating habits are the main risk factors for poor health worldwide (Carl et al., 2017). They are determined by socioeconomic and political factors (Darmon and Drewnowski, 2015). Eating behavior influences food choices, food intake, eating habits, and food preferences (Debra et al., 2025). In recent decades, eating behaviors have shifted globally toward more processed, high-energy, and less varied diets (Leite et al., 2022). These diets are generally high levels of sugar, saturated fat, and sodium, and are of low nutritional quality. Furthermore, consuming high-quality, nutritious foods is essential for leading a healthy life and preventing disease (Catherine et al., 2025). Previous studies have shown that frequent consumption of nutrient-rich and healthy foods is associated with improved academic performance among children (Mou et al., 2023). School-aged children need adequate nutrition, both in quantity and quality, to maintain good health and achieve better academic results (Rufina et al., 2018). They spend most of their time at school and consume food estimated to provide nearly one-third of their daily energy intake (Tugault-Lafleur et al., 2017; Tugault-Lafleur and Black, 2020).

In Côte d'Ivoire, studies have been conducted on the nutritional status of primary school students and their academic performance (Touré et al., 2023; Zahe et al., 2017), but very little data exists on the nutritional quality of food sold in schools. This food constitutes an important part of the school food system in Côte d'Ivoire and warrants further investigation. The objective of this research is to evaluate the nutritional potential of food sold and consumed by students in primary schools in the Tiassalé Department.

Material and Methods:-

This cross-sectional food consumption study was conducted from October to November 2024 in four (4) primary schools in the Tiassalé Department. The Tiassalé Department is located in southern Côte d'Ivoire, 120 km from Abidjan. It is divided into four (4) sub-prefectures: N'douci, Morokro, Tiassalé, and Gbolouville. The sites were selected from two (2) sub-prefectures (Tiassalé and N'douci) that had a high number of schools and students, according to the 2023-2024 data from the Preschool and Primary Education Inspectorates (IEPP) of Tiassalé and N'douci. The survey sites in the two sub-prefectures were selected based on inclusion and exclusion criteria. All schools that simultaneously have a food production and sales area at the entrance and/or inside the school were included in the study. A total of four (4) schools per sub-prefecture were selected, two public and two private.

Survey on food consumption in schools:-

The survey was conducted at the Plateau School Group in N'douci, the Methodist School in N'douci (urban area), the N'zianouan School Group, and the AT-TABIAT School in N'zianouan (rural area). It consisted of observing and interviewing vendors and school principals about the various food items sold in the primary schools. Using a document, the listed food items were recorded, and twenty-seven (27) of them were collected as samples for physicochemical analysis.

Determination of the physicochemical composition of food:-

The method used for determining moisture and ash content is that described by AOAC (1990). The crude protein content of the samples was determined according to the Kjeldhal AOAC method (1990), from the determination of total nitrogen. The lipid content of the samples was determined according to the method described by AFNOR (1996), using the Soxhlet as the extractor. The fiber content of the food samples was determined by the Wolf method (1968). The total carbohydrate content was determined by difference according to the calculation method recommended by the FAO (2002). The theoretical energy value of the samples was calculated from the specific coefficients for proteins, lipids and total carbohydrates (FAO,2002).

Principal Component Analysis (PCA):-

Principal component analysis (PCA) is an exploratory statistical method used in the statistical analysis of physicochemical data. It was described by Gilbert and Ndèye(2003).

Statistical analysis:-

For statistical analysis, the data were entered into an Excel file and analyzed with the R software (version 4.5.1). The statistical tests used in the data processing were ANOVA followed by Tukey's HSD multiple comparison test at the 0.05 threshold. This latter test made it possible to compare the variance of the means of the different physicochemical parameters when the analysis of variance revealed significant differences (marked by the asterisk symbol).

Principal Component Analysis (PCA) was performed with the physico-chemical parameter data and the results are presented in table and figure form.

Results:-

Survey on food consumption in schools:-

The food consumption survey conducted in schools in the two sub-prefectures revealed a variety of foods that differed from one school to another. These foods included prepared dishes, baked breads, juices, fritters, and snacks. Prepared dishes consisted of attieké (cassava semolina) with fish, beef liver and stomach or chicken meat, rice with fat, attoupkou (steamed cassava cake) with sauce, baked bread with condiments, and fritters. A local dish called "NOUGOU," made with broken rice and moringa (*Moringa oleifera*) leaves, was also found in two schools. Juices included hibiscus, tomi, baobab, and mint-flavored milk, as well as imported juices in sachets or plastic containers. Snacks included candies, cookies, and chips.

Determination of the physicochemical composition of food:-

Table I presents the physico-chemical composition of food consumed by students in schools. Foods such as seasoned rice and broken rice combined with moringa leaves had the highest moisture content, at 65.00 ± 1.36 g/100 g dry matter (DM) and 59.20 ± 6.20 g/100 g DM, respectively. Following these were attieke served with chicken, fish, or cooked beef liver, with moisture content levels of 53.20 ± 0.85 g/100 g DM, 52.82 ± 3.02 g/100 g DM, and 51.76 ± 3.71 g/100 g DM, respectively. Fried bread was the least moist food, with a moisture content of 7.90 ± 3.12 g/100 g DM. The observed difference in moisture content between the foods was significant ($P < 0.05$). Regarding fat content, all foods had a high content except for the attoupkou and sauce dish and the rice with seasoning, with respective mean values of 4.30 ± 0.14 g/100 g DM and 9.85 ± 3.42 g/100 g DM. Fried bread had a high fat content, with an mean of 30.30 ± 7.74 g/100 g DM. The observed difference in fat content between the foods was significant ($P < 0.05$). Dietary fiber was most abundant in fried bread, attoupkou and chicken, and the other foods, with respective mean of 21.12 ± 3.40 g/100 g DM and 19.50 ± 0.70 g/100 g DM. Attoupkou and sauce was the food with the lowest fiber content (2.25 ± 0.35 g/100 g DM). The results of the analysis show a statistically significant difference ($P < 0.05$). Foods such as bakery bread and cooked beef liver or egg had the highest protein content, at 15.70 ± 0.03 g/100 g DM and 15.65 ± 0.03 g/100 g DM, respectively. The observed difference was significant ($P < 0.05$). The results showed that some foods were rich in carbohydrates, with mean values ranging from 41.57 ± 0.89 g/100 g DM to 54.18 ± 5.63 g/100 g DM. The energy value of the foods was high, ranging from 174.05 ± 26.80 kcal/100 g DM to 512.90 ± 52.77 kcal/100 g DM. The observed difference between carbohydrate values was significant ($P < 0.05$). The same was true for energy values.

Table I : Physicochemical composition of different foods

Food	Moisture (g/100g MF)	Lipid (g/100g DM)	Fiber (g/100g DM)	Ash (g/100g DM)	Protein (g/100g DM)	Total carbohydrate (g/100g DM)	Energy values (kcal/100g DM)
Atoukpou_sauce	50,90±0,42 _{bcd}	4,30±0,14 ^c	2,25±0,35 _b	2,40±0,2 _{8^{ab}}	0,83±0,04 ^e	41,57±0,89 _{ab}	208,30±2,12 _d
Attieke_liver	51,76±3,71 _{bc}	14,36±3,1 ^b _c	9,50±3,52 ^a _b	2,13±0,3 _{7^b}	3,71±1,34 _{de}	28,02±1,70 _c	256,23±27,4 _{8^{cd}}
Attieke_fish	52,82±3,02 _{bc}	14,32±2,82 _{bc}	16,62±9,0 _{7^{ab}}	3,60±0,9 _{1^a}	6,80±1,19 ^c	22,45±4,33 _{cde}	245,93±21,8 _{6^{cd}}
Attieke_chicken	53,20±0,85 _{abc}	12,00±0,57 _{bc}	19,50±0,7 _{0^{ab}}	2,90±0,1 _{4^{ab}}	4,95±0,04 ^e _{de}	26,96±0,46 _{cd}	235,60±6,79 _{cd}
Attieke_beef tripe	50,60±0,00 _{bcd}	15,10±0,99 _{bc}	18,00±0,0 _{0^{ab}}	2,90±0,1 _{4^{ab}}	6,75±0,04 ^e _d	24,65±0,89 _{cde}	261,50±5,52 _{cd}
Fritters	15,23±11,0 _{5^e}	20,90±8,58 _{ab}	14,67±3,6 _{4^{ab}}	1,43±0,7 _{3^b}	8,25±2,03 ^c	54,18±5,63 _a	437,83±89,0 _{8^{ab}}
Avocado_bread	47,20±3,11 _{bcd}	17,70±0,42 _{abc}	4,5±0,70 ^b	3,20±0,0 _{0^{ab}}	9,84±0,01 _{bc}	22,07±3,53 _{cde}	286,90±10,3 _{2^{cd}}
Liver_bread	35,10±0,14 _d	16,60±1,13 _{bc}	14,50±2,1 _{2^{ab}}	2,50±0,4 _{2^{ab}}	15,70±0,0 _{4^a}	30,11±1,73 _{bc}	332,60±3,39 _{bc}
Fried_bread	7,90±3,12 ^e	30,30±7,74 _a	21,12±3,4 _{0^a}	1,75±0,4 _{1^b}	8,24±1,12 ^c	51,81±3,11 _a	512,90±52,7 _{7^a}
Egg_bread	44,50±0,14 _{cd}	12,80±0,00 _{bc}	6,75±0,35 ^a _b	2,60±0,2 _{8^{ab}}	15,65±0,0 _{4^a}	24,46±0,46 _{cde}	275,60±1,70 _{cd}
Fish_ver_bread	40,73±1,09 _{cd}	17,57±3,88 _{bc}	10,33±5,2 _{2^{ab}}	2,83±0,1 _{5^{ab}}	11,48±0,9 _{8^{ab}}	27,39±5,04 _c	313,57±17,3 _{6^c}
Fat_rice_seas	65,00±1,36 _a	9.85±3,42 ^c	4,87±4,77 _b	3,80±1,1 _{7^a}	8,80±0,30 _{bc}	12,54±1,37 _{de}	174,05±26,8 _{0^d}
Moringa_rice	59,20±6,20 _{ab}	13,30±2,61 _{bc}	15,62±5,0 _{5^{ab}}	2,00±0,5 _{6^b}	8,70±2,66 _{bc}	16,79±6,81 _e	221,70±29,4 _{9^d}
Pr(>F)	2e-16 ***	2.4e-06 ***	0.000223 ***	4.35e-06 ***	2.44e-13 ***	1.6e-06 ***	0.000567 ***
Food	Moisture (g/100g MF)	Lipid (g/100g DM)	Fiber (g/100g DM)	Ash (g/100g DM)	Protein (g/100g DM)	Total carbohydrate (g/100g DM)	Energy values (kcal/100g DM)
Atoukpou_sauce	50,90±0,42 _{bcd}	4,30±0,14 ^c	2,25±0,35 _b	2,40±0,2 _{8^{ab}}	0,83±0,04 ^e	41,57±0,89 _{ab}	208,30±2,12 _d
Attieke_liver	51,76±3,71 _{bc}	14,36±3,1 ^b _c	9,50±3,52 ^a _b	2,13±0,3 _{7^b}	3,71±1,34 _{de}	28,02±1,70 _c	256,23±27,4 _{8^{cd}}
Attieke_fish	52,82±3,02 _{bc}	14,32±2,82 _{bc}	16,62±9,0 _{7^{ab}}	3,60±0,9 _{1^a}	6,80±1,19 ^c	22,45±4,33 _{cde}	245,93±21,8 _{6^{cd}}
Attieke_chicken	53,20±0,85 _{abc}	12,00±0,57 _{bc}	19,50±0,7 _{0^{ab}}	2,90±0,1 _{4^{ab}}	4,95±0,04 ^e _{de}	26,96±0,46 _{cd}	235,60±6,79 _{cd}
Attieke_beef tripe	50,60±0,00 _{bcd}	15,10±0,99 _{bc}	18,00±0,0 _{0^{ab}}	2,90±0,1 _{4^{ab}}	6,75±0,04 ^e _d	24,65±0,89 _{cde}	261,50±5,52 _{cd}
Fritters	15,23±11,0 _{5^e}	20,90±8,58 _{ab}	14,67±3,6 _{4^{ab}}	1,43±0,7 _{3^b}	8,25±2,03 ^c	54,18±5,63 _a	437,83±89,0 _{8^{ab}}
Avocado_bread	47,20±3,11 _{bcd}	17,70±0,42 _{abc}	4,5±0,70 ^b	3,20±0,0 _{0^{ab}}	9,84±0,01 _{bc}	22,07±3,53 _{cde}	286,90±10,3 _{2^{cd}}
Liver_bread	35,10±0,14 _d	16,60±1,13 _{bc}	14,50±2,1 _{2^{ab}}	2,50±0,4 _{2^{ab}}	15,70±0,0 _{4^a}	30,11±1,73 _{bc}	332,60±3,39 _{bc}

Fried_bread	7,90±3,12 ^e	30,30±7,74 ^a	21,12±3,40 ^a	1,75±0,41 ^b	8,24±1,12 ^c	51,81±3,11 ^a	512,90±52,77 ^a
Egg_bread	44,50±0,14 ^{cd}	12,80±0,00 ^{bc}	6,75±0,35 ^a	2,60±0,28 ^{ab}	15,65±0,04 ^a	24,46±0,46 ^{cde}	275,60±1,70 ^{cd}
Fish_ver_bread	40,73±1,09 ^{cd}	17,57±3,88 ^{bc}	10,33±5,22 ^{ab}	2,83±0,15 ^{ab}	11,48±0,98 ^{ab}	27,39±5,04 ^c	313,57±17,36 ^c
Fat_rice_seas	65,00±1,36 ^a	9.85±3,42 ^c	4,87±4,77 ^b	3,80±1,17 ^a	8,80±0,30 ^{bc}	12,54±1,37 ^{de}	174,05±26,80 ^d
Moringa_rice	59,20±6,20 ^{ab}	13,30±2,61 ^{bc}	15,62±5,05 ^{ab}	2,00±0,56 ^b	8,70±2,66 ^{bc}	16,79±6,81 ^e	221,70±29,49 ^d
Pr(>F)	2e-16 ***	2.4e-06 ***	0.000223 ***	4.35e-06 ***	2.44e-13 ***	1.6e-06 ***	0.000567 ***

Pr(>F) 2e-16 *** 2.4e-06 *** 0.000223 *** 4.35e-06 *** 2.44e-13 *** 1.6e-06 *** 0.000567 *** Ver: vermicelli; seas : seasoning; DM: Dry Matter; MF: Fresh Matter The data are expressed as mean ± standard error (M±SEM). a, b, c, d, e: Values assigned different letters in the same column are significantly different (P<0.05).

Table II: Principal Component Analysis of the Physicochemical Parameters of Food

Main components	1	2	3	4	5
Equity	3.97	1.22	0.89	0.57	0.36
Variance (%)	56.76	17.37	12.63	8.13	5.11
Cumulative variance (%)	56.76	74.13	86.76	94.89	100

Principal Component Analysis:-

Principal Component Analysis (PCA) was performed to classify and process information on the physicochemical parameters of the food by examining the correlations between variables. It revealed five (5) principal components, whose eigenvalues for the axes or dimensions (Dim 1, Dim 2, Dim 3, Dim 4, and Dim 5) are shown in (Table II) and (Figure 1). The two dimensions, Dim 1 and Dim 2, describe the correlations between the variables related to the physicochemical parameters, representing 74.13% of the total information, with respective values of 56.76% and 17.37%. The correlation circle resulting from the PCA allowed for the examination of the relationships between the variables in the factorial space defined by the first two principal components (Dim 1 and Dim 2) (Figure 2). This circle shows a positive correlation between lipids and energy value, between fiber and lipids, and between energy value and total carbohydrates.

There is a negative correlation between protein and total carbohydrates, between protein and energy value, and between moisture and other physicochemical components. These variables are well represented in Dim1 and Dim2 and indicate a strong contribution to the explained variance (74.13%). The factorial design, represented by the two dominant dimensions of total variance (Dim 1 and Dim 2) with respective values of 56.76% and 17.37%, showed the distribution of food consumption by study area. Foods such as attoukpou and sauce, bread and avocado, attieké and cooked beef tripe, bread and cooked beef liver, fried bread, and attieké and fried chicken are consumed specifically in schools in both rural and urban areas. However, foods such as fritter, bread and egg, bread and vermicelli, rice with moringa leaves, attieké and fish, and attieké and cooked beef liver are consumed in schools in both rural and urban areas and are highly represented (Figure 3).




Figure 1: Distribution of inertia between dim1 and dim2




Figure 2: Correlation circle of physicochemical parameters




Figure 3: Factorial design of the foods studied

Atoukpou_sa : Attoukpou_sauce ; Attieke_chick : Attieke_chicken ; Avocado_bre : Avocado_bread ; Fish_ver_bre : Fish_vermicelli_bread ; Fat_rice_sea : Fat_rice_seasoning.

Discussion:-

This study examined the food sold to students in primary schools. This food was diverse and varied, including prepared meals, bread, fritters, juices, and sweets. Prepared meals, bread, fritters, and juices were the most frequently consumed foods by students in these schools. Similar observations were made in schools in Yaoundé 1st, Cameroon, and N'Djamena, Chad, where the categories of food sold included prepared meals, pastries, fritters, vegetables, snacks, drinks, and ice cream. Prepared meals accounted for 43% and pastries for 24% in Yaoundé, while cereal products represented 60% in N'Djamena (Annie-Claude et al., 2017; Naibe, 2019).

The physicochemical analysis of commonly consumed foods, such as prepared dishes, baked breads, and fritters, revealed their macronutrient and energy composition. The lipid content of the studied foods was high, with mean values ranging from 12.00 ± 0.57 to 30.30 ± 7.74 g/100 g DM, except for attoukpou with sauce and rice with seasoning, which had average values of 4.30 ± 0.14 and 9.85 ± 3.42 g/100 g DM, respectively. This high lipid content is primarily due to the significant use of refined palm oil or other oils in preparation. Attoukpou (steamed cassava cake) had a low lipid content, which came from the sauce. Research conducted by Lecerf has shown that palm oil contains 50% saturated fatty acids, generally in positions 1 and 3, and is rich in carotenoids and vitamin E. Copra oil (coconut oil) contains 94% saturated fatty acids, and cocoa butter 60%. Saturated fatty acids increase both HDL and LDL cholesterol, the latter (LDL) being considered hypercholesterolemic. High LDL cholesterol levels are associated with an increased risk of cardiovascular disease (Lecerf, 2013).

The fiber content was high in some of the foods studied, ranging from 15.62 ± 5.05 to 21.12 ± 3.40 g/100 g DM. Fried bread had the highest content (21.12 ± 3.40 g/100 g DM). Studies conducted in Addis Ababa, Ethiopia, on school meals provided by programs showed an adequate daily fiber intake. Fiber-rich foods came from cereal-based meals such as white bread. However, these studies reported that the fat content of the meals was less than one-third of the recommended daily intake (Zelalem et al., 2018). Indeed, dietary fiber absorbs water and solidifies waste for healthy bowel movements and helps prevent certain nutritional disorders (Chhabra, 2018). Regarding protein

content, the foods studied had low levels, except for bakery bread and cooked beef liver or egg, which had high levels of 15.70 ± 0.03 g/100 g DM and 15.65 ± 0.03 g/100 g DM, respectively. Indeed, the foods studied mostly contained fish, meat, eggs, and leafy green vegetables. However, this low protein content could be explained by the small portions used in food preparation. Furthermore, the foods contained less plant-based protein. The protein values of the foods studied are lower than the recommended daily intake of protein for children and adolescents (FAO/WHO/UNU, 1973). These foods do not meet the protein requirements of students.

The total carbohydrate content of the foods ranged from 12.54 ± 1.37 g/100 g DM to 54.18 ± 5.63 g/100 g DM. More than half of the foods had a low total carbohydrate content according to WHO recommendations. The WHO guidelines on carbohydrate intake for adults and children indicate that total carbohydrate intake from foods represents approximately 40% to 70% of total energy intake and is considered compatible with a healthy diet (WHO, 2023). Furthermore, the study by Seidelmann et al. (2018) on dietary carbohydrate intake and mortality in an adult population showed that diets high (> 70%) or low (< 40%) in carbohydrates were associated with increased mortality. Carbohydrates provide energy to the body's cells, especially the brain, which is a glucose-dependent organ (Rufina et al., 2018).

The energy content of these foods was high, ranging from 174.05 ± 26.80 to 512.90 ± 52.77 kcal/100g DM. This intake could meet the energy requirements of children as indicated in the recommendations of FAO/WHO/UNU (2001). Principal Component Analysis (PCA), performed to analyze and describe the correlations between the variables of the physicochemical parameters of the foods, showed that 74.13% of the information was represented by the two dimensions (Dimension 1 and Dimension 2). The variances show that axis 1, represented by Dimension 1, is the most important, as it describes more than half of the information (56%). The positive correlation between the physicochemical parameters indicates a trend in the same direction. Conversely, the negative correlation between the parameters indicates a trend in opposite directions. The factorial plane defined by the two axes (Dimension 1 and Dimension 2) shows the distribution of foods by consumption area (urban or rural schools). Thus, some foods are consumed in rural or urban schools, and others are consumed in both study areas. Although the foods sold to schoolchildren provides energy and certain nutrients, they do not adequately meet their protein and dietary fiber requirements. Consequently, students may be exposed to both forms of malnutrition due to nutrient excess or deficiency. However, quality school meals contribute to improving students' nutritional status. This study suggests improving the quality of food sold in schools and to strengthen nutrition education programs for both vendors and students. The micronutrient content of these foods was not evaluated in this study and remains an important point to develop in further research.

Conclusion:-

This study identified the foods sold in primary schools. These foods are diverse and varied in nature. Physicochemical analysis revealed a high energy content due to their high lipid content, but low protein and fiber content. Consuming these foods is important for providing energy and helping schoolchildren stay focused during class. However, their high lipid content and low protein content make these foods less nutritious, and excessive or prolonged consumption of some of them can lead to nutritional deficiencies in children.

Acknowledgements:-

I would like to express my deep gratitude and thanks to the head of the Nutrition and Pharmacology Department at Félix HOUPOUËT BOIGNY University and the scientific director for supervising and directing this research. I would especially like to thank Dr. Urbain Monney, Dr. Sylvere Zahe, and Ph.D. student Franck Kouame for their contributions to this work.

Bibliographical References:-

1. AFNOR, 1996. Water testing: protocol for evaluating an alternative method of quantitative physicochemical analysis in relation to a reference method. In French Standardization Association (ed). XP T ISSN 0335-3931: Paris, France; 210.
2. Annie-Claude P., Malika M., Majesté MP, Germain N., 2017. Current State of the Food Supply System in Some Primary Schools in the City of Yaoundé (Cameroon). *European Scientific Journal*, 13 (18): 1857-7881
3. AOAC, 1990. Official methods of analysis. Association of Official Analytical Chemists. Washington DC, USA, 684 P.

4. Carl L., Jessica E., Raneria, b., Katherine WS, Patrick K., Patrick V. DammeCD., Chat V., Daniela P., Wouter V., Gina K., Danny H., Francis OO, Gervais N.-B., Bernard DB, Disna R., Hoang LK, Roseline R., Céline T., 2017. Food species richness as a measure of food biodiversity and the nutritional quality of diets. 115 (1) 127–132.
5. Catherine JA, Kristina M., Alexandre G., Lydia D., Layra Cintrón-R., 2025. The nutritional quality of university dining options varies according to location, convenience level, and accessibility: Perspectives from a pilot study on the evaluation of university food environments. <https://doi.org/10.1016/j.jafr.2025.102057>.
6. Chhabra S. 2018. Dietary fiber nutrition and health benefits. *Function Food Human Health*:15-25. Singapore, Singapore: Springer Singapore.
7. Darmon N, Drewnowski A., 2015. Contribution of food prices and food costs to socioeconomic disparities in diet quality and health: a systematic review and analysis. *Nutr Rev*73:643–660.
8. Debra KM, Tacad.,B., Kamil B., Nancy L. Keim. 2025. Differential associations of eating behavior traits, food preference, motivations of food choice on diet intake and diet quality in adult females and males from the USDA nutritional phenotyping study.<https://doi.org/10.1016/j.appet.2025.108048>
9. FAO. 2002. Food energy methods of analysis and conversion factors. FAO Ed, Rome, Italy, 97P.
10. FAO/WHO/UNU Rome, 17-24 October 2001. Human energy requirements. Technical reports on food and nutrition. Rome, 103P.
11. FAO/WHO/UNU. 1973. Energy and protein requirements: Technical report of a joint expert consultation. WHO, Geneva. 123p.
12. Gilbert Saporta, Ndèye Niang. 2003. Principal component analysis. Gérard Govaert. Data analysis, Hermes, pp.19-42, 978-2-7462-0643-4.
13. Laurencia T., S.,O., Kourfom G., Aly S., Fabrice B., Bréhima D. 2016. Evaluation of the nutritional suitability of foods used in complementary feeding of young children in Burkina Faso.<https://www.researchgate.net/publication/309784941>
14. Lecerf J.-M., 2013. Palm oil, such a bad oil? *Science et pseudo-science*; 306:42-49.
15. Leech RM, McNaughton SA, Timperio A. 2014. The clustering of diet, physical activity, and sedentary behavior in children and adolescents: a review. *Proceedings of the International Journal of Behavioral and Nutritional Health*;11:4. doi:10.1186/1479-5868-11-4
16. Leite, FHM, Khandpur, N., Andrade, GC., Anastasiou, K., Baker, P., Lawrence, M. and Monteiro, CA., 2022. Ultra-processed foods should be at the heart of the dialogue on global food systems and action for biodiversity. *BMJ Global Health*, 7 (3).<https://doi.org/10.1136/bmjgh-2021-008269>.
17. Mou, Y., Block, E., Barroso, M., Jansen, PW, White, T., Voortman, T., 2023. Models, brain morphology and cognitive performance in children: results of a prospective study in the general population. *Eur. J. Epidemiol.* 38, 669–687.
18. Naibe MS, 2019. Assessment of food quality in schools in the city of N'Djamena, Chad. Master's thesis, University of N'Djamena-Chad, Faculty of Graduate and Postdoctoral Studies. p127.
19. Rufina NB Ayogu, Paul EE, Vivien CA, Henrietta NE, Uche VA 2018. Nutritional value of school meals and their contributions to energy and nutrient intakes of rural school children in Enugu and Anambra States, Nigeria.<https://doi.org/10.1186/s40795-018-0216-0>.
20. Seidemann SB, Claggett B, Cheng S, Henglin M, Shah A, Steffen LM, Solomon Sc.D. 2018. Dietary carbohydrate intake and mortality: a prospective cohort study and meta-analysis. *Lancet Public Health*.3(9):e419–e428.
21. Touré M., F., Zahe K., Y., Assamois F., F., Marie L., Bitty, Séraphin C., 2023. Evaluation of the nutritional status of schoolchildren in the commune of Abobo. *EAS Journal of Nutrition and Food Sciences*. 5(5):2663-7308.
22. Tugault-Lafleur, CN., and Black, JL., 2020. Lunch on school days. Review of contributions to nutrient and food group intake and differences according to consumption setting. *J. Acad. Nutr. Diet.* 120. doi:10.1016/j.jand.2020.01.011.
23. Tugault-Lafleur, CN., Black, JL., Barr, SI., 2017. Study of food intake during the school day in Canadian children. *Applied Physiology, Nutrition and Metabolism*.42(10):1064-1072. doi:10.1139/apnm-2017-0125.
24. WHO. 2023. Carbohydrate intake for adults and children: WHO guideline summary. Geneva, World Health Organization. License: CC BY-NC-SA 3.0 IGO
25. Wolf, JP1968. Manual of food analysis. Techniques et Documentation-Lovoisier, Paris.
26. Zahé KYAS., Méité A., Ouattara H., Dally T., Kouamé KG., Aké-Tano O. Kati-Coulibaly S. 2017. Diet, nutritional status and school performance of schoolchildren in the district of Abidjan: the case of Yopougon and Bingerville. *Journal of Nutrition and Health Sciences*, 4 (1): 2393-9060.
27. Zelalem D., Eshetu W., Samuel K., Mulugeta E., Yohannes C., Melkamu T., Meti T., Hanna S., Dilu S., Mogessie A.2018. School feeding contributed valuable dietary energy and nutrients despite suboptimal supply to school-age children and adolescents at primary schools in Addis Ababa, Ethiopia. *Nutrition* 102 (2022) 111693.