



RESEARCH ARTICLE

TO STUDY PHENOTYPIC IDENTIFICATION OF CANDIDA ISOLATES AMONG CLINICAL SAMPLES AT TERTIARY LEVEL HOSPITAL, WESTERN RAJASTHAN

Devaki Nandan Kushwah, Sanju Pannu, Richa Agrawal and Prabhu Prakash

Manuscript Info

Manuscript History

Received: 8 April 2026

Final Accepted: 10 May 2026

Published: June 2026

Key words:-

Candida, Candidiasis, MALDI-TOF MS, Phenotypic Identification, Non-albicans Candida, Candichrome Agar.

Abstract

Background: Candida species are important opportunistic fungal pathogens responsible for a wide spectrum of infections ranging from superficial mucosal disease to invasive candidiasis. Accurate and timely species identification is essential because different Candida species exhibit varying antifungal susceptibility patterns. The increasing occurrence of non-albicans Candida species has further emphasized the need for reliable identification methods.

Objectives: To study the phenotypic identification of Candida isolates using Sabouraud Dextrose Agar (SDA), Candichrome agar, and Matrix-Assisted Laser Desorption/Ionization Time-of-Flight Mass Spectrometry (MALDI-TOF MS), and to evaluate their demographic and sample-wise distribution among clinical specimens.

Materials and Methods: A hospital-based cross-sectional observational study was conducted in the Department of Microbiology, Dr. S.N. Medical College, Jodhpur, from February 2025 to May 2025. A total of 400 clinical specimens including sputum, urine, blood, endotracheal aspirates (ETA), pus, and cerebrospinal fluid (CSF) were processed. Candida isolates were identified by conventional phenotypic methods and confirmed by MALDI-TOF MS.

Results: Of the 400 clinical samples processed, 74 (18.5%) yielded Candida isolates. Male patients constituted 55.41% of cases, while 44.59% were females. Urban residents accounted for 58.11% of isolates. Most cases were obtained from hospitalized patients (71.62%). The highest frequency of isolation was observed in children ≤ 10 years (33.78%) and adults >60 years (27.02%). Sputum was the most common source of Candida isolates (29.72%), followed by urine and ETA (20.27% each). Phenotypic identification and MALDI-TOF MS revealed Candida albicans as the predominant species (62.16%), followed by Candida tropicalis (29.72%). Less frequently isolated species included Candida kefyr (2.70%), Candida glabrata (2.70%), Candida parapsilosis (1.35%), and Candida krusei (1.35%).

"© 2026 by the Author(s). Published by IJAR under CC BY 4.0. Unrestricted use allowed with credit to the author."

Conclusion: Candida albicans remained the leading species isolated from clinical samples, although non-albicans Candida species constituted a substantial proportion of isolates. MALDI-TOF MS proved to be a rapid and accurate

method for species-level identification and can serve as an effective adjunct to conventional phenotypic techniques. Continuous surveillance of *Candida* species distribution is essential for guiding appropriate antifungal therapy and improving patient outcomes.

Introduction:-

Candida species are ubiquitous fungi that can exist as commensals in the human body but have the potential to cause a range of infections, collectively termed candidiasis. These infections can vary from superficial mucocutaneous conditions to invasive systemic diseases, particularly in immunocompromised individuals¹. The increasing prevalence of candidemia, a bloodstream infection caused by *Candida* species, is a significant concern in healthcare settings due to its association with high morbidity and mortality rates. Studies have shown that candidemia accounts for considerable morbidity and mortality, especially among hospitalized patients in critical care units.^[1] *Candida* species are widespread fungi that commonly exist as commensals within the human microbiota, particularly in the mucosal surfaces of the gastrointestinal and genitourinary tracts.^[2] Although generally harmless, these organisms have the capacity to become opportunistic pathogens under certain conditions, especially in individuals with weakened immune systems. This can result in candidiasis, a spectrum of diseases ranging from superficial mucocutaneous infections to life-threatening systemic conditions such as candidemia, which is a bloodstream infection caused by *Candida* spp.^[2] Candidemia has emerged as a serious healthcare-associated infection, particularly among critically ill and hospitalized patients, due to its high rates of morbidity and mortality. A growing concern in recent years is the shifting epidemiology of candidemia, where non-*albicans* *Candida* (NAC) species such as *C. tropicalis*, *C. glabrata*, *C. parapsilosis*, and *C. krusei* are being isolated with increasing frequency.^[1]

These NAC species often exhibit reduced susceptibility to standard antifungal drugs, complicating treatment protocols and highlighting the need for accurate, species-level identification.^[2] Of particular concern is the emergence of *Candida auris*, a multidrug-resistant yeast capable of causing hospital outbreaks due to its persistence on surfaces and resistance to multiple antifungal classes.^[3] *Candida* infections are increasing in hospital settings, especially among immunocompromised patients due to prolonged use of antibiotic, immunosuppressive therapy, disruption of skin or mucosal barriers, corticosteroids, chronic illnesses, malignancies, organ transplantation, hemodialysis, premature birth, recent surgery, trauma, and the presence of central vascular catheters. In such cases, the non-specific clinical presentation of fungal infections necessitates early consideration of a fungal etiology to improve patient outcomes.^[3] The accurate and timely identification of *Candida* species is paramount for effective patient management.

Different *Candida* species exhibit varying antifungal susceptibility profiles; thus, species-level identification informs appropriate antifungal therapy, which is crucial for improving patient outcomes and reducing mortality^[2]. Traditional identification methods, such as culture-based techniques and biochemical assays, are often time-consuming and may lack the specificity required for precise species differentiation.^[3] To address these diagnostic challenges, Matrix-Assisted Laser Desorption/Ionization Time-of-Flight Mass Spectrometry (MALDI-TOF MS) has revolutionized yeast identification. Unlike conventional culture-based or biochemical techniques, which are time-consuming and less specific, MALDI-TOF MS enables rapid and accurate species-level identification by analyzing the unique protein spectra of yeast-like pathogens.^[3] Its integration into diagnostic workflows, particularly in tertiary care centers, can significantly enhance timely antifungal therapy which is critical in managing candidemia thus reduce mortality. Furthermore, understanding the local epidemiology of *Candida* infections through precise identification aids in monitoring trends, detecting outbreaks, and formulating effective infection control policies.

Material and Methods:-

This study was done for a period of 4 Months i.e. February 2025 to May 2025.

Type of Study: This was a cross-sectional observational hospital-based study.

Sample Size: Total 400 clinical samples were studied.

Source of Data: This study was done in department of microbiology Dr. S.N. Medical College, Jodhpur and their associated groups of hospitals.

Inclusion Criteria:-

1. IPD or OPD patients of all age groups and genders attending tertiary level hospitals, their clinical samples like sputum, Endotracheal aspirate (ETA), Bronchoalveolar lavage (BAL), urine, pleural fluid, blood and CSF of positive fungal culture were included.

Exclusion Criteria:-

1. Patients who were already on anti-fungal treatments.

Sample collection and processing – All clinical samples like blood, urine, CSF, BAL, Pus, other sterile fluids were received in associated groups of hospitals. As per standard laboratory protocol samples were processed and pure growth yeast isolates were further identified and confirmed by MALDI-TOF MS.

Clinical Sample Collection and Processing:-

1. **Blood Samples:** Blood was collected aseptically after disinfecting the venipuncture site with 70% alcohol and povidone-iodine. After drying, 1–3 mL of blood from pediatric and 8–10 mL from adult patients were drawn using sterile syringes and inoculated directly into the BACTEC system bottles for microbial detection.
2. **Urine Samples:** Midstream morning urine was collected in sterile containers. In bedridden patients, urine was drawn aseptically from Foley’s catheter ports using sterile syringes and transferred into 413 labeled sterile containers.
3. **Pus and Swab Samples:** Samples from infected sites like wounds, throat were taken using sterile swabs after surface cleaning to avoid contamination. Specimens were sent to the lab within 1 hour, or refrigerated at 4°C if delayed (not exceeding 24 hours).
4. **Respiratory Samples:** Sputum was collected early morning before meals after rinsing the mouth. Patients coughed deeply into sterile containers, which were properly 413 labeled.
5. **CSF and Pleural Fluid:** CSF was collected in sterile tubes under aseptic conditions and sent immediately. Pleural fluid was aspirated using sterile syringes and placed in anticoagulant-containing tubes for culture.
6. **BAL Fluid:** Bronchoalveolar lavage was collected during bronchoscopy under sterile conditions and transported immediately to the microbiology lab.

Specimen Transport and Storage:-

All clinical samples were transported under controlled conditions immediately after collection to ensure timely and accurate microbiological diagnosis. If immediate processing was not feasible, samples were stored under appropriate conditions to preserve their integrity and prevent contamination or degradation.

Identification Techniques:-

Direct Microscopy:-

Gram Staining:-

Principle:-

Gram staining is based on the ability of bacterial cells to retain certain dyes depending on the structural composition of their cell walls. Gram-positive bacteria possess a thick peptidoglycan layer, which traps the crystal violet-iodine complex, resulting in a purple appearance under the microscope even after alcohol-based decolorization. In contrast, Gram-negative bacteria, with their thin peptidoglycan layer and outer membrane, lose the initial dye during decolorization and take up the counterstain (usually safranin), appearing pink or red. Though originally intended for bacterial identification, Gram staining is also useful in visualizing yeast cells, which generally appear Gram-positive due to the characteristics of their cell walls.

Procedure:

1. Prepare a smear on a slide.
2. Heat-fix over a flame.
3. Stain with crystal violet for 1 minute.
4. Then stain with iodine for 1 minute.
5. Decolorization step - wash the smear with ethanol/acetone for 5-10 seconds.
6. Counterstain with safranin for 1 minute.
7. Rinse the slide let it air dry and examine under oil immersion objectives.

Result Interpretation:-

- Used to visualize **yeasts and pseudohyphae of Candida species.**
- Presence of pseudohyphae indicates tissue invasion this is highly significant finding on direct grams staining
- Yeasts typically appear **Gram-positive** (4–8 µm), but over-decolorization may cause a **false Gram-negative** appearance.

Phenotypic Identification of Yeast Isolates

Culture Media:^[12]:-**SDA (Sabouraud Dextrose Agar):** Supports general fungal growth.

Composition: SDA agar with antibiotic

- Dextrose 40g/L.
- Peptone 10g/L.
- Agar 15g/L.
- pH ~5.6 ±0.2 at 25°C
- Chloramphenicol is added to inhibit bacterial growth.

Culture Procedure:-

1. Inoculate specimen using sterile loop on Sabouraud Dextrose Agar medium
2. Streak gently for isolated colony growth.
3. Incubate aerobically at 25°C or 37° for 24-48 hours.
4. Observe colony characteristics – color, size, texture, **and** pigment.

Species	Colony characteristics of Candida species on SDA agar
Candida albicans	Convex, shiny, creamy
Candida tropicalis	Convex, shiny, creamy
Candida pseudotropicalis	Convex, hard, ivory coloured with mycelia fringe
Candida krusei	Flat, dry
Candida parapsilosis	Wrinkled
Candida guilliermondii	Grayish pink flat, shiny

CandiChrome Agar: Chromogenic medium that differentiates **Candida spp.** Based on colony color.**CandiChrome Agar Testing:-**

Principle: CandiChrome Agar is a fast and efficient plate-based method used for the simultaneous isolation and preliminary identification of different Candida species. This modern medium differentiates Candida species based on distinct colony colors produced through specific biochemical reactions. It enables the identification of multiple Candida types—such as *C. albicans*, *C. krusei*, *C. tropicalis*, *C. glabrata*, *C. parapsilosis*, *C. dubliniensis* and other yeast-like organisms in a single test.

Composition:-

- Peptone.
- Dextrose
- chromogenic substrates
- agar

Procedure:-

1. Inoculate fresh culture isolate on plate using sterile inoculating loop.
2. Streak in zig-zag method on candichrome agar medium
3. Incubate at 25–30°C for 24–48 hours.
4. Observe colony color and morphology to presumptively identify Candida species

Species	Colony Characteristics on Candi Chrome of Candida species
Candida albicans	Apple green colonies, consistent
Candida tropicalis	Dull blue, to purple colour that diffused into surrounding agar plate with pale pink edges.
Candida parapsilosis	White to pale pink colonies
Candida krusei	Large flat, spreading, pale pink colonies with matt surfaces
Candida glabrata	White large glossy pale pink to violet colonies
Candida kefyr	Small pink to purple colonies

Germ Tube Test

Purpose: The germ tube test is a rapid method for identifying **Candida albicans** and distinguishing it from other **Candida** species. When incubated in serum **C. albicans** forms germ tubes which are elongated outgrowths that originate from yeast cells. These structures are an early indication of hyphal development and are considered a characteristic feature of **Candida albicans**.

Procedure:-

1. Suspend yeast colony in 0.5 mL of human/fetal serum.
2. Incubate at 35–37°C for 2–3 hours (no shaking).
3. Prepare a wet mount from yeast serum suspension on a glass slide, examine under 40x objective under microscope.

Result:-

- **Positive:** Germ tubes without basal constriction indicate **Candida albicans**.
- **Negative:** Suggests non albicans **Candida** species.

MALDI-TOF (Matrix-Assisted Laser Desorption/Ionization Time of Flight)^[1]:-

Principle – MALDI-TOF Mass Spectrometry works by mixing the analyte with a matrix compound and applying it to a target plate. A laser beam ionizes the sample, and the resulting ions are accelerated into a drift tube. The time it takes for ions to reach the detector is measured and used to determine their mass-to-charge ratio (m/z), allowing for the identification and characterization of the analytes

Requirements:-

1. ATCC Escherichia coli 8739 strain (as a control) 24 hours fresh sub culture on sheep blood agar plate.
2. CHCA matrix (Cyano-4- hydroxycinnamic acid).
3. Formic acid.
4. Pure and 24 hours fresh Yeast culture isolates.
5. MALDI-TOF Slides.
6. Pipette (0-10 µl).
7. Toothpick for colony picking.

Procedure:-

1. **Sample Preparation:** Pick a yeast colony from an SDA agar plate.
2. **Formic Acid Treatment:** First, add 0.5 µL of 70% formic acid to the yeast colony on the target plate. Allow the formic acid to dry completely. This helps to lyse the cells and release intracellular proteins.
3. **Matrix Application:** After the formic acid dries, add 1-2 µL of the matrix solution, typically α-Cyano-4-hydroxycinnamic acid (CHCA), and let it dry.
4. **Control:-** pick a colony from ATCC E.Coli 8973 strain as a control and make a smear on control spot at center.
5. **Drying and Crystallization:** Let the sample air dry at room temperature. Proper drying will result in the formation of matrix crystals, which is essential for successful analysis.
6. **Loading the Plate:** Insert the prepared target plate into the MALDI-TOF instrument.
7. **Result:** MALDI-TOF then analyses the time of flight and gives the organisms identification report by comparing the time of flight data of analysed isolate with its own database.



Figure 1 – *Candida albicans* SDA Agar



Figure 2 – *Candida* species on Candichrome




Figure 3 - Gram's Stain budding yeast cell



Figure 4 – Germ tube positive



Figure 5 – ATCC *Esch. coli* 8739 strain on sheep blood agar



Figure 6 – MALDI-TOF slide




Figure 7 – Matrix (CHACA) & Formic acid



Figure 8 – Sample processing for MALDI-TOF



Figure 9 – Barcode scanner




Figure 10 – MALDI-TOF Data entry performa



Figure 11 – MALDI-TOF MS



Figure 12 – BOD incubator

Result:-

This is the prospective study for a period of 4 months from February to May 2025. This study was conducted in the Department of Microbiology at Dr. S.N. Medical College & associated group of hospitals in Jodhpur (Rajasthan).

Table-1 Phenotypic identification results of Candida isolates

Total	Positive	%
400	74	18.50

A total of 400 samples were collected and processed for culture during study time period. Out of this 74 (18.50%) samples were positive for Candida isolates, these were studied further.(Table-1)

Chart 1 - Phenotypic identification results of Candida isolates




Table 2: Gender-wise Distribution of Candida isolates

Gender	Number of Cases (N=400)	%
Male	41	55.41
Female	33	44.59
Total	74	100

Table 2 Shown gender wise distribution of Candida isolates. Out of 74 positive isolates, 41(55.41%) were male and 33(44.59%) were female.

Chart 2 - Gender-wise Distribution of Candida isolates




Table 3: Demographic distribution of Candida isolates

Demography	Number of Cases (N=400)	%
Urban	43	58.11
Rural	31	41.89
Total	74	100

Table 3 Shown the demographic distribution of Candida isolates. Out of 74 positive isolates, 43(58.11%) were urban and 31(41.89%) were from rural areas.

Chart 3 -Demographic distribution of Candida isolates




Table 4: OPD and IPD Distribution of Candida isolates

Department	Number of Cases (N=400)	%
OPD	21	28.38
IPD	53	71.62
Total	74	100

As shown in Table 4 out of 74 positive isolates, 21 (23.38) were from OPD and 53(71.62%) were from IPD. This is showing high rate of Candida infection among hospitalized patients.

Chart 4 -OPD and IPD Distribution of Candida isolates




Table 5: Age wise distribution of Candida isolates

Age Group(in years)	Number of Cases(N=30)	%
≤10	25	33.78
11–20	5	6.75
21–30	7	9.45
31–40	5	6.75
41–50	8	10.81
51–60	4	5.40
>60	20	27.02
Total	74	100

Table 5 shown Age wise distribution of Candida isolates the maximum positivity was seen in the age group ≤10years (33.78%) followed by >60 years (27.02)&41-50 years age group (5.40%).

Chart 5 - Age wise distribution of Candida isolates




Table 6: Sample-wise Distribution of Candida Species

Sample type	Total	Positive	%
Sputum	173	22	29.72
Urine	97	15	20.27
Blood	40	11	14.86
ETA	43	15	20.27
Pus	37	10	13.51
CSF	10	1	1.35
Total	400	74	100

Table6 presents the distribution of Candida isolates by Sample wise. Maximum cases were seen in sputum(29.72%) followed by urine (20.27%) & ETA (20.27%), blood (14.86%).

Chart 6 -Sample-wise Distribution of Candida Species




Table 7. Phenotypic identification of Candida by growth on Candichrome agar

Species	Growth Characteristics on Candi-chrome agar	N	%
Candida albicans	Apple green colonies,	46	62.16%
Candida tropicalis	blue, to purple colour colonies	22	29.72%
Candida kefyf	Small pink colonies	2	2.70%
Candida glabrata	Small, pink to mauve colonies	2	2.70%
Candida parapsilosis	White to pale pink colonies	1	1.35%
Candida krusei	Small pink colonies	1	1.35%

This table presents the phenotypic identification of Candida species based on their characteristic growth and color on Candichrome agar. A total of 74 Candida isolates were identified. Candida albicans was the most prevalent species, accounting for 62.16% (46 isolates), and was characterized by apple green colonies. The next most common species was Candida tropicalis, representing 29.72% (22 isolates), which produced blue to purple colonies. Other less frequently identified species included Candida kefyf (2 isolates, 2.70%) and Candida glabrata (2 isolates, 2.70%), both forming small pink to mauve colonies. Candida parapsilosis (1 isolate, 1.35%) displayed white to pale pink colonies, and Candida krusei (1 isolate, 1.35%) also presented as small pink colonies

Chart 7 Phenotypic identification of Candida by growth on Candichrome agar





Table: 8 Species wise distribution of Candida isolates by MALDI-TOF

Species	Candida albicans		Candida tropicalis		Candida kefyfyr		Candida glabrata		Candida parapsilosis		Candida krusei		Total	
	N	%	N	%	N	%	N	%	N	%	N	%	N	%
Sputum	18	81.81%	4	18.18%	0	0%	0	0%	0	0%	0	0%	22	29.72%
Urine	7	46.66%	5	33.33%	1	6.66%	2	13.33%	0	0%	0	0%	15	20.27%
Blood	6	54.54%	4	36.36%	0	0%	0	0%	0	0%	1	9.09%	11	14.86%
ETA	9	60%	5	33.33%	1	6.66%	0	0%	0	0%	0	0%	15	20.27%
Pus	5	50%	4	40%	0	0%	0	0%	1	10%	0	0%	10	13.52%
CSF	1	100%	0	0%	0	0%	0	0%	0	0%	0	0%	1	1.36%
Total	46		22		2		2		1		1		74	100%

This table provides a detailed breakdown of Candida species distribution across various clinical sample types, identified using MALDI-TOF. A total of 74 Candida isolates were analyzed. Candida albicans was the predominant species across most sample types, representing 81.81% (18 isolates) in sputum, 60% (9 isolates) in ETA, and 100% (1 isolate) in CSF. It also formed a significant proportion in blood (54.54%, 6 isolates) and pus (50%, 5 isolates). Candida tropicalis was the second most common species, showing notable presence in sputum (18.18%, 4 isolates), urine (33.33%, 5 isolates), blood (36.36%, 4 isolates), ETA (33.33%, 5 isolates), and pus (40%, 4 isolates). Other Candida species were detected in smaller numbers. Candida kefyfyr was found in urine (1 isolate, 6.66%) and ETA (1 isolate, 6.66%). Candida glabrata was identified in urine (2 isolates, 13.33%). Candida parapsilosis was present in pus (1 isolate, 10%), and Candida krusei was detected in blood (1 isolate, 9.09%). Overall, sputum samples yielded the highest number of isolates (22, 29.72%), followed by urine (15, 20.27%) and ETA (15, 20.27%). Blood samples accounted for 11 isolates (14.86%), while pus samples contributed 10 isolates (13.52%). A single isolate was found in CSF (1.36%).

Chart 8 -Species wise distribution of Candida isolates by MALDI-TOF



Discussion:-

The present cross-sectional study, conducted at Dr. S.N. Medical College, Jodhpur, aimed to investigate the phenotypic identification and species distribution of Candida isolates using both conventional methods and MALDI-TOF MS. This analysis sheds light on the evolving epidemiology of candidiasis, particularly in Western Rajasthan. Out of 400 clinical samples, 74 cases (18.5%) tested positive for Candida infections (Table 1). However, other studies like Agarwal et al. [8] found positivity rate of 25% and Singh et al. [25] recorded lower prevalence rate of between 10% and 20%. The variation in the finding could be influenced by factors such as local healthcare practices, environmental conditions, and population vulnerability.

Gender wise Distribution of Candida isolates:-

The gender distribution in this study showed a male predominance (55.41%), aligning with the findings of Rajni et al., who suggested that males may have greater exposure to risk factors like catheterization or underlying comorbidities such as diabetes [2]. Urban cases (58.11%) were more prevalent than rural ones (41.89%), possibly due to environmental and occupational exposure, our results emphasize the need for awareness and public health strategies in both urban and rural populations to effectively address the fungal disease burden.

IPD and OPD wise distribution of Candida isolates:

A significant majority (71.62%) of Candida-positive cases were detected in hospitalized (IPD) patients, affirming findings by Rudramurthy et al. and Mellinghoff et al., who documented elevated candidiasis prevalence among ICU and hospitalized populations due to immunosuppressive therapies, invasive procedures, and prolonged hospital stays [2,4]. Such patients are often subject to multiple risk factors, including prior antibiotic use, steroid therapy, and the presence of indwelling medical devices, which facilitate Candida colonization and infection.

Age group wise distribution of Candida isolates:-

A bimodal age distribution was noted, with the highest incidence in the pediatric age group (<10 years, 33.78%) and older adults (>60 years, 27.02%). While candidiasis is typically more common in the elderly due to immunocompromise status and chronic illness, the spike in pediatric cases may reflect heightened diagnostic scrutiny or underlying pediatric conditions. Similar age-related vulnerability was noted in studies by Agarwal and Chakrabarti,^[2] who reported increased incidence among older adults.^[5]

Sample-wise Distribution of Candida Isolates:-

Table 6 highlights the distribution of Candida isolates across different sample types. Out of 400 total clinical samples, 74 (18.5%) were positive for Candida. Sputum samples yielded the highest positivity rate (29.72%, 22 isolates), followed by urine and ETA (Endotracheal Aspirate) samples, both contributing 20.27% (15 isolates each). Blood samples accounted for 14.86% (11 isolates), while pus samples comprised 13.51% (10 isolates). Notably, only one Candida isolate (1.35%) was detected from CSF. This distribution concordant with findings from other studies in tertiary care settings where respiratory tract samples (sputum, ETA) and urinary samples are frequently implicated in candidiasis, often due to colonization in critically ill or catheterized patients^[41,43,29]. The presence of Candida in blood and CSF, though less frequent in our study, signifies invasive candidiasis, which is associated with high morbidity and mortality^[2,18,32].

Phenotypic Identification of Candida Species:-

Table 7 illustrates the phenotypic identification of Candida species using Candichrome agar. This chromogenic medium allows for presumptive identification based on distinct color production, which is a valuable tool for rapid differentiation in a clinical microbiology laboratory^[12]. Our study identified *Candida albicans* as the most prevalent species (62.16%, 46 isolates), characterized by its typical apple green colonies. This finding is consistent with global epidemiology, where *C. albicans* traditionally remains the most common Candida species causing human infections.^[9,10,11] *Candida tropicalis* was the second most common species (29.72%, 22 isolates), identified by its blue to purple colonies. The increasing prevalence of non-*albicans* Candida (NAC) species, including *C. tropicalis*, *C. glabrata*, and *C. parapsilosis*, has been a growing concern in recent years, as they often exhibit reduced susceptibility to commonly used antifungals^[1,5,17]. The detection of *C. kefyr* (2.70%), *C. glabrata* (2.70%), *C. parapsilosis* (1.35%), and *C. krusei* (1.35%) in our study, although in smaller numbers, underscores the importance of accurately identifying these species for guiding appropriate antifungal therapy.^[5,6]

Species-wise Distribution of Candida Isolates by MALDI-TOF:

Table 8 provides a more definitive species-level identification using Matrix-Assisted Laser Desorption/Ionization-Time of Flight Mass Spectrometry (MALDI-TOF MS). MALDI-TOF is a rapid, accurate, and cost-effective method for identifying microorganisms, significantly improving the turnaround time for pathogen identification compared to traditional methods.^[1,3] Our MALDI-TOF results largely corroborated the phenotypic identifications from Candichrome agar, while providing a more precise breakdown of species distribution across different sample types. Consistent with the overall prevalence, *Candida albicans* was the dominant species in sputum (81.81%) and ETA (60%) samples, reinforcing its role as a significant colonizer and pathogen in the respiratory tract [9]. Its 100% presence in the single CSF isolate is clinically relevant, indicating its potential to cause severe central nervous system infections. In urine samples, *C. albicans* was still the most common (46.66%), but *C. tropicalis* (33.33%) and *C. glabrata* (13.33%) also contributed significantly, reflecting the varied etiology of candiduria [41, 42].

The distribution in blood samples is particularly important given the severity of candidemia. Here, *C. albicans* (54.54%) and *C. tropicalis* (36.36%) were the primary isolates. While *C. albicans* is traditionally the leading cause of candidemia, the high proportion of *C. tropicalis* aligns with global trends showing an increasing incidence of *C. tropicalis* candidemia, which is often associated with higher mortality rates in some settings^[1,2,5]. The detection of *Candida krusei* (9.09%) in blood, though only one isolate, is notable because this species is intrinsically resistant to fluconazole, a commonly used antifungal^[13,17]. Similarly, *C. parapsilosis* in pus samples highlights its potential role in wound infections and device-related infections^[3,10]. This study underscores the critical need for species-level identification in managing candidiasis, especially given the varying antifungal susceptibility profiles of NAC species. Timely identification supports appropriate antifungal therapy and resistance prevention.

These observations are in line with recommendations by Tajaneet al.^[4] and Tiwari & Verma,^[6] who advocate for integrated diagnostics combining phenotypic and molecular tools for effective fungal management. The findings from our study are crucial for developing effective antifungal stewardship programs and improving patient outcomes in our tertiary care hospital. The prevalence of *C. albicans* reaffirms the need for continued surveillance and

appropriate treatment protocols for this species. However, the significant presence of NAC species, particularly *C. tropicalis*, necessitates vigilant identification and antifungal susceptibility testing to guide targeted therapy, especially considering their varying susceptibility profiles.^[6,13]

The application of advanced diagnostic techniques like MALDI-TOF MS in routine laboratory practice provides rapid and accurate identification, facilitating timely intervention and reducing the reliance on empirical antifungal therapy^[1,3]. This is particularly important in managing critically ill patients with underlying risk factors such as diabetes mellitus, immunosuppression (e.g., HIV, organ transplantation), and malignancy, who are highly susceptible to invasive candidiasis^[26, 27, 29, 30, 33, 36, 38, 39, 40]. Furthermore, the emerging threat of multidrug-resistant *Candida* species, such as *Candida auris*, though not detected in our study, emphasizes the need for ongoing surveillance and robust infection control measures^[15,18,25].

Summary:-

This cross-sectional study was conducted over a four-month period at Dr. S.N. Medical College, Jodhpur, with the aim of investigating the phenotypic identification and species distribution of *Candida* isolates from various clinical samples. Out of 400 processed samples, 74 (18.5%) tested positive for *Candida*, with the most frequent isolates found in sputum and urine samples. The use of both conventional culture methods and advanced techniques like MALDI-TOF MS allowed accurate species-level identification. A total of 74 *Candida* isolates were identified. *Candida albicans* was the predominant species, accounting for 46 (62.16%) isolates, followed by *Candida tropicalis* with 22 (29.72%) isolates. Less frequently isolated species included *Candida kefyr* and *Candida glabrata* (2.70% each), while *Candida parapsilosis* and *Candida krusei* accounted for 1.35% each. This reflects an epidemiological transition from *C. albicans* dominance to the rise of NAC species, emphasizing the need for precise diagnostic tools. The gender distribution showed a slight male predominance (55.41%), while urban cases slightly outnumbered rural ones (58.11% vs. 41.89%), suggesting more frequent healthcare access or reporting in urban regions.

Age-wise, children aged 0–10 years (33.78%) and the elderly population above 60 years (27.02%) were most affected, indicating bimodal vulnerability due to immune immaturity and immunosenescence, respectively. The study also found a much higher infection rate among hospitalized patients (71.62%) compared to outpatients, underlining the risk factors linked to hospital settings such as catheterization, ICU stays, and immunosuppression. The application of MALDI-TOF MS significantly improved the speed and accuracy of species-level identification, especially for NAC isolates. This reinforces the importance of integrating advanced diagnostic modalities alongside traditional phenotypic methods to facilitate prompt and appropriate antifungal treatment, particularly in tertiary care setups.

Conclusion:-

This study provides valuable insights into the epidemiological profile of *Candida* infections in a tertiary care setting. The data underscores the continued dominance of *C. albicans* while highlighting the significant contribution of NAC species, particularly *C. tropicalis*, across various clinical samples. The use of advanced identification methods like MALDI-TOF is indispensable for accurate and timely diagnosis, which is paramount for effective antifungal management and combating the evolving challenge of candidiasis in healthcare. Continued surveillance and research are vital to monitor changes in *Candida* epidemiology and antifungal resistance patterns to inform clinical guidelines and improve patient care.

Limitation of study:-

In this present study only identification of *Candida* isolates was done. Antifungal susceptibility test for *Candida* isolates was not done.

References:-

1. K, Shetty AK, Antony B. MALDI-TOF MS Analysis to Detect the Prevalence of Non-albicans Candidemia with a Special Emphasis on *Candida auris* from a Tertiary Care Centre in Coastal Karnataka. Online J Health Allied Scs. 2024;23(2):9
2. Rajni E, Chaudhary P, Garg VK, Sharma R, Malik M. A complete clinico-epidemiological and microbiological profile of candidemia cases in a tertiary-care hospital in Western India. Antimicrobe Steward Healthcare Epidemiolgy. 2022 Mar 7;2(1):e37
3. Samantaray S, Singh R. Evaluation of MALDI-TOF MS for Identification of Species in the *Candida parapsilosis* Complex from Candidiasis Cases. J Appl Lab Med. 2022 Jun 30;7(4):889-900

4. Tajane SB, Pawar S, Patil S. Revisiting the History of Candidiasis. *Cureus*. 2025 Feb 11;17(2):e78878
5. M A, JD J (February 25, 2025) Species Distribution and Antifungal Susceptibility Patterns of Candida Isolates: A Cross-Sectional Study From a Tertiary Care Hospital in South India. *Cureus* 17(2)
6. Tiwari, K., & Verma, A. K. (2020). Antifungal susceptibility pattern of clinical isolates of Candida from a tertiary care hospital in Chhattisgarh, India. *ResearchGate*.
7. Yaguchi T, Tanaka R. [Classification and phylogeny of pathogenic fungi]. *Nihon Rinsho*. 2008 Dec;66(12):2261-7
8. Schauer F, Hanschke R. Zur Taxonomie und Ökologie der Gattung Candida [Taxonomy and ecology of the genus Candida]. *Mycoses*. 1999;42 Suppl 1:12-21. German
9. Christina Tsui, Eric F. Kong, Mary Ann Jabra-Rizk, Pathogenesis of Candida albicans biofilm, *Pathogens and Disease*, Volume 74, Issue 4, June 2016, ftw018
10. utler, G., Rasmussen, M., Lin, M. et al. Evolution of pathogenicity and sexual reproduction in eight Candida genomes. *Nature* 459,657(2009)
11. Lopes, J. P., & Lionakis, M. S. (2021). Pathogenesis and virulence of Candida albicans. *Virulence*, 13(1), 89–121.
12. Medina N, Soto-Debrán JC, Seidel D, Akyar I, Badali H, Barac A, et al. MixInYeast: A Multicenter Study on Mixed Yeast Infections. *J Fungi (Basel)*. 2020 Dec 29;7(1):13.
13. Chander J. textbook of medical mycology. 4th edition. New delhi : Mehta publisher;2008 ASDCV.
14. Clinical and Laboratory Standards Institute (CLSI). Method for Antifungal Disk Diffusion Susceptibility Testing of Yeasts; Approved Guideline. CLSI document M44. Wayne, PA: Clinical and Laboratory Standards Institute; 2022.
15. Hoeningl M, Gangneux JP, Segal E, Alanio A, Chakrabarti A, Chen SCA, et al. Global guidelines and initiatives from the European Confederation of Medical Mycology to improve patient care and research worldwide: New leadership is about working together. *Mycoses*. 2018 Nov;61(11):885-94
16. Dorko E, Baranová Z, Jenca A, Kizek P, Pilipcinec E, Tkáčiková L. Diabetes mellitus and candidiasis. *Folia Microbiol (Praha)*. 2005;50(3):255-61.
17. Gonçalves SS, Souza ACR, Chowdhary A, Meis JF, Colombo AL. Epidemiology and molecular mechanisms of antifungal resistance in Candida and Aspergillus. *Mycoses*. 2016 Apr;59(4):198-219.
18. Mathur P, Hasan F, Singh PK, Malhotra R, Walia K, Chowdhary A. Five-year profile of candidaemia at an Indian trauma centre: High rates of Candida auris blood stream infections. *Mycoses*. 2018 Sep;61(9):674-80.
19. Prakash A, Randhawa HS, Khan ZU, Ahmad S, Hagen F, Meis JF, et al. Environmental distribution of Cryptococcus species and some other yeast-like fungi in India. *Mycoses*. 2018 May;61(5):305-13.
20. Chow NA, Muñoz JF, Gade L, Berkow EL, Li X, Welsh RM, et al. Tracing the Evolutionary History and Global Expansion of Candida auris Using Population Genomic Analyses. *mBio*. 2020 Apr 28;11(2):e03364-19.
21. Chowdhary A, Sharma C, Meis JF. Azole-Resistant Aspergillosis: Epidemiology, Molecular Mechanisms, and Treatment. *J Infect Dis*. 2017 Aug 15;216(suppl_3):S436-44.
22. Kordalewska M, Zhao Y, Lockhart SR, Chowdhary A, Berrio I, Perlin DS. Rapid and Accurate Molecular Identification of the Emerging Multidrug-Resistant Pathogen Candida auris. *J Clin Microbiol*. 2017 Aug;55(8):2445-52.
23. Mellinshoff SC, Hartmann P, Cornely FB, Knauth L, Köhler F, Köhler P, et al. Analyzing candidemia guideline adherence identifies opportunities for antifungal stewardship. *Eur J Clin Microbiol Infect Dis* 2018 Aug;37(8):1563-71
24. Kanj SS, Haddad SF, Meis JF, Verweij PE, Voss A, Rautemaa-Richardson R, et al. The battle against fungi: lessons in antifungal stewardship from COVID 19 times. *Int J Antimicrob Agents*. 2023 Jul;62(1):106846.
25. Chowdhary A, Sharma C, Meis JF. Candida auris: A rapidly emerging cause of hospital-acquired multidrug-resistant fungal infections globally. *PLoS Pathog*. 2017 May; 13(5):e1006290.
26. Chowdhary A, Gupta N, Wurster S, Kumar R, Mohabir JT, Tatarvarthy S, et al. Multimodal analysis of the COVID-19-associated mucormycosis outbreak in Delhi, India indicates the convergence of clinical and environmental risk factors. *Mycoses*, 2023 Jun;66(6):515-26.
27. Banerjee S, Denning DW, Chakrabarti A. One Health aspects & priority roadmap for fungal diseases: A mini-review. *Indian J Med Res*. 2021 Mar, 153(3):311-9
28. Lanternier F, Lortholary O. Zygomycosis and diabetes mellitus. *Clin Microbiol Infect*. 2009 Oct;15 Suppl 5:21-5.
29. Rawson TM, Moore LSP, Zhu N, Ranganathan N, Skolimowska K, Gilchrist M, et al. Bacterial and Fungal Coinfection in Individuals With Coronavirus: A Rapid Review To Support COVID-19 Antimicrobial Prescribing. *Clin Infect Dis*. 2020 Dec 3;71(9):2459-74.

30. Thomas-Rüddel DO, Schlattmann P, Pletz M, Kurzai O, Bloos F. Risk Factors for Invasive Candida Infection in Critically Ill Patients: A Systematic Review and Meta-analysis. *Chest*. 2022 Feb;161(2):345-55.
31. Chukwuanukwu RC, Uchenna N, Mbagwu SI, Chukwuanukwu TO, Charles O. Cryptococcus neoformans seropositivity and some haematological parameters in HIV seropositive subjects. *J Infect Public Health*. 2020 Jul;13(7):1042-6.
32. Chesdachai S, Yetmar ZA, Lahr BD, Vege SS, Vergidis P. Clinical characteristics and outcomes of pancreatic fungal infection in patients with necrotizing pancreatitis. *Med Mycol*. 2023 Jul 6;61(7):myad068.
33. Thompson GR, Jenks JD, Baddley JW, Lewis JS, Egger M, Schwartz IS, et al. Fungal Endocarditis: Pathophysiology, Epidemiology, Clinical Presentation, Diagnosis, and Management. *Clin Microbiol Rev*. 2023 Sep 21;36(3):20001923.
34. Richardson MD, Kokki MH. Diagnosis and prevention of fungal infection in the immunocompromized patient. *Blood Rev*. 1998 Dec;12(4):241-54.
35. Cornely OA, Maertens J, Winston DJ, Perfect J, Ullmann AJ, Walsh TJ, et al. Posaconazole vs. fluconazole or itraconazole prophylaxis in patients with neutropenia. *N Engl J Med*. 2007 Jan 25;356(4):348-59.
36. 106. Jørgensen KJ, Gøtzsche PC, Johansen HK. Voriconazole versus amphotericin B in cancer patients with neutropenia. *Cochrane Database Syst Rev*. 2006 Jan 25;(1):CD004707.
37. Tolkoff-Rubin NE, Rubin RH. Opportunistic fungal and bacterial infection in the renal transplant recipient. *J Am Soc Nephrol*. 1992 Jun;2(12 Suppl):S264-269.
38. Al Salmi I, Metry AM, Al Ismaili F, Hola A, Al Riyami M, Khamis F. et al. Transplant tourism and invasive fungal infection. *Int J Infect Dis*. 2018 Apr 69:120-9.
39. Kim L, Ferraz C, Corbisiero MF, Gorvetzian S, Franco-Paredes C, Krak M. et al. Glucocorticoids as a risk factor for infection and adverse outcomes in non-HIV and non-transplant patients with cryptococcal meningitis. *Mycoses*, 2024 Mar 67(3)ke13709
40. Chen CY, Sheng WH, Tien FM, Lee PC, Huang SY, Tang L, et al Clinical characteristics and treatment outcomes of pulmonary invasive fungal infection among adult patients with hematological malignancy in a medical centre in Taiwan, 2008-2013. *J Microbiol Immunol Infect*. 2020 Feb;53(1):106-14.
41. Singh GK, Capoor MR, Nair D, Bhowmik KT. Spectrum of fungal infection in head and neck cancer patients on chemoradiotherapy. *J Egypt Natl Canc Inst*. 2017 Mar;29(1):33-7.
42. Odabasi Z, Mert A. Candida urinary tract infections in adults. *World J Urol*. 2020 Nov;38(11):2699-707.
43. Malani AN, Kauffman CA. Candida urinary tract infections: treatment options. *Expert Rev Anti Infect Ther*. 2007 Apr;5(2):277-84.
44. Kim J, Kim DS, Lee YS, Choi NG. Fungal urinary tract infection in burn patients with long-term foley catheterization. *Korean J Urol*. 2011 Sep;52(9):626-31.
45. Thomas L, Tracy CR. Treatment of Fungal Urinary Tract Infection. *Urol Clin North Am*. 2015 Nov;42(4):473-83.
46. Maseda E, Rodríguez AH, Aguilar G, Pemán J, Zaragoza R, Ferrer R, et al. EPICO 3.0. Recommendations on invasive candidiasis in patients with complicated intra-abdominal infection and surgical patients with ICU extended stay. *Rev IberoamMicol*. 2016;33(4):196-205.
47. Vandewoude KH, Blot SI, Benoit D, Colardyn F, Vogelaers D. Invasive aspergillosis in critically ill patients: attributable mortality and excesses in length of ICU stay and ventilator dependence. *J Hosp Infect*. 2004 Apr;56(4):269-76.
48. Fróes LAR, Toma TS, Jachiet M, Rousset L, Poderoso RE, Trindade MAB. Bacterial, fungal and parasitic co-infections in leprosy: A scoping review. *PLoS Negl Trop Dis*. 2023 May; 17(5):e0011334.