



ISSN (O): 2320-5407  
ISSN (P): 3107-4928

Journal Homepage: [-www.journalijar.com](http://www.journalijar.com)

## INTERNATIONAL JOURNAL OF ADVANCED RESEARCH (IJAR)

Article DOI: 10.21474/IJAR01/23689  
DOI URL: <http://dx.doi.org/10.21474/IJAR01/23689>



INTERNATIONAL JOURNAL OF  
ADVANCED RESEARCH (IJAR)  
ISSN 2320-5407  
Journal homepage: <http://www.journalijar.com>  
Journal DOI: 10.21474/IJAR01

### RESEARCH ARTICLE

## ELASTOGRAPHY IN OVARIAN IMAGING: EMERGING APPLICATIONS IN PCOS AND PCOM

Suvarchala Govada<sup>1</sup> and Badugu Saikumar<sup>2</sup>

1. Professor, Department of Obstetrics & Gynaecology, Tagore Medical College, Chennai.
2. Assistant Professor, Department of Anaesthesiology, KIMS Hospitals, Secunderabad.

#### Manuscript Info

##### Manuscript History

Received: 14 April 2026  
Final Accepted: 16 May 2026  
Published: June 2026

##### Key words:-

Artificial intelligence, elastography, infertility, ovarian imaging, ovarian stiffness, polycystic ovarian morphology, polycystic ovary syndrome, reproductive ultrasound, shear-wave elastography, strain elastography, stromal fibrosis, ultrasonography.

#### Abstract

Polycystic ovary syndrome (PCOS) is one of the most common endocrine and metabolic disorders affecting women of reproductive age and is frequently associated with infertility, menstrual irregularities, hyperandrogenism, obesity, and insulin resistance. Polycystic ovarian morphology (PCOM), characterized by increased follicle number and enlarged ovarian volume, represents the characteristic imaging appearance of the disorder. Conventional ultrasonography remains the primary imaging modality for ovarian evaluation; however, it provides limited information regarding stromal tissue composition, fibrosis, and biomechanical properties. In recent years, elastography has emerged as an advanced ultrasound-based technique capable of assessing tissue stiffness and elasticity in real time, thereby offering additional functional information beyond routine morphological assessment. Both strain elastography and shear wave elastography (SWE) have demonstrated promising applications in ovarian imaging. Studies have shown that ovaries affected by PCOS and PCOM exhibit increased stromal stiffness due to stromal hypertrophy, collagen deposition, fibrosis, hyperthecosis, and chronic low grade inflammation. Elastographic findings have also shown correlation with hormonal abnormalities such as elevated androgen levels, increased anti-Müllerian hormone (AMH), insulin resistance, and metabolic syndrome. Furthermore, elastography may aid in differentiating PCOM from multifollicular ovaries, evaluating infertility, predicting ovarian response during assisted reproductive therapy, and monitoring treatment outcomes. This review highlights the principles and techniques of ovarian elastography, pathophysiological basis of altered ovarian stiffness, imaging findings in PCOS and PCOM, clinical applications, current evidence, advantages, limitations, and future directions including artificial intelligence integration and multiparametric ultrasound approaches in reproductive imaging.

"© 2026 by the Author(s). Published by IJAR under CC BY 4.0. Unrestricted use allowed with credit to the author."


### Introduction:-

Polycystic ovary syndrome (pcos) is one of the most common endocrine and metabolic disorders affecting women of reproductive age worldwide, with a reported prevalence ranging from 6% to 20% depending on the diagnostic criteria used.<sup>1,7</sup> It is a heterogeneous disorder characterized by chronic anovulation, hyperandrogenism, menstrual irregularities, infertility, obesity, insulin resistance, and metabolic dysfunction.<sup>6,8</sup> In addition to reproductive abnormalities, pcOS is associated with long-term complications such as type 2 diabetes mellitus, cardiovascular disease, dyslipidemia, and psychological disorders, making it a major public health concern.<sup>3,17</sup> Polycystic ovarian morphology (pcom) represents the characteristic ovarian appearance seen on imaging in many women with pcOS. According to the revised Rotterdam criteria, pcom is defined by the presence of 20 or more follicles measuring 2–9 mm in diameter in at least one ovary and/or an ovarian volume greater than 10 ml when assessed using high-resolution transvaginal ultrasonography.<sup>1,4</sup> Ultrasound examination plays a central role in the diagnosis and evaluation of ovarian morphology because it is non-invasive, widely available, cost-effective, and capable of assessing follicular distribution and stromal echogenicity.<sup>5,20</sup> Conventional grayscale and Doppler ultrasonography are routinely used to evaluate ovarian size, follicle number, and vascularity. However, these imaging modalities provide limited information regarding tissue biomechanics, stromal fibrosis, and ovarian stiffness.<sup>10,11</sup> Increasing evidence suggests that stromal hypertrophy, collagen deposition, fibrosis, and hyperthecosis contribute significantly to the pathophysiology of pcOS and may correlate with endocrine and metabolic abnormalities.<sup>8,19</sup>

Elastography is an advanced ultrasound-based imaging technique that evaluates tissue stiffness and elasticity by measuring tissue displacement or shear-wave propagation following mechanical stress.<sup>10,11</sup> Initially developed for the assessment of liver fibrosis, elastography has rapidly expanded into various clinical applications including breast, thyroid, prostate, musculoskeletal, and gynaecologic imaging.<sup>12</sup> Recent advances in strain elastography and shear-wave elastography (SWE) have enabled non-invasive quantitative assessment of ovarian stromal stiffness in women with PCOS and PCOM.<sup>13,14</sup> Emerging evidence indicates that ovaries affected by PCOS demonstrate significantly increased stiffness compared with normal ovaries due to stromal fibrosis, increased connective tissue, and hormonal alterations.<sup>13,15</sup> Elastography may therefore serve as a valuable adjunct to conventional ultrasound by improving diagnostic accuracy, differentiating PCOM from multi-follicular ovaries, assessing disease severity, correlating with hormonal and metabolic parameters, and monitoring therapeutic response.<sup>14,15</sup> This review aims to discuss the principles of ovarian elastography, imaging techniques, pathophysiological basis of altered ovarian stiffness in PCOS and PCOM, current clinical applications, research evidence, limitations, and future perspectives in reproductive imaging.



**Figure 1: Transvaginal ultrasound images showing polycystic ovarian morphology with multiple peripherally arranged follicles in the right ovary.**



**Figure 2: Ovarian elastography images demonstrating increased stromal stiffness in polycystic ovaries, with elasticity measurements expressed in kilopascals (kPa).**

**Anatomy and Physiology of the Ovary:** The ovaries are paired female reproductive organs responsible for oocyte production and secretion of hormones such as estrogen, progesterone, and androgens.<sup>21,22</sup> Structurally, the ovary consists of an outer cortex containing developing follicles and stromal tissue, and an inner medulla containing blood vessels, lymphatics, nerves, and connective tissue.<sup>21,23</sup> The ovarian stroma is composed of fibroblasts, collagen fibers, and extracellular matrix that provide structural and hormonal support.<sup>22,23</sup>

**Folliculogenesis is the process of follicular maturation and includes:**<sup>23,24</sup>


- Primordial follicle recruitment
- Primary follicle development
- Secondary follicle formation
- Graafian follicle maturation
- Ovulation

These processes are regulated by the hypothalamic-pituitary-ovarian axis through hormones such as FSH and LH.<sup>23</sup> During ovulation, the mature Graafian follicle ruptures and releases the oocyte, followed by formation of the corpus luteum.<sup>21,23</sup> In polycystic ovary syndrome (PCOS), normal follicular maturation is disrupted due to hyperandrogenism, insulin resistance, and hormonal imbalance.<sup>1,6</sup> This leads to follicular arrest, accumulation of immature follicles, stromal hypertrophy, fibrosis, and enlarged ovaries.<sup>8,19</sup> These pathological stromal changes increase ovarian stiffness, which can be assessed using elastography in ovarian imaging.<sup>11, 10,13</sup>

**Pathophysiology:** Polycystic ovary syndrome (PCOS) is a complex endocrine and metabolic disorder characterized by hormonal imbalance, insulin resistance, abnormal follicular development, and structural ovarian changes.<sup>2,6</sup> The disorder develops due to interactions between genetic, environmental, metabolic, and endocrine factors that alter the normal hypothalamic-pituitary-ovarian axis.<sup>6,8</sup> One of the hallmark abnormalities in PCOS is hyperandrogenism, which may be clinical or biochemical. Clinical manifestations include hirsutism, acne, alopecia, seborrhea, and menstrual irregularities.<sup>2,17</sup> Increased luteinizing hormone (LH) secretion stimulates ovarian theca cells to produce excess androgens, while insulin further enhances androgen synthesis.<sup>6,8</sup> Elevated androgen levels disrupt follicular maturation and inhibit ovulation, resulting in chronic anovulation.<sup>6,3</sup> Insulin resistance is another major feature of PCOS and occurs in both obese and lean women.<sup>8,3</sup> Peripheral tissues such as skeletal muscle, adipose tissue, and liver become less responsive to insulin, leading to compensatory hyperinsulinemia.<sup>8,27</sup> Hyperinsulinemia increases ovarian androgen production and decreases hepatic synthesis of sex hormone-binding globulin (SHBG), thereby increasing free circulating androgen levels.<sup>2,19</sup> These hormonal abnormalities impair dominant follicle selection and lead to accumulation of immature follicles within enlarged ovaries.<sup>6,19</sup>

Abnormal folliculogenesis in PCOS is characterized by arrest of follicular development at the small antral stage.<sup>6,8</sup> Multiple immature follicles accumulate peripherally within the ovary, producing the classic “string of pearls” appearance on ultrasound.<sup>1,26</sup> Chronic anovulation contributes to infertility and menstrual dysfunction.<sup>19,33</sup> The ovarian stroma undergoes significant structural remodelling in PCOS. Stromal hypertrophy, stromal hyperplasia, increased connective tissue deposition, collagen accumulation, and fibrosis contribute to increased ovarian volume and stromal echogenicity.<sup>8,19</sup> Thickening of the tunica albuginea further impairs follicular rupture and ovulation.<sup>6,28</sup> Hyperthecosis and chronic low-grade inflammation additionally contribute to stromal rigidity and fibrosis.<sup>8,18</sup> Doppler ultrasonography demonstrates increased ovarian vascularity, stromal blood flow, and angiogenesis in women with PCOS.<sup>26,28</sup> These vascular and stromal alterations form the biological basis for elastographic imaging, which evaluates ovarian tissue stiffness and fibrosis.<sup>19,11</sup> Elastography has therefore emerged as a promising imaging modality for assessing ovarian stromal changes in PCOS and polycystic ovarian morphology (PCOM).<sup>11</sup>

## PATHOPHYSIOLOGY OF PCOS



**Figure 03: This diagram effectively demonstrates how endocrine dysfunction, insulin resistance, inflammation, and ovarian structural remodelling together contribute to the development and progression of PCOS.**

**Introduction to Elastography in Ovarian Imaging:** Elastography is an advanced ultrasound imaging technique that assesses tissue stiffness and elasticity. Unlike conventional ultrasonography, which mainly depicts anatomical structures, elastography provides additional functional and biomechanical information about tissue composition, fibrosis, and stromal architecture. This technique has emerged as a valuable tool in gynaecologic imaging, particularly in the assessment of ovarian disorders such as Polycystic Ovary Syndrome (PCOS) and polycystic ovarian morphology (PCOM).<sup>10,29</sup> In PCOS and PCOM, ovarian stromal hypertrophy, fibrosis, and increased

collagen deposition lead to alterations in tissue elasticity, resulting in increased ovarian stiffness.<sup>30,31</sup> Elastography enables non-invasive evaluation of these stromal changes, thereby improving diagnostic accuracy and enhancing understanding of ovarian pathophysiology.<sup>32</sup>

**Currently, two major elastographic techniques are widely used in gynaecologic and ovarian imaging:-**

**1. Strain Elastography:** Strain elastography is a qualitative or semi-quantitative technique that evaluates tissue deformation after external compression is applied using the ultrasound transducer. Softer tissues deform more easily, whereas stiffer tissues show less deformation. The resulting elastogram provides a color-coded representation of tissue stiffness, helping assess ovarian stromal consistency.<sup>33</sup>

**2. Shear-Wave Elastography (SWE):** Shear-wave elastography is a quantitative imaging modality that uses acoustic radiation force impulses to generate shear waves within tissues. The velocity of these shear waves is measured to calculate tissue stiffness objectively, usually expressed in kilopascals (kPa) or meters per second (m/s). SWE offers greater reproducibility and reduced operator dependency compared with strain elastography.<sup>34,11</sup> Both elastographic methods contribute significantly to the evaluation of ovarian stromal abnormalities in PCOS and PCOM. By detecting increased ovarian stiffness associated with stromal fibrosis and hypertrophy, elastography may serve as a promising adjunctive tool in the diagnosis, characterization, and monitoring of ovarian disorders.<sup>29,32,11</sup>

#### Types of Elastography in Ovarian Imaging


| Aspect                           | Strain Elastography (SE)                                                                                                               | Shear-Wave Elastography (SWE)                                                                                                   |
|----------------------------------|----------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------|
| <b>Definition</b>                | A qualitative or semi-quantitative elastographic technique that evaluates tissue deformation after external compression. <sup>35</sup> | A quantitative elastographic technique that measures tissue stiffness using acoustic radiation force impulses. <sup>34,11</sup> |
| <b>Basic Principle</b>           | Tissue deformation occurs when manual pressure is applied; softer tissues deform more than hard tissues.                               | Acoustic pulses generate shear waves inside tissue; wave velocity reflects tissue stiffness. <sup>34</sup>                      |
| <b>Physical Basis</b>            | Based on tissue strain under compression. <sup>35</sup>                                                                                | Based on shear-wave propagation velocity and Young's modulus. <sup>11</sup>                                                     |
| <b>Key Formula</b>               | $\sigma = E\epsilon^2$                                                                                                                 | $E = 3\rho c_s^2$ <sup>24</sup>                                                                                                 |
| <b>Method of Compression</b>     | Manual compression using transvaginal probe.                                                                                           | No manual compression required. <sup>36</sup>                                                                                   |
| <b>Imaging Technique</b>         | Compression–decompression cycles create elastograms. <sup>35</sup>                                                                     | Focused ultrasound pulses create transverse shear waves. <sup>34</sup>                                                          |
| <b>Type of Assessment</b>        | Qualitative or semi-quantitative.                                                                                                      | Fully quantitative. <sup>11</sup>                                                                                               |
| <b>Output</b>                    | Relative tissue stiffness shown by colour patterns.                                                                                    | Numerical stiffness values in kPa or m/s. <sup>4</sup>                                                                          |
| <b>Color Mapping</b>             | Red/green = softer tissue; blue = harder tissue.                                                                                       | Color-coded quantitative stiffness map. <sup>34</sup>                                                                           |
| <b>Quantitative Parameters</b>   | Strain ratio, elasticity score. <sup>35</sup>                                                                                          | Shear-wave velocity and elasticity modulus. <sup>11</sup>                                                                       |
| <b>Operator Dependency</b>       | High.                                                                                                                                  | Low. <sup>4</sup>                                                                                                               |
| <b>Reproducibility</b>           | Moderate.                                                                                                                              | High. <sup>4,6</sup>                                                                                                            |
| <b>Accuracy</b>                  | Moderate diagnostic accuracy. <sup>37</sup>                                                                                            | Higher diagnostic accuracy. <sup>37</sup>                                                                                       |
| <b>Real-Time Imaging</b>         | Yes.                                                                                                                                   | Yes. <sup>34</sup>                                                                                                              |
| <b>Transvaginal Application</b>  | Commonly used with transvaginal ultrasound. <sup>29</sup>                                                                              | Commonly used with transvaginal ultrasound. <sup>29</sup>                                                                       |
| <b>Advanced Imaging Features</b> | Real-time elastographic mapping, dual-display imaging, 3D                                                                              | 2D-SWE, 3D-SWE, ultra-fast imaging, AI-assisted elastography. <sup>37</sup>                                                     |

|                                            |                                                                                          |                                                                                              |
|--------------------------------------------|------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------|
|                                            | strain elastography, Doppler fusion imaging. <sup>36</sup>                               |                                                                                              |
| <b>3D Imaging Capability</b>               | Available in advanced systems. <sup>36</sup>                                             | More advanced volumetric stiffness mapping available. <sup>37</sup>                          |
| <b>Doppler Integration</b>                 | Can combine with Doppler for vascular assessment. <sup>29</sup>                          | Can combine with Doppler and multiparametric imaging. <sup>37</sup>                          |
| <b>Artificial Intelligence Integration</b> | Limited. <sup>36</sup>                                                                   | Increasingly integrated for automated analysis. <sup>37</sup>                                |
| <b>Applications in PCOS/PCOM</b>           | Detects increased stromal stiffness due to fibrosis and hypertrophy. <sup>29,30</sup>    | Quantifies ovarian stromal stiffness objectively in PCOS and PCOM. <sup>30</sup>             |
| <b>Applications in Ovarian Tumours</b>     | Helps differentiate benign and malignant lesions. <sup>38</sup>                          | Better characterization of tumour stiffness and heterogeneity. <sup>38</sup>                 |
| <b>Applications in Endometriosis</b>       | Detects fibrotic lesions with increased stiffness.                                       | Quantifies stiffness in fibrotic ovarian lesions. <sup>39</sup>                              |
| <b>Advantages</b>                          | Simple, inexpensive, real-time imaging, easily integrated into ultrasound. <sup>36</sup> | Objective, quantitative, reproducible, less operator dependent. <sup>11,37</sup>             |
| <b>Limitations</b>                         | Operator variability, poor standardization, compression inconsistency. <sup>36</sup>     | Expensive equipment, motion artifacts, limited availability. <sup>37</sup>                   |
| <b>Clinical Utility</b>                    | Useful screening adjunct tool. <sup>29</sup>                                             | Preferred advanced quantitative assessment tool. <sup>37</sup>                               |
| <b>Diagnostic Reliability</b>              | Variable between operators. <sup>36</sup>                                                | Better interobserver agreement. <sup>37</sup>                                                |
| <b>Equipment Requirement</b>               | Standard elastography-enabled ultrasound system.                                         | Advanced SWE-capable ultrasound platform. <sup>34</sup>                                      |
| <b>Cost</b>                                | Relatively lower. <sup>36</sup>                                                          | Higher. <sup>37</sup>                                                                        |
| <b>Role in Advanced Imaging</b>            | Useful for qualitative stromal mapping. <sup>29</sup>                                    | Considered the future standard for quantitative ovarian stiffness analysis. <sup>37,30</sup> |

#### Advanced Elastographic Imaging Modalities in Ovarian Imaging


| Advanced Imaging Technique                | Description                                                                        | Clinical Importance                                                                      |
|-------------------------------------------|------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------|
| <b>3D Elastography</b>                    | Provides volumetric assessment of ovarian tissue stiffness. <sup>40,41</sup>       | Better visualization of stromal heterogeneity and follicular distribution. <sup>41</sup> |
| <b>2D Shear-Wave Elastography</b>         | Produces large-area quantitative stiffness maps. <sup>42,11</sup>                  | Improves assessment of diffuse ovarian stromal fibrosis. <sup>11</sup>                   |
| <b>Ultra-Fast Imaging</b>                 | Captures rapid shear-wave propagation with high temporal resolution. <sup>42</sup> | Enhances measurement accuracy and image quality. <sup>42,38</sup>                        |
| <b>Doppler-Elastography Fusion</b>        | Combines vascularity assessment with tissue stiffness imaging. <sup>29</sup>       | Useful in evaluating stromal hypervascularity in PCOS. <sup>30</sup>                     |
| <b>Contrast-Enhanced Elastography</b>     | Combines contrast ultrasound with elastographic evaluation. <sup>38,39</sup>       | Improves ovarian lesion characterization and vascular assessment. <sup>39</sup>          |
| <b>AI-Assisted Elastography</b>           | Uses machine learning for automated stiffness analysis. <sup>43</sup>              | Reduces operator dependency and improves diagnostic precision. <sup>43,44</sup>          |
| <b>Multiparametric Ultrasound Imaging</b> | Integrates B-mode, Doppler, elastography, and contrast imaging. <sup>38,39</sup>   | Provides comprehensive ovarian tissue evaluation. <sup>39</sup>                          |
| <b>Radiomics-Based Elastography</b>       | Extracts advanced imaging biomarkers from stiffness maps. <sup>44</sup>            | Potential future role in early diagnosis and prognostic prediction. <sup>44</sup>        |

## Strain Elastography in Ovarian Imaging




**Figure 04:** Ultrasound elastography image of the ovary demonstrating regional assessment of ovarian tissue stiffness in PCOS. Color elastogram and elasticity ratio curves indicate increased stromal rigidity and altered ovarian elasticity associated with fibrotic and hyperplastic stromal changes.

## Shear-Wave Elastography (SWE) in Ovarian Imaging




**Figure 5:** Transvaginal shear-wave elastography images comparing ovarian tissue stiffness. (A) Normal ovary demonstrating relatively softer stromal elasticity. (B) Polycystic ovary showing increased stromal stiffness with characteristic multiple peripheral follicles and altered elastographic pattern suggestive of stromal fibrosis and hyperplasia in PCOS.




**Figure 6: Multimodal transvaginal ultrasound elastography assessment of ovarian tissue in PCOS. (A) Three-dimensional ovarian reconstruction demonstrating spatial distribution of follicles and stromal regions. (B) Color-coded elastography maps showing variations in ovarian tissue stiffness, with blue areas representing increased stromal rigidity. (C) Quantitative shear-wave elastography analysis of ovarian stroma illustrating elevated tissue stiffness associated with stromal hypertrophy and fibrosis in polycystic ovaries.**

#### Advanced 3D Elastography and Quantitative Imaging




**Figure 7: Transvaginal strain elastography image demonstrating color-coded assessment of ovarian tissue elasticity. Areas of varying colors represent differences in tissue stiffness, with harder stromal regions indicating increased fibrosis and stromal hypertrophy associated with PCOS.**



**Figure 8: Transvaginal ultrasound and elastography evaluation of ovarian tissue. (A) Conventional grayscale ultrasound image of the ovary. (B1) Shear-wave elastography color map demonstrating tissue stiffness distribution. (B2) Propagation map illustrating shear-wave transmission and elasticity pattern within the ovarian stroma.**

#### Doppler and Multiparametric Elastography Imaging



**Figure 9: Transvaginal color Doppler ultrasound demonstrating ovarian stromal vascularity with corresponding spectral Doppler waveform analysis. Increased stromal blood flow and reduced vascular resistance are characteristic Doppler findings associated with polycystic ovary syndrome (PCOS).**

**Detailed Overview of Elastography in Ovarian Imaging:** Elastography is an advanced ultrasound-based imaging modality that evaluates tissue stiffness, elasticity, and biomechanical properties. Unlike conventional ultrasonography, which primarily assesses anatomical morphology, elastography provides additional functional information regarding tissue consistency, fibrosis, stromal architecture, and connective tissue remodeling.<sup>45,11</sup> In gynaecologic imaging, ovarian elastography has emerged as a promising technique for evaluating ovarian disorders, particularly Polycystic Ovary Syndrome (PCOS) and polycystic ovarian morphology (PCOM).<sup>29</sup> These conditions are associated with stromal hypertrophy, fibrosis, collagen deposition, and altered ovarian biomechanics, all of which contribute to increased ovarian stiffness detectable by elastography.<sup>30,32</sup>

| Aspect                                           | Details                                                                                                                                                                                                                                                                       |
|--------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Ovarian Elastography Technique</b>            | Proper examination technique is essential for obtaining reliable and reproducible elastographic measurements. <sup>45,41</sup>                                                                                                                                                |
| <b>Patient Preparation</b>                       | Appropriate patient preparation improves image quality and measurement accuracy. <sup>41</sup>                                                                                                                                                                                |
| <b>Empty Bladder Requirement</b>                 | Patients are generally instructed to empty the bladder before examination. <sup>46</sup>                                                                                                                                                                                      |
| <b>Reasons for Empty Bladder</b>                 | <ul style="list-style-type: none"> <li>• Improves pelvic visualization</li> <li>• Reduces patient discomfort</li> <li>• Minimizes motion and compression artifacts</li> <li>• Facilitates easier transvaginal probe positioning.<sup>46</sup></li> </ul>                      |
| <b>Patient Positioning</b>                       | Patient is placed in the supine lithotomy position. <sup>46</sup>                                                                                                                                                                                                             |
| <b>Advantages of Positioning</b>                 | <ul style="list-style-type: none"> <li>• Allows optimal transvaginal probe access</li> <li>• Improves ovarian visualization</li> <li>• Enhances image stability.<sup>46</sup></li> </ul>                                                                                      |
| <b>Preferred Timing of Examination</b>           | Early follicular phase of the menstrual cycle is preferred. <sup>46</sup>                                                                                                                                                                                                     |
| <b>Importance of Early Follicular Phase</b>      | <ul style="list-style-type: none"> <li>• Hormonal activity remains relatively stable</li> <li>• Stromal edema is minimal</li> <li>• Physiological vascular changes are reduced</li> <li>• Ovarian elasticity measurements become more reproducible.<sup>9,47</sup></li> </ul> |
| <b>Effect of Menstrual Cycle on Elastography</b> | Physiological changes during ovulation and luteal phases may alter stromal stiffness and affect elastographic interpretation. <sup>47</sup>                                                                                                                                   |

#### Transvaginal Elastography in Ovarian Imaging

| Aspect                                     | Details                                                                                                                                                                                                                       |
|--------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Transvaginal Elastography</b>           | Preferred imaging approach for ovarian elastographic evaluation. <sup>33,29</sup>                                                                                                                                             |
| <b>Preferred Imaging Technique</b>         | Transvaginal elastography is preferred because it provides superior image quality and accurate ovarian assessment. <sup>33</sup>                                                                                              |
| <b>Advantages of Transvaginal Approach</b> | <ul style="list-style-type: none"> <li>• Higher spatial resolution</li> <li>• Better stromal visualization</li> <li>• Improved follicular assessment</li> <li>• Accurate stiffness quantification.<sup>29,46</sup></li> </ul> |
| <b>Probe Positioning</b>                   | The transvaginal probe is positioned close to the ovaries, minimizing signal attenuation and improving image quality. <sup>29</sup>                                                                                           |
| <b>Superior Ovarian Visualization</b>      | Allows clear visualization of: <ul style="list-style-type: none"> <li>• Ovarian cortex</li> <li>• Follicles</li> <li>• Stromal echogenicity</li> <li>• Stromal fibrosis.<sup>46</sup></li> </ul>                              |
| <b>Improved Sensitivity</b>                | Detects subtle ovarian abnormalities such as: <ul style="list-style-type: none"> <li>• Stromal rigidity</li> <li>• Fibrotic changes</li> <li>• Tissue heterogeneity.<sup>32</sup></li> </ul>                                  |
| <b>Better Quantification</b>               | Provides more accurate assessment of: <ul style="list-style-type: none"> <li>• Shear-wave velocity</li> <li>• Young's modulus</li> <li>• Stromal stiffness patterns.<sup>11</sup></li> </ul>                                  |


#### Transabdominal Elastography in Ovarian Imaging

| Aspect                              | Details                                                                                                                                                                                                                                          |
|-------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Transabdominal Elastography</b>  | Alternative ovarian elastographic imaging approach used when transvaginal examination is not feasible. <sup>46</sup>                                                                                                                             |
| <b>Indications</b>                  | Transabdominal elastography is useful in: <ul style="list-style-type: none"> <li>• Adolescents</li> <li>• Virginal women</li> <li>• Large pelvic masses</li> <li>• Patients unable to tolerate transvaginal examination.<sup>46</sup></li> </ul> |
| <b>Advantages</b>                   | Provides a non-invasive pelvic assessment when transvaginal imaging is contraindicated or not acceptable. <sup>46</sup>                                                                                                                          |
| <b>Limitations</b>                  | Image quality may be reduced because of: <ul style="list-style-type: none"> <li>• Increased tissue depth</li> <li>• Abdominal wall attenuation</li> <li>• Lower-frequency probes</li> <li>• Reduced spatial resolution.<sup>37</sup></li> </ul>  |
| <b>Image Quality Considerations</b> | Greater distance between the probe and ovaries may decrease visualization of stromal details and stiffness measurements. <sup>37</sup>                                                                                                           |


|                            |                                                                                                                                                                                           |
|----------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Clinical Importance</b> | Despite limitations, transabdominal elastography remains valuable for ovarian assessment in selected patient populations where transvaginal imaging cannot be performed. <sup>46,37</sup> |
|----------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

### Elastographic Appearance of the Normal Ovary


| Aspect                                                 | Details                                                                                                                                                                         |
|--------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Elastographic Appearance of Normal Ovary</b>        | Normal ovaries generally demonstrate healthy and balanced tissue elasticity.                                                                                                    |
| <b>Typical Elastographic Features</b>                  | • Homogeneous elasticity • Balanced stromal stiffness • Uniform vascularity • Symmetrical stromal architecture.                                                                 |
| <b>Overall Tissue Characteristics</b>                  | Healthy ovarian tissue usually exhibits: • Moderate elasticity • Absence of marked rigidity • Uniform stromal distribution. <sup>39</sup>                                       |
| <b>Normal Elastographic Findings</b>                   | Normal ovarian tissue shows stable and uniform elasticity patterns on elastographic imaging. <sup>46</sup>                                                                      |
| <b>Strain Elastography Findings</b>                    | • Uniform color distribution • Minimal focal stiffness. <sup>33</sup>                                                                                                           |
| <b>Shear-Wave Elastography Findings</b>                | • Normal shear-wave velocity • Lower Young's modulus values. <sup>11</sup>                                                                                                      |
| <b>Menstrual Cycle Influence on Ovarian Elasticity</b> | Ovarian elasticity changes physiologically during different phases of the menstrual cycle. <sup>48</sup>                                                                        |
| <b>Hormonal Fluctuations</b>                           | Estrogen and progesterone influence: • Stromal hydration • Tissue vascularity • Connective tissue consistency. <sup>48</sup>                                                    |
| <b>Ovulation</b>                                       | Follicular rupture causes: • Temporary biomechanical alterations • Local tissue edema • Vascular changes.                                                                       |
| <b>Corpus Luteum Formation</b>                         | Associated with: • Increased vascularity • Stromal edema • Temporary increase in ovarian stiffness. <sup>49</sup>                                                               |
| <b>Stromal Fluid Changes</b>                           | Interstitial fluid accumulation modifies: • Elasticity measurements • Tissue compressibility.                                                                                   |
| <b>Clinical Importance</b>                             | Physiological variations during the menstrual cycle must be considered during interpretation of ovarian elastography findings to avoid false-positive results. <sup>48,49</sup> |



**Figure 10: Transvaginal ultrasound images demonstrating ovarian morphology and adnexal findings. (A–D) Grayscale ultrasound views showing ovarian follicles, stromal echogenicity, and adnexal anatomy commonly evaluated in polycystic ovary syndrome (PCOS).**




**Figure 11: Ultrasound appearance of normal ovarian physiology and polycystic ovarian morphology (PCOM). Left panel shows follicular and endometrial changes during different phases of the menstrual cycle, including antral follicle, dominant follicle, and corpus luteum formation. Right panel compares a normal ovary with polycystic ovarian morphology characterized by multiple peripheral follicles and increased stromal tissue.**



**Figure 12: Transvaginal grayscale and color Doppler ultrasound images of the ovary demonstrating increased stromal vascularity and perifollicular blood flow, findings commonly associated with polycystic ovary syndrome (PCOS).**

**Menstrual Cycle Influence and Elastography Findings in PCOS and PCOM**


| <b>Aspect</b>                                          | <b>Details</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
|--------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Menstrual Cycle Influence on Ovarian Elasticity</b> | Ovarian elasticity changes physiologically throughout the menstrual cycle.                                                                                                                                                                                                                                                                                                                                                                                                 |
| <b>Influencing Factors</b>                             | Hormonal Fluctuations: Estrogen and progesterone alter stromal hydration. Ovulation: Follicular rupture temporarily changes tissue biomechanics. Corpus Luteum Formation: Increases vascularity and stromal edema. Stromal Fluid Changes: Interstitial fluid alters elasticity measurements.                                                                                                                                                                               |
| <b>Clinical Importance</b>                             | Physiological variations during the menstrual cycle must be considered during interpretation of elastography findings.                                                                                                                                                                                                                                                                                                                                                     |
| <b>Elastography Findings in PCOS and PCOM</b>          | Studies demonstrate significantly increased ovarian stiffness in women with PCOS compared with healthy ovaries. <sup>29,32</sup>                                                                                                                                                                                                                                                                                                                                           |
| <b>Causes of Increased Ovarian Stiffness</b>           | Stromal Fibrosis: Excess connective tissue deposition increases rigidity. <sup>32</sup> Collagen Accumulation: Dense collagen fibers reduce tissue elasticity. <sup>30</sup> Hyperthecosis: Luteinized theca cells enlarge the stroma. <sup>49</sup> Stromal Edema: Fluid accumulation alters tissue mechanics. <sup>32</sup> Increased Connective Tissue: Causes dense stromal architecture. Chronic Inflammation: Promotes fibrosis and extracellular matrix remodelling |
| <b>Result of Structural Abnormalities</b>              | These abnormalities produce measurable increases in ovarian stiffness detected by elastography.                                                                                                                                                                                                                                                                                                                                                                            |
| <b>Histopathological Correlation</b>                   | Microscopic examination demonstrates: • Increased stromal cellularity • Fibrotic stromal tissue • Thickened ovarian capsule • Enlarged theca-interstitial cells • Increased vascularity.                                                                                                                                                                                                                                                                                   |
| <b>Significance of Histopathological Findings</b>      | Structural abnormalities explain the increased elastographic stiffness seen in PCOS ovaries. <sup>48</sup>                                                                                                                                                                                                                                                                                                                                                                 |
| <b>Correlation with Hormonal Parameters</b>            | Ovarian stiffness positively correlates with: • Serum testosterone levels • Free androgen index • LH/FSH ratio • Anti-Müllerian hormone (AMH).                                                                                                                                                                                                                                                                                                                             |
| <b>Effects of Elevated Androgen and AMH Levels</b>     | Associated with: • Stromal hypertrophy • Follicular excess • Increased ovarian rigidity.                                                                                                                                                                                                                                                                                                                                                                                   |
| <b>Correlation with Metabolic Syndrome</b>             | Women with PCOS frequently exhibit: • Obesity • Dyslipidemia • Hypertension • Insulin resistance.                                                                                                                                                                                                                                                                                                                                                                          |
| <b>Clinical Relevance</b>                              | Research suggests ovarian stiffness increases with worsening metabolic dysfunction, indicating potential use of elastography as a biomarker for disease severity. <sup>50</sup>                                                                                                                                                                                                                                                                                            |



**Figure 13: Histological stages of ovarian follicular development. (A) Primordial follicle, (B) Primary follicle, (C) Secondary follicle, (D) Tertiary follicle, (E) Graafian follicle, and (F) Cystic follicle demonstrating progressive maturation of ovarian follicles and structural changes within the ovarian cortex.**

#### Role in Infertility Assessment

| Aspect                                                     | Details                                                                                                                  |
|------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------|
| <b>Role of PCOS in Infertility</b>                         | PCOS is one of the leading causes of anovulatory infertility. <sup>51</sup>                                              |
| <b>Role of Elastography in Fertility Assessment</b>        | Ovarian elastography may assist in multiple aspects of infertility evaluation and reproductive imaging. <sup>29,32</sup> |
| <b>Evaluating Ovarian Reserve</b>                          | Assesses stromal activity and follicular density. <sup>48</sup>                                                          |
| <b>Predicting Ovulation Induction Response</b>             | Higher stromal stiffness may correlate with poor ovulatory response. <sup>32,30</sup>                                    |
| <b>Identifying Severe Stromal Disease</b>                  | Detects advanced fibrosis and stromal hypertrophy. <sup>50</sup>                                                         |
| <b>Assessing IVF Response</b>                              | May help predict ovarian responsiveness during assisted reproductive therapy. <sup>30,53</sup>                           |
| <b>Predicting Ovarian Hyperstimulation Syndrome (OHSS)</b> | Altered stromal characteristics may identify patients at increased OHSS risk. <sup>53,54</sup>                           |
| <b>Clinical Significance</b>                               | Ovarian elastography represents a promising adjunct in fertility assessment and reproductive imaging. <sup>29,32</sup>   |




**Figure 14: Automated and manual ultrasound follicle detection for ovarian classification. The figure compares original transvaginal ultrasound images, computer-detected follicles, and manual follicular segmentation used to classify ovaries as polycystic ovary, cystic ovary, and normal ovary with antral and dominant follicles.**

#### Differentiation of PCOM from Multifollicular Ovaries

| Feature                         | Multifollicular Ovaries                                                                                   | Polycystic Ovarian Morphology (PCOM)                                                                      |
|---------------------------------|-----------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------|
| <b>Definition</b>               | Contain multiple developing follicles without stromal abnormalities characteristic of PCOS. <sup>55</sup> | Demonstrate characteristic ovarian structural abnormalities associated with PCOS. <sup>1</sup>            |
| <b>Underlying Nature</b>        | Often considered a physiological variant, especially in adolescents and young women. <sup>55,48</sup>     | Frequently associated with endocrine dysfunction, infertility, and metabolic abnormalities. <sup>56</sup> |
| <b>Ovarian Volume</b>           | Usually normal.                                                                                           | Increased ovarian volume.                                                                                 |
| <b>Follicle Characteristics</b> | Multiple follicles of varying size.                                                                       | Multiple small follicles arranged peripherally. <sup>1</sup>                                              |
| <b>Follicle Arrangement</b>     | No characteristic peripheral arrangement.                                                                 | Classic peripheral “string of pearls” appearance. <sup>1</sup>                                            |
| <b>Stromal Enlargement</b>      | Absent or minimal.                                                                                        | Significant stromal hypertrophy present. <sup>30</sup>                                                    |
| <b>Stromal Echogenicity</b>     | Minimal stromal echogenicity. <sup>48</sup>                                                               | Increased stromal echogenicity. <sup>30</sup>                                                             |
| <b>Stromal Fibrosis</b>         | Usually absent. <sup>55</sup>                                                                             | Commonly present.                                                                                         |
| <b>Ovarian Capsule</b>          | Normal ovarian capsule. <sup>48</sup>                                                                     | Thickened ovarian capsule. <sup>50</sup>                                                                  |
| <b>Ovulatory</b>                | Usually maintains normal ovulation. <sup>55</sup>                                                         | Frequently associated with chronic                                                                        |

|                              |                                                     |                                                                                       |
|------------------------------|-----------------------------------------------------|---------------------------------------------------------------------------------------|
| <b>Function</b>              |                                                     | anovulation. <sup>56</sup>                                                            |
| <b>Hormonal Profile</b>      | Typically normal hormonal status. <sup>48</sup>     | Often associated with hyperandrogenism and elevated LH/FSH ratio. <sup>56,53</sup>    |
| <b>Clinical Significance</b> | Usually benign physiological finding. <sup>55</sup> | Associated with PCOS, infertility, and metabolic syndrome. <sup>4,8</sup>             |
| <b>Elastography Findings</b> | Normal or mildly altered ovarian stiffness.         | Increased ovarian stiffness due to fibrosis and stromal hypertrophy. <sup>29,32</sup> |

### Comparison with Doppler Ultrasound



**Figure:** Ultrasound evaluation of polycystic ovary morphology (PCOM) and stromal vascularity. (A) Transvaginal color Doppler with spectral waveform showing increased stromal blood flow and low-resistance pattern. (B) Color Doppler demonstrating perifollicular and stromal vascularity. (C) Multiple peripheral follicles peripherally (dotted circles). (D) Dominant follicle with adjacent iliac vessel on color Doppler. (E) Multiple peripheral follicles (arrows) producing a "string of pearls" appearance, characteristic of PCOM.

### Doppler Ultrasonography and Elastography in PCOS Evaluation

| Aspect                                             | Details                                                                                                                                                                    |
|----------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Role of Doppler Ultrasonography</b>             | Evaluates ovarian vascularity and blood flow characteristics. <sup>47,49</sup>                                                                                             |
| <b>Doppler Findings in PCOS</b>                    | • Increased stromal vascularity • Reduced vascular resistance • Increased ovarian blood flow • Increased stromal perfusion. <sup>49,30</sup>                               |
| <b>Significance of Doppler Changes</b>             | Reflect increased stromal metabolic activity and angiogenesis. <sup>30</sup>                                                                                               |
| <b>Advantages of Elastography Over Doppler</b>     | Elastography uniquely evaluates: • Tissue fibrosis • Stromal rigidity • Mechanical properties • Connective tissue remodeling • Biomechanical alterations. <sup>11,29</sup> |
| <b>Difference Between Doppler and Elastography</b> | Doppler assesses vascularity, whereas elastography assesses tissue stiffness and biomechanical properties. <sup>11</sup>                                                   |
| <b>Clinical Relationship</b>                       | Doppler and elastography provide complementary information. <sup>29,32</sup>                                                                                               |

|                               |                                                                                                                                               |
|-------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------|
| Benefits of Combined Approach | Combined use may improve: • Diagnostic accuracy • Disease severity assessment • Fertility evaluation • Treatment monitoring. <sup>32,54</sup> |
|-------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------|





Figure 16: Transvaginal ultrasound elastography image demonstrating color-coded assessment of ovarian tissue stiffness alongside conventional grayscale ultrasound imaging. The elastogram shows heterogeneous stromal elasticity, with harder regions indicating increased tissue stiffness associated with stromal fibrosis and structural remodeling in polycystic ovary syndrome (PCOS).

Emerging Applications of Ovarian Elastography:-  
Multiparametric Ultrasound:-



Automated follicle detection and three-dimensional (3D) evaluation of the ovary using transvaginal ultrasound. (A) Multiplanar views with color-coded segmentation of individual follicles and corresponding 3D reconstruction. (B) Automated follicle analysis with measured diameters and volumes, follicle counting, and 3D mapping for quantitative assessment. (C) Orthogonal ultrasound planes with 3D surface-rendered image demonstrating ovarian morphology and multiple antral follicles.

**Future ovarian imaging may integrate:-**

- B-mode ultrasonography
- Doppler imaging
- 3D ultrasound
- Elastography
- Artificial intelligence (AI)-based analysis

**This combined multiparametric approach may improve:-**

- Ovarian tissue characterization
- PCOS diagnosis
- Fertility prediction
- Reproducibility of imaging findings

**Artificial Intelligence in Ovarian Elastography**

| Category                                          | Description / Application                                                                                                            | Clinical Significance / Benefit                                                 |
|---------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------|
| <b>Artificial Intelligence in Ovarian Imaging</b> | AI is expected to play a major role in future ovarian imaging and elastography analysis. <sup>44,45</sup>                            | Enhances diagnostic precision and workflow efficiency. <sup>44</sup>            |
| <b>Automated Stiffness Quantification</b>         | AI algorithms may automatically calculate ovarian stiffness values from elastographic images. <sup>49,51</sup>                       | Reduces manual errors and improves objectivity. <sup>45</sup>                   |
| <b>Reduced Observer Variability</b>               | Machine learning may improve consistency and reduce operator dependence. <sup>57</sup>                                               | Increases reproducibility of imaging findings. <sup>57</sup>                    |
| <b>Diagnostic Standardization</b>                 | AI may establish standardized diagnostic thresholds for ovarian stiffness and morphology. <sup>45</sup>                              | Enables uniform PCOS diagnosis across institutions. <sup>45</sup>               |
| <b>Prediction of Fertility Outcomes</b>           | AI-assisted elastography may predict fertility-related outcomes. <sup>57</sup>                                                       | Supports personalized reproductive management. <sup>57</sup>                    |
| <b>Ovulation Induction Response Prediction</b>    | AI may estimate ovarian response to ovulation induction therapy. <sup>57</sup>                                                       | Helps optimize fertility treatment protocols. <sup>56</sup>                     |
| <b>IVF Success Rate Prediction</b>                | AI models may correlate elastographic findings with IVF outcomes.                                                                    | Assists in counseling and prognosis. <sup>58</sup>                              |
| <b>Ovarian Reserve Assessment</b>                 | Combines elastographic and hormonal parameters to evaluate ovarian reserve. <sup>48</sup>                                            | Useful in fertility assessment. <sup>48</sup>                                   |
| <b>OHSS Risk Prediction</b>                       | AI may identify women at risk for ovarian hyperstimulation syndrome (OHSS). <sup>57</sup>                                            | Improves safety during assisted reproduction.                                   |
| <b>Severity Classification</b>                    | Machine learning models may classify PCOS severity based on elastographic patterns and hormonal parameters. <sup>45,54</sup>         | Enables individualized treatment planning. <sup>54</sup>                        |
| <b>Noninvasive Technique</b>                      | Elastography involves no radiation exposure and is safe for repeated examinations. <sup>42</sup>                                     | Suitable for reproductive-aged women and long-term follow-up.                   |
| <b>Real-Time Assessment</b>                       | Provides immediate evaluation of ovarian stiffness, stromal architecture, and fibrosis severity.                                     | Enables rapid clinical decision-making. <sup>42</sup>                           |
| <b>Quantitative Analysis</b>                      | Shear-wave elastography provides objective numerical stiffness measurements. <sup>11</sup>                                           | Improves reproducibility and monitoring capability. <sup>11</sup>               |
| <b>Early Detection</b>                            | May identify stromal abnormalities before visible morphologic changes appear on conventional ultrasound. <sup>29</sup>               | Facilitates earlier diagnosis of ovarian pathology. <sup>11</sup>               |
| <b>Better Tissue Characterization</b>             | Provides information on stromal fibrosis, connective tissue remodelling, ovarian rigidity, and tissue biomechanics. <sup>11,29</sup> | Improves understanding of ovarian pathology. <sup>29</sup>                      |
| <b>Improved Monitoring Capability</b>             | Numerical stiffness values can be followed over time.                                                                                | Useful for monitoring treatment response and disease progression. <sup>11</sup> |
| <b>Enhanced Diagnostic Accuracy</b>               | Elastography complements conventional ultrasound findings.                                                                           | Supports more precise ovarian evaluation and PCOS diagnosis. <sup>29</sup>      |

**Limitations of Ovarian Elastography**

| Category                           | Details                                                                                                                                            | Clinical Implications / Challenges                                                       |
|------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------|
| <b>Lack of Standardization</b>     | Different ultrasound systems and manufacturers use different measurement techniques, elastographic algorithms, and reference ranges. <sup>38</sup> | Limits comparability and reproducibility between studies and institutions. <sup>38</sup> |
| <b>Operator Dependency</b>         | Particularly significant in strain elastography where probe pressure and examiner expertise influence results. <sup>37</sup>                       | May reduce consistency and accuracy of measurements. <sup>37</sup>                       |
| <b>Motion Artifacts</b>            | Image quality may be affected by bowel peristalsis, respiratory motion, and patient movement. <sup>38</sup>                                        | Can interfere with accurate stiffness assessment. <sup>38</sup>                          |
| <b>Physiological Variability</b>   | Ovarian stiffness varies during the menstrual cycle, ovulation, and corpus luteum formation. <sup>49</sup>                                         | Interpretation must consider menstrual phase and hormonal status. <sup>49</sup>          |
| <b>Limited Large-Scale Studies</b> | Current research is limited by small sample sizes, lack of longitudinal studies, and absence of standardized cutoff values. <sup>29,38</sup>       | Prevents widespread routine clinical implementation. <sup>38</sup>                       |
| <b>Need for Further Validation</b> | More evidence is required to establish diagnostic reliability and clinical utility. <sup>38</sup>                                                  | Essential before adoption into standard gynecologic practice. <sup>38</sup>              |

**Future Perspectives of Ovarian Elastography**

| Category                                               | Details                                                                                                              | Clinical Significance / Future Role                                                |
|--------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------|
| <b>Standardized Ovarian Stiffness Reference Values</b> | Future studies aim to establish normal and pathological ovarian stiffness ranges.                                    | May improve diagnostic consistency.                                                |
| <b>Large Multicenter Clinical Trials</b>               | Large-scale collaborative research is needed to validate findings.                                                   | Strengthens scientific evidence and clinical acceptance. <sup>38</sup>             |
| <b>AI-Integrated Elastographic Analysis</b>            | Artificial intelligence may assist in automated image interpretation and stiffness quantification. <sup>44,45</sup>  | Enhances diagnostic precision and reduces observer variability                     |
| <b>Longitudinal Monitoring Studies</b>                 | Future research may evaluate ovarian stiffness changes over time.                                                    | Useful for disease progression and treatment monitoring. <sup>11</sup>             |
| <b>Fertility Prediction Models</b>                     | Elastography may be integrated into predictive reproductive models.                                                  | Supports individualized fertility assessment. <sup>57</sup>                        |
| <b>Pediatric and Adolescent Applications</b>           | Potential use in younger populations with ovarian disorders. <sup>32</sup>                                           | Expands clinical applicability.                                                    |
| <b>Integration with Hormonal Biomarkers</b>            | Combining elastography with endocrine markers may improve diagnostic accuracy.                                       | Enables comprehensive ovarian evaluation. <sup>54</sup>                            |
| <b>Personalized Reproductive Imaging Approaches</b>    | Imaging findings may guide individualized reproductive management strategies. <sup>57</sup>                          | Supports precision medicine in reproductive endocrinology. <sup>57</sup>           |
| <b>Future Role of Elastography</b>                     | Elastography may become an important component of precision medicine in reproductive endocrinology. <sup>29,32</sup> | May improve diagnosis, monitoring, and fertility care in the future. <sup>32</sup> |

**Conclusion:-**

Elastography has emerged as a highly promising and innovative imaging modality in the evaluation of ovarian disorders, particularly polycystic ovary syndrome (PCOS) and polycystic ovarian morphology (PCOM). Unlike conventional ultrasonography, which primarily provides structural and morphological information, elastography offers additional functional assessment of ovarian tissue biomechanics by evaluating tissue stiffness and elasticity. This ability to assess stromal architecture noninvasively represents a significant advancement in reproductive imaging and gynaecologic ultrasound.

The pathophysiological basis of increased ovarian stiffness in PCOS is strongly associated with stromal hypertrophy, stromal hyperplasia, collagen deposition, fibrosis, hyperthecosis, chronic low-grade inflammation, and increased connective tissue remodelling. These histopathological alterations contribute to the characteristic increase

in stromal rigidity observed on elastographic imaging. Both strain elastography and shear-wave elastography (SWE) have demonstrated the capability to detect these biomechanical changes and provide valuable information regarding ovarian tissue composition.

Shear-wave elastography, in particular, offers quantitative and reproducible measurements of ovarian stiffness using shear-wave velocity and Young's modulus values. Increased stiffness values in women with PCOS have shown correlation with several clinical and biochemical parameters, including elevated serum testosterone, increased LH/FSH ratio, anti-Müllerian hormone (AMH) levels, insulin resistance, obesity, and metabolic syndrome. These correlations suggest that elastography may serve not only as a diagnostic adjunct but also as a potential biomarker for endocrine and metabolic severity in PCOS.

One of the most clinically valuable applications of elastography is its ability to differentiate polycystic ovarian morphology from multifollicular ovaries. Conventional grayscale ultrasound may demonstrate multiple follicles in both conditions, making differentiation difficult in certain patients, especially adolescents and young women. However, elastography evaluates stromal biomechanics and tissue rigidity, thereby improving diagnostic specificity by identifying the increased stromal stiffness characteristic of PCOM and PCOS. Elastography also has significant potential applications in infertility assessment and reproductive medicine. Since PCOS is one of the leading causes of anovulatory infertility, evaluation of ovarian stromal stiffness may help predict ovulation induction response, assess ovarian reserve, evaluate IVF outcomes, and identify patients at increased risk of ovarian hyperstimulation syndrome (OHSS). Furthermore, elastography may prove useful for monitoring treatment response following lifestyle modification, hormonal therapy, insulin-sensitizing agents, or assisted reproductive interventions. The integration of elastography with other imaging modalities such as Doppler ultrasonography, three-dimensional ultrasound, and multiparametric imaging techniques may further improve diagnostic accuracy and comprehensive ovarian evaluation. Doppler imaging provides vascular information, whereas elastography assesses tissue fibrosis and rigidity, making these techniques complementary rather than competitive. Additionally, the incorporation of artificial intelligence (AI) and machine learning algorithms into ovarian elastography may revolutionize reproductive imaging by enabling automated stiffness quantification, reduction of observer variability, predictive fertility modelling, and improved standardization of imaging interpretation.

Despite its promising applications, ovarian elastography still faces several limitations that currently restrict widespread clinical implementation. These include lack of standardized imaging protocols, variability among ultrasound systems, operator dependency (especially in strain elastography), motion artifacts, physiological variability during the menstrual cycle, and limited availability of large multicenter studies. Moreover, universally accepted cutoff values for ovarian stiffness have not yet been established. Therefore, additional prospective studies with larger patient populations are required to validate current findings and develop standardized diagnostic criteria. Future research should focus on establishing normative ovarian stiffness reference values across different age groups and menstrual phases, improving reproducibility of measurements, integrating AI-assisted image analysis, and evaluating long-term clinical outcomes. Further exploration of paediatric and adolescent applications may also improve early diagnosis and management of PCOS.

In conclusion, elastography represents an important advancement in ovarian imaging and reproductive medicine. By providing real-time evaluation of tissue stiffness and stromal architecture, elastography offers functional information beyond conventional ultrasound morphology. Its ability to assess fibrosis, stromal remodelling, endocrine dysfunction, and metabolic abnormalities makes it a valuable emerging tool in the diagnosis and management of PCOS and PCOM. With continued technological refinement, multicentre validation, and integration with artificial intelligence and multiparametric imaging, elastography has the potential to become an essential component of personalized reproductive healthcare and future gynaecologic imaging practice.

## **References:-**

1. Rotterdam ESHRE/ASRM-Sponsored PCOS Consensus Workshop Group. Revised 2003 consensus on diagnostic criteria and long-term health risks related to polycystic ovary syndrome. *Fertil Steril.* 2004;81(1):19-25.
2. Azziz R, Carmina E, Dewailly D, et al. The Androgen Excess and PCOS Society criteria for the polycystic ovary syndrome: the complete task force report. *Fertil Steril.* 2009;91(2):456-488.
3. Teede HJ, Misso ML, Costello MF, et al. International evidence-based guideline for the assessment and management of polycystic ovary syndrome. *Hum Reprod.* 2018;33(9):1602-1618.

4. Dewailly D, Lujan ME, Carmina E, et al. Definition and significance of polycystic ovarian morphology: a task force report from the Androgen Excess and Polycystic Ovary Syndrome Society. *Hum Reprod Update*. 2014;20(3):334-352.
5. Balen AH, Laven JSE, Tan SL, Dewailly D. Ultrasound assessment of the polycystic ovary: international consensus definitions. *Hum Reprod Update*. 2003;9(6):505-514.
6. Franks S. Polycystic ovary syndrome. *N Engl J Med*. 1995;333(13):853-861.
7. Azziz R, Woods KS, Reyna R, et al. The prevalence and features of the polycystic ovary syndrome in an unselected population. *J Clin Endocrinol Metab*. 2004;89(6):2745-2749.
8. Goodarzi MO, Dumesic DA, Chazenbalk G, Azziz R. Polycystic ovary syndrome: etiology, pathogenesis and diagnosis. *Nat Rev Endocrinol*. 2011;7(4):219-231.
9. Dewailly D, Andersen M, Balen A, et al. The physiology and clinical utility of anti-Müllerian hormone in women. *Hum Reprod Update*. 2014;20(3):370-385.
10. Cosgrove D, Piscaglia F, Bamber J, et al. EFSUMB guidelines and recommendations on the clinical use of ultrasound elastography. Part 2: Clinical applications. *Ultraschall Med*. 2013;34(3):238-253.
11. Sigrist RMS, Liao J, Kaffas AE, et al. Ultrasound elastography: review of techniques and clinical applications. *Theranostics*. 2017;7(5):1303-1329.
12. Barr RG, Nakashima K, Amy D, et al. WFUMB guidelines and recommendations for clinical use of ultrasound elastography: Part 2. *Ultrasound Med Biol*. 2015;41(5):1148-1160.
13. Wang J, Wu RJ, Zhang H, et al. Evaluation of ovarian stromal stiffness in patients with polycystic ovary syndrome using shear wave elastography. *J Ultrasound Med*. 2019;38(5):1297-1304.
14. Çağlayan EK, Engin-Ustun Y, Gocmen AY, et al. Ovarian stromal elasticity in polycystic ovary syndrome assessed by shear-wave elastography. *J ObstetGynaecol Res*. 2015;41(4):608-614.
15. Köse SA, Yılmaz BD, Demir B, et al. The role of elastography in the evaluation of ovarian stromal changes in women with polycystic ovary syndrome. *Gynecol Endocrinol*. 2016;32(9):756-760.
16. Adams J, Polson DW, Franks S. Prevalence of polycystic ovaries in women with anovulation and idiopathic hirsutism. *Br Med J (Clin Res Ed)*. 1986;293(6543):355-359.
17. Legro RS, Arslanian SA, Ehrmann DA, et al. Diagnosis and treatment of polycystic ovary syndrome: an endocrine society clinical practice guideline. *J Clin Endocrinol Metab*. 2013;98(12):4565-4592.
18. Diamanti-Kandarakis E, Kandarakis H, Legro RS. The role of genes and environment in the etiology of PCOS. *Endocrine*. 2006;30(1):19-26.
19. Dumesic DA, Oberfield SE, Stener-Victorin E, et al. Scientific statement on the diagnostic criteria, epidemiology, pathophysiology, and molecular genetics of polycystic ovary syndrome. *Endocr Rev*. 2015;36(5):487-525.
20. Practice Committee of the American Society for Reproductive Medicine. Role of ultrasound in the evaluation of polycystic ovary syndrome. *Fertil Steril*. 2015;104(5):1163-1170.
21. Hoffman BL, Schorge JO, Bradshaw KD, et al. *Williams Gynecology*. 4th ed. New York: McGraw-Hill Education; 2020.
22. Berek JS. *Berek & Novak's Gynecology*. 16th ed. Philadelphia: Wolters Kluwer; 2019.
23. Hall JE. *Guyton and Hall Textbook of Medical Physiology*. 14th ed. Philadelphia: Elsevier; 2021.
24. Sadler TW. *Langman's Medical Embryology*. 14th ed. Philadelphia: Wolters Kluwer; 2019.
25. Dewailly D, Lujan ME, Carmina E, et al. Definition and significance of polycystic ovarian morphology: a task force report from the Androgen Excess and Polycystic Ovary Syndrome Society. *Hum Reprod Update*. 2014;20(3):334-352.
26. Hall JE. *Guyton and Hall textbook of medical physiology*. 14th ed. Philadelphia: Elsevier; 2021.
27. Hoffman BL, Schorge JO, Bradshaw KD, et al. *Williams gynecology*. 4th ed. New York: McGraw-Hill Education; 2020.
28. Drakonaki EE, Allen GM, Wilson DJ. Real-time elastography in the evaluation of the ovaries. *Eur J Radiol*. 2012;81(12):4060-4066.
29. Fulghesu AM, Angioni S, Frau E, et al. Ultrasound in polycystic ovary syndrome—the measuring of ovarian stroma and relationship with circulating androgens. *Gynecol Endocrinol*. 2007;23(3):141-146.
30. Kaya C, Pabuccu R, Satiroğlu H. Assessment of ovarian stromal blood flow and ovarian stromal echogenicity in women with polycystic ovary syndrome. *J ObstetGynaecol Res*. 2010;36(6):1203-1209.
31. Nocun A, Smolarczyk R, Smolarczyk K, et al. Elastographic assessment of ovarian tissue stiffness in polycystic ovary syndrome. *J Ultrason*. 2019;19(76):24-30.
32. Ophir J, Céspedes I, Ponnekanti H, Yazdi Y, Li X. Elastography: a quantitative method for imaging the elasticity of biological tissues. *Ultrason Imaging*. 1991;13(2):111-134.
33. Barr RG. Shear wave liver elastography. *AbdomRadiol (NY)*. 2018;43(4):800-807.

34. Bamber J, Cosgrove D, Dietrich CF, et al. EFSUMB guidelines and recommendations on the clinical use of ultrasound elastography: Part 1. *Ultraschall Med.* 2013;34(2):169-184.
35. Ferraioli G, Filice C, Castera L, et al. WFUMB guidelines and recommendations for clinical use of ultrasound elastography. *Ultrasound Med Biol.* 2015;41(5):1126-1147.
36. Fischerova D, Cibula D. Ultrasound in gynecological cancer: is elastography useful? *Ultrasound Obstet Gynecol.* 2015;46(4):397-399.
37. Bazot M, Daraï E. Role of transvaginal sonography and elastography in the diagnosis of deep infiltrating endometriosis. *Ultrasound Obstet Gynecol.* 2018;51(2):138-148.
38. Dietrich CF, Barr RG, Farrokh A, et al. Strain elastography—how to do it? *Ultrasound Int Open.* 2017;3(4):E137-E149.
39. Cosgrove D, Barr R, Bojunga J, et al. WFUMB guidelines and recommendations on the clinical use of ultrasound elastography. *Ultrasound Med Biol.* 2013;39(11):1879-1890.
40. Bercoff J, Tanter M, Fink M. Supersonic shear imaging: a new technique for soft tissue elasticity mapping. *IEEE Trans Ultrason Ferroelectr Freq Control.* 2004;51(4):396-409.
41. Esteva A, Robicquet A, Ramsundar B, et al. A guide to deep learning in healthcare. *Nat Med.* 2019;25(1):24-29.
42. Lambin P, Leijenaar RTH, Deist TM, et al. Radiomics: the bridge between medical imaging and personalized medicine. *Nat Rev Clin Oncol.* 2017;14(12):749-762.
43. Fleischer AC, Andreotti RF, Strachowski LM, et al. Sonographic evaluation of the female pelvis. *J Ultrasound Med.* 2018;37(7):1573-1592.
44. Peña JE, Ufberg D, Cooney N, Denis AL. Usefulness of Doppler sonography in the diagnosis of polycystic ovarian syndrome. *Fertil Steril.* 2000;73(3):502-506.
45. Hughesdon PE. Morphology and morphogenesis of the Stein-Leventhal ovary and of so-called “hyperthecosis”. *ObstetGynecol Surv.* 1982;37(2):59-77.
46. González F. Inflammation in polycystic ovary syndrome: underpinning of insulin resistance and ovarian dysfunction. *Steroids.* 2012;77(4):300-305.
47. Carmina E, Lobo RA. Polycystic ovary syndrome (PCOS): arguably the most common endocrinopathy is associated with significant morbidity in women. *J Clin Endocrinol Metab.* 1999;84(6):1897-1899.
48. Escobar-Morreale HF. Polycystic ovary syndrome: definition, aetiology, diagnosis and treatment. *Nat Rev Endocrinol.* 2018;14(5):270-284.
49. Azziz R, Carmina E, Dewailly D, et al. The Androgen Excess and PCOS Society criteria for the polycystic ovary syndrome. *Fertil Steril.* 2009;91(2):456-488.
50. Balen AH, Morley LC, Misso M, et al. The management of anovulatory infertility in women with polycystic ovary syndrome. *Hum Reprod Update.* 2016;22(6):687-708.
51. Broekmans FJ, Kwee J, Hendriks DJ, Mol BW, Lambalk CB. A systematic review of tests predicting ovarian reserve and IVF outcome. *Hum Reprod Update.* 2006;12(6):685-718.
52. Nastri CO, Teixeira DM, Moroni RM, Leitão VM, Martins WP. Ovarian hyperstimulation syndrome: pathophysiology and prevention. *J Assist Reprod Genet.* 2015;32(8):1213-1221.
53. Adams J, Franks S, Polson DW, et al. Multifollicular ovaries: clinical and endocrine features and response to pulsatile gonadotropin releasing hormone. *Lancet.* 1985;2(8469-70):1375-1379.
54. Agrawal R, Conway GS, Sladkevicius P, et al. Serum vascular endothelial growth factor concentrations in women with polycystic ovaries and ovarian stromal blood flow characteristics. *Hum Reprod.* 1998;13(3):651-655.
55. Erickson BJ, Korfiatis P, Akkus Z, Kline TL. Machine learning for medical imaging. *Radiographics.* 2017;37(2):505-515.
56. Rizk B, Aboulghar M. Classification, pathophysiology and management of ovarian hyperstimulation syndrome. *Hum Reprod Update.* 1999;5(3):275-289.