

**RESEARCH ARTICLE****COMPARATIVE DOSIMETRIC ASSESSMENT OF AMBULATORY AND INPATIENT IODINE-131 THERAPY USING MONTE CARLO SIMULATIONS**

Placide Koffi Allangba^{1,2,3,4}, Kouassi-Bagui Affiba Florance¹, N'guessan Kouamé Hermann^{1,3} and Guy Muller Okra^{1,2}

1. Laboratory of Fundamental and Applied Physics (LFAP), University Nangui Abrogoua, Abidjan, Côte d'Ivoire.
2. Physics Teaching Unit, Laboratory of Environmental Sciences and Technologies, University Jean Lorougnon Guede, Daloa, Côte d'Ivoire.
3. Institute of Nuclear Medicine of Abidjan (IMENA), Biosecurity and Biosafety Pole, Medical Physics and Radiation Protection Unit, Abidjan, Côte d'Ivoire.
4. Department of Medical Physics, University Trieste and International Centre for Theoretical Physics (ICTP), Trieste, Italy.

Manuscript Info**Manuscript History**

Received: 16 June 2026

Final Accepted: 18 July 2026

Published: August 2026

Key words:-

Iodine-131, ambulatory therapy, inpatient therapy, dosimetry, Monte Carlo simulation, radiation protection.

Abstract

This study presents a regulatory and dosimetric comparison of Three iodine 131 treatment scenarios, involving the administration of 200, 3700 and 7400 MBq for scenarios A, B, and C, respectively. Monte Carlo modeling using an ICRP male phantom was employed to assess dose distribution across tissues and organs. International radiation protection guidelines were applied according to the specific scenarios. Dose comparisons across the scenarios reveal a trend proportional to the administered activity. Occupational exposures and doses received by healthcare staff measured at a distance of 1 meter from the patient, were evaluated against ICRP and IAEA discharge criteria, resulting in dose rates of 7.64, 140, and 280 $\mu\text{Sv/h}$ for scenarios A, B, and C, respectively. The results confirm the safety and feasibility of outpatient treatment for activities of around 786 MBq corresponding to 30 $\mu\text{Sv/h}$ at 1 meter, under controlled exposure conditions, whereas the administration of higher activities necessitates hospitalization in isolation due to cumulative exposure considerations.

"© 2026 by the Author(s). Published by IJAR under CC BY 4.0. Unrestricted use allowed with credit to the author."

Introduction:-

Iodine-131 radiotherapy is widely used for hyperthyroidism and differentiated thyroid carcinoma. Historically, patients receiving high activities were hospitalized due to concerns regarding radiation exposure to staff and the public.

Corresponding Author:-Placide Koffi Allangba

Address:-1.Laboratory of Fundamental and Applied Physics (LFAP), University Nangui Abrogoua, Abidjan, Côte d'Ivoire 2.Physics Teaching Unit, Laboratory of Environmental Sciences and Technologies, University Jean Lorougnon Guede, Daloa, Côte d'Ivoire. 3. Institute of Nuclear Medicine of Abidjan (IMENA), Biosecurity and Biosafety Pole, Medical Physics and Radiation Protection Unit, Abidjan, Côte d'Ivoire. 4. Department of Medical Physics, University Trieste and International Centre for Theoretical Physics (ICTP), Trieste, Italy.

However, evolving radiation protection standards now permit outpatient therapy under strict regulatory conditions. The ICRP recommends patient release based on dose rate measurements and public exposure constraints [1,2], while the IAEA establishes quantitative release criteria, including a dose rate threshold of approximately 30 $\mu\text{Sv/h}$ at 1 meter [3]. Harmonized guidelines from EANM and SNMMI also support ambulatory therapy up to 3.7 GBq provided that appropriate radiation safety instructions are delivered [4,5,6].

External exposure to patients around I-131 has been extensively studied. Measurements show rapid dose rate decay with distance, supporting inverse square law considerations and favoring short duration patient interactions [7]. Monte Carlo simulations have been widely used to predict organ specific absorbed doses in individuals exposed to patients treated with iodine-131 [8,9]. Given these regulatory and radiation protection considerations, comparing staff and caregiver exposures across different therapeutic activities is essential for clinical decision making.

Materials and Methods:-

Monte Carlo modeling assumptions:-

The treated patient was modeled as an external photon emitting source corresponding to the iodine-131 decay scheme. A simplified ICRP adult male anthropomorphic geometry was assumed and absorbed and effective doses were calculated for an external receptor located at 1 m from the patient. This configuration reflects standard radiation protection assessment conditions used for staff and caregiver exposure evaluations. Organ absorbed doses (μGy) and effective doses (μSv) were obtained from Visual Monte Carlo dose calculation software based simulation data sets provided by the scenarios.

Realistic scenario:-

Three activity scenarios were analyzed: Scenario A: 200 MBq, Scenario B: 3700 MBq, Scenario C: 7400 MBq. Exposure durations ranged from 5 to 60 minutes at a fixed distance of 1 meter. These durations represent conservative upperbound scenarios intended to assess worst case radiation protection conditions [10, 11, 12]. Monte Carlo approaches have been validated for similar external dose assessments in the nuclear medicine environment [8,13, 14, 15,16]. These simulations based dosimetric values were cross referenced with published staff exposure measurements [7,16].

Results:-

The results of the Monte Carlo simulation calculations are presented in Tables 1, 2, and 3. The histograms in Figures 1, 2, and 3, derived from these tables, respectively illustrate the Biodistribution of absorbed doses by each organ and tissue, as well as the effective whole-body dose.

Dosimetric simulation of scenario A:-

Table 1: Scenario A, a person is exposed at 1 m from a patient who has absorbed 200 MBq of iodine 131 for durations of 5, 15, 20, 30 and 60 min.

Scenario	Sc1A	Sc2A	Sc3A	Sc4A	Sc5A	Sc6A
¹³¹ I activity (MBq)	200	200	200	200	200	200
Exposure duration (min)	5	10	15	20	30	60
Distance source(x :-100cm,y :26.5cm ,z :126.5cm) -worker (m)	1	1	1	1	1	1
Organs and tissues absorbed dose (μGy)						
Red bonemarrow	0.45	0.90	1.35	1.80	2.70	5.41
colon	0.62	1.24	1.86	2.48	3.72	7.45
lung	0.64	1.28	1.91	2.55	3.83	7.65
stomach	0.75	1.51	2.26	3.01	4.52	9.04
Breast	0.83	1.65	2.48	3.31	4.96	9.92
Remainder	0.55	1.09	1.64	2.18	3.28	6.55
Testes	0.48	0.95	1.43	1.91	2.86	5.72
bladder Wall	0.73	1.47	2.20	2.94	4.40	8.81
Oesophagus	0.63	1.27	1.90	2.53	3.80	7.59

Liver	0.64	1.28	1.92	2.56	3.84	7.67
Thyroid	0.97	1.95	2.92	3.89	5.84	11.68
Bone surface	0.45	0.91	1.36	1.82	2.72	5.45
Brain	0.43	0.87	1.30	1.73	2.60	5.19
Salivary gland	0.52	1.07	1.57	2.09	3.14	6.28
Skin	0.51	1.02	1.54	2.05	3.07	6.14
Eye lens	-	-	-	-	-	-
Wholeeye	0.59	1.17	1.76	2.34	3.51	7.03
Effective dose (μSv)	0.64	1.27	1.91	2.55	3.82	7.64




Figure 1: Absorbed dose of the exposed person in scenario A

Dosimetric simulation of scenario B:-

Table 2: Scenario B: a person is exposed at 1 m from a patient who has absorbed 3700 MBq of iodine 131 for durations of 5, 15, 20, 30 and 60 min.

Scenario	Sc1B	Sc2B	Sc3B	Sc4B	Sc5B	Sc6B
^{131}I activity (MBq)	3700	3700	3700	3700	3700	3700
Exposure duration (min)	5	10	15	20	30	60
Distance source(x :-100cm,y :26.5cm ,z :126.5cm) - worker (m)	1	1	1	1	1	1
Organs and tissues absorbed dose (μGy)						
Red bonemarrow	8.21	16.43	24.64	32.86	49.28	100
colon	12.72	25.45	38.17	50.89	76.34	150
lung	11.40	22.80	34.19	45.59	68.39	140
stomach	12.40	24.80	37.19	49.59	74.39	150
Breast	13.23	26.46	39.69	52.92	79.38	160
Remainder	11.20	22.40	33.61	44.81	67.21	130
Testes	10.42	20.83	31.25	41.67	62.50	120
bladder Wall	10.86	21.72	32.59	43.45	65.17	130

Oesophagus	15.66	31.31	46.97	62.62	93.94	190
Liver	11.83	23.66	35.49	47.32	70.98	140
Thyroid	12.51	25.03	37.54	50.05	75.08	150
Bone surface	8.40	16.80	25.19	33.59	50.39	100
Brain	7.41	14.82	22.23	29.64	44.47	90
Salivary gland	11.87	23.74	35.60	47.47	71.21	140
Skin	9.42	18.83	28.25	37.66	56.50	110
Eye lens	-	-	-	-	-	-
Wholeeye	10.15	20.31	30.46	40.62	60.93	110
Effective dose (μ Sv)	11.54	23.08	34.62	46.15	69.23	140




Figure 2: Absorbed dose of the exposed person in scenario B

Dosimetric simulation of scenario C:-

Table 3: Scenario C, a person is exposed at 1 m from a patient who has absorbed 7400 MBq of iodine 131 for durations of 5, 15, 20, 30 and 60 min.

Scenario	Sc1C	Sc2C	Sc3C	Sc4C	Sc5C	Sc6C
¹³¹ I activity (MBq)	7400	7400	7400	7400	7400	7400
Exposure duration (min)	5	10	15	20	30	60
Distance source (x :-100cm,y :26.5cm ,z :126.5cm) -worker (m)	1	1	1	1	1	1
Organs and tissues absorbed dose (μ Gy)						
Red bonemarrow	16.43	32.86	49.28	65.72	100	200
colon	25.46	50.89	76.34	101.78	150	310
lung	22.80	45.59	68.39	91.18	140	270
stomach	24.80	49.59	74.39	99.18	150	300
Breast	26.46	52.92	79.38	105.84	160	320
Remainder	22.40	44.81	67.21	89.62	130	270
Testes	20.83	41.67	62.50	83.34	120	250

bladder Wall	21.72	43.45	65.17	86.9	130	260
Oesophagus	31.31	62.62	93.94	125.24	120	380
Liver	23.66	47.32	70.98	94.64	190	280
Thyroid	25.03	50.05	75.08	100.1	150	300
Bone surface	16.80	33.59	50.39	67.18	100	200
Brain	14.82	39.64	44.47	79.28	90	180
Salivary gland	23.74	47.47	71.21	94.94	140	280
Skin	18.83	37.66	56.50	75.32	110	230
Eye lens	-	-	-	-	-	-
Wholeeye	30.31	40.62	60.93	81.24	120	240
Effective dose (μSv)	23.08	46.16	69.23	92.32	140	280




Figure 3: Absorbed dose of the exposed person in scenario C

Discussion:-

Time-dose linearity:-

Doses increase linearly with exposure duration for all organs, confirming time-dose proportionality. Comparison of Scenarios A, B, and C shows that doses scale almost exactly with administered activity, consistent with empirical measurements around I-131 patients [6,17]. The maximum effective dose at 1 hour for the Scenario A (200 MBq), Scenario B (3700 MBq) and Scenario C (7400 MBq) equal to $7.64 \mu\text{Sv} \approx 0.00764 \text{ mSv}$, $140 \mu\text{Sv} \approx 0.14 \text{ mSv}$ and $280 \mu\text{Sv} \approx 0.28 \text{ mSv}$, respectively. These results align with published external exposure patterns seen in nuclear medicine workers and caregivers [8,12]. Even the highest scenario (C) remains extremely low compared to the annual regulatory limits but beware of the repetition or cumulative duration of exposures.

Scenario A:-

At 200 MBq, absorbed doses remain in the low microgray range (0.4-11 μGy). The maximum effective dose of $7.64 \mu\text{Sv}$ for 1 hour is negligible compared with annual regulatory limits [1,2]. These results correlate with low activity thyroid therapy studies reporting minimal external exposure for staff [18]. Considering the minimum exposure, even 1 hour of close exposure remains at a few μSv . The impact on personnel could be explained by the fact that an individual could be exposed several thousand times before approaching 20 mSv/year. The impact on close contacts would show that the doses remain well below 1 mSv, even with relatively prolonged exposure times. The dose rate for patient release, according to Table 1, is approximately $7.64 \mu\text{Sv/h}$, significantly lower than $30 \mu\text{Sv/h}$ [3]. From a

clinical perspective, the scenario falls within a wide safety zone, which justifies very flexible outpatient use, subject to basic guidelines and an assessment of the home environment. Outpatient management is feasible under standard radiation protection conditions.

Scenario B:-

The 3700 MBq scenario corresponds to standard outpatient ablation practice. This represents an approximately 18 times increase compared to scenario A. The 60 minutes effective dose about 140 μSv or 0.14 mSv is consistent with occupational exposure surveys in nuclear medicine professionals [12], and caregiver measurements after outpatient therapy [16]. This dose is very low, but not negligible if multiplied by numerous interventions or prolonged contacts. Also, this dose level remains significantly below annual occupational limits (20 mSv/year) and below caregiver specific limits (5 mSv per treatment episode) [2,19]. This supports the feasibility of ambulatory administration at 3.7 GBq when distance and time precautions are followed [4]. For relatives of the treated patient, remaining within 1 meter for cumulative durations, doses could reach a few mSv over several days. The dose rate for patient release, according to Table 2, is approximately 140 $\mu\text{Sv/h}$, which is higher than 30 $\mu\text{Sv/h}$ [3]. This would necessitate hospitalization in some cases. However, in some developed countries, radiation protection regulations recommend patient release under strict conditions. Other studies have shown doses to caregivers or accompanying people in these orders of magnitude, always with a safety margin for well-managed, one-off interventions. For strict safety reasons under controlled exposure conditions, the activity for outpatient treatment corresponding to a dose rate of 30 $\mu\text{Sv/h}$ at a distance of 1 meter from the caregiver is around 786 MBq.

Scenario C:-

7400 MBq represents high-activity inpatient therapy. Effective doses scale predictably (about 280 μSv at 60 min), matching Monte Carlo predictions for high-dose I-131 treatments [11,20]. While still below occupational thresholds, cumulative exposure must be carefully managed for personnel entering patient rooms repeatedly [12,14]. Regulatory recommendations generally advise inpatient isolation for activities $\geq 5.5\text{--}7.4$ GBq [21, 22, 23]. The dose is approximately 37 times higher than in scenario A and approximately 2 times higher than in scenario B. For staff, even 1 minute of exposure equates to approximately 0.0047 mSv (assuming linearity), but actual management involves multiple interventions. The dose rate for patient release, according to Table 3, is approximately 280 $\mu\text{Sv/h}$, significantly higher than 30 $\mu\text{Sv/h}$ [3]. This would necessitate mandatory hospitalization of the patient. For the patient's family, hospitalization clearly reduces all exposure, which is the safest option when activity levels are high [24]. In a clinical setting, scenario C may be necessary for certain metastatic diseases or high-dose treatments, but the hospital approach directly impacts radiation protection and exposure minimization recommendations, making the reduction of staff time spent on site or the number of admissions a priority.

Most exposed organs:-

Regardless of scenario A, B, or C, the frontal and anterior organs (thyroid, breast, stomach, esophagus, liver) receive the highest doses, see figure 1,2 and 3. This reflects their relative proximity to the source (the treated patient) and the minimal path taken through the patient's body. The absorption gradient between superficial and deep organs (such as bone marrow, brain, testicles) demonstrates attenuation and anatomical distance. Deeper organs, being more distant or more covered, receive less, but this effect is still linear with duration and activity.

Dosimetric analysis of Monte Carlo simulation:-

Recent studies further support the robustness of Monte Carlo-based external dose assessments in nuclear medicine. Monte Carlo derived organ doses in iodine-131 therapy are in close agreement with analytical and measurement-based approaches, reinforcing confidence in simulation driven decision making [25]. Similarly, voxel-based human phantom models have improved the anatomical realism of dose calculations, particularly for organs at varying depths, thereby refining estimates of caregiver and staff exposure [26, 27]. Additional investigations confirm that external doses to caregivers remain well below regulatory constraints when time-distance principles are respected [28]. Taking together, these findings provide supplementary evidence that ambulatory iodine-131 therapy up to 3.7 GBq is dosimetrically justified under controlled conditions, while higher activities benefit from inpatient management to optimize cumulative exposure control [29]. The International Commission on Radiological Protection (ICRP) 103 recommends 5 mSv as dose limit for caregivers although that for individuals who are not directly caring for the patient should be restricted to 1 mSv/year [30]

Conclusion:-

Comparative dosimetric evaluation confirms the linear relationship between activity level, exposure duration, and external dose. Ambulatory therapy at 200 and 3700 MBq is safe when proper radiation protection measures are implemented. Our study recommends an activity level of around 786 MBq. Higher activity treatments (7400 MBq) remain safer in controlled inpatient environments where staff interaction time and proximity can be optimized. These findings are fully aligned with international recommendations regarding patient release and occupational radiation protection. Field studies have proposed different thresholds or limits depending on the context, sometimes reaching approximately 370 MBq for exit in certain countries, to limit doses to close contacts and the public while taking local realities into account. They confirm that the external dose measured at 1 m is a crucial indicator for deciding on hospitalization or outpatient care, with thresholds based on conventions, studies and local adaptation.

Funding:-

This study received no external funding.

Acknowledgements:-

The authors would like to thank the Medical Physics team of Nuclear Medicine Institute of Abidjan, Nangui Abrogoua University of Côte d'Ivoire, University of Trieste and International Centre for Theoretical Physics (ICTP) of Italy, as well as the anonymous reviewers, for their contributions.

Declaration of Authors' Contributions:-

The study and the drafting of the manuscript were carried out collaboratively. However, as detailed below, Allangba, Hermann, Affiba and Okrawere were involved in dose calculation of the study. Allangba and Affiba collected the data and drafted the first version of the manuscript. Allangba, Hermann, Affiba and Okra carried out the statistical analyses and data interpretation.

Conflicts of Interest:-

The study was conducted without any conflicts of interest.

References:-

- [1] ICRP. Release of patients after therapy with unsealed radionuclides. ICRP Publication 94. Annals of the ICRP. 2004;34(2).
- [2] ICRP Publication 103, The Recommendations of the International Commission on Radiological Protection. 2007.
- [3] IAEA, Radiation Protection in Medical Uses of Ionizing Radiation, SSG-46, 2018.
- [4] Campenni A., Avram, A.M., Verburg, F.A. et al., « The EANM guideline on radioiodine therapy of benign thyroid disease », Eur J Nucl Med Mol Imaging, vol. 50, no 11, p. 3324-3348, sept. 2023, doi: 10.1007/s00259-023-06274-5.
- [5] Anca M Avram, Luca Giovanella, Bennett Greenspan, Susan A Lawson, Markus Luster, Douglas Van Nostrand, Justin G Peacock, Petra Petranović Ovčariček, Edward Silberstein, Mark Tulchinsky, Frederik A Verburg, Alexis Vrachimis. Vrachimis, SNMMI Procedure Standard/EANM Practice Guideline for Nuclear Medicine Evaluation and Therapy of Differentiated Thyroid Cancer: Abbreviated Version. J Nucl Med 2022 Jun;63(6):15N-35N.
- [6] M. Andersson et S. Mattsson, Improved Patient Dosimetry at Radioiodine Therapy by Combining the ICRP Compartment Model and the EANM Pre-Therapeutic Standard Procedure for Benign Thyroid Diseases, Front. Endocrinol., vol. 12, p. 634955, mars 2021, doi: 10.3389/fendo.2021.634955.
- [7] D. Broggio, « The Nuclear Medicine Patient as a Line Source: The Source Length Is Certainly Not the Patient Height, But It Is a Reasonable Approximation », Health Physics, vol. 123, no 3, p. 208-217, sept. 2022, doi: 10.1097/HP.0000000000001587.
- [8] K. Charles Kouman, K. N'guessan Placide Gabin Allangba, Y. K. H. Fagnidi, et O. Guy Müller Banquet, « Monte Carlo Simulation of Organ Absorbed Dose of Worker's Radiation Exposure in Bone Scintigraphy », J. Drug Delivery Ther., vol. 14, no 11, p. 105-110, nov. 2024, doi: 10.22270/jddt.v14i11.6896.
- [9] D. Shahbazi-Gahrouei et S. Ayat, « Determination of Organ Doses in Radioiodine Therapy using Monte Carlo Simulation », World J Nucl Med, vol. 14, no 01, p. 16-18, janv. 2015, doi: 10.4103/1450-1147.150517.
- [10] A. Abdulhasan Kadhim, P. Sheikhzadeh, S. Farzanefar, S. Yavari, M. Mousa Jber, et M. R. Ay, Measurement of Radiation Exposure to Caregivers of Patients with Thyroid Diseases Treated with Iodine-131: A Review », fbt, nov. 2020, doi: 10.18502/fbt.v7i3.4622.

- [11] Kyu Hwan Jeong, Jae Won Jung, Chang Bum Kim, Byeong-Cheol Ahn, Hyun Kuk Lee, Song Jae Yoo, Orville Day, Jai Ki Lee, Estimation of External Radiation Dose to Caregivers of Patients Treated with Radioiodine after Thyroidectomy, *Health Physics*, vol. 106, no 4, p. 466-474, avr. 2014, doi: 10.1097/HP.0b013e3182a415eb.
- [12] S Dewji, M Bellamy, N Hertel, R Leggett, S Sherbini, M Saba, K Eckerman, Estimated dose rates to members of the public from external exposure to patients with ^{131}I thyroid treatment, *Medical Physics*, vol. 42, no 4, p. 1851-1857, avr. 2015, doi: 10.1118/1.4915084.
- [13] J. S. Hendricks, KJ Adam, Thomas Edward Booth, J.F. Briesmeister, Present and future capabilities of MCNP, *Applied Radiation and Isotopes*, vol. 53, no 4-5, p. 857-861, nov. 2000, doi: 10.1016/S0969-8043(00)00231-1.
- [14] Christelle Huet, Jonathan Eakins, Maria Zankl, José María Gómez-Ros, Jan Jansen, Montserrat Moraleda, Lara Struelens, Deepak K. Akar f, Jorge Borbinha, Hrvoje Brkić, Duc Ky Bui, Kevin Capello, Thi My Linh Dang, Laurent Desorgher, Salvatore Di Maria, Lior Epstein, Dario Faj, Karin Fantinova, Paolo Ferrari, Sebastian Gossio, Katarzyna Tyimińska., « Monte Carlo calculation of organ and effective doses due to photon and neutron point sources and typical X-ray examinations: Results of an international intercomparison exercise, *Radiation Measurements*, vol. 150, p. 106695, janv. 2022, doi: 10.1016/j.radmeas.2021.106695.
- [15] S. P. Jollota, A. U. Khan, J. L. Radtke, O. Kwon, B. P. Bednarz, et L. A. DeWerd, Experimental determination and Monte Carlo validation of ^{225}Ac absorbed dose, *Metrologia*, vol. 62, no 6, p. 065004, déc. 2025, doi: 10.1088/1681-7575/ae1b0a.
- [16] Suphalak Khamruang Marshall, Awatif Hayee abdu nromae, Nadia Noomad, Wanita Durawee, Nantakorn Sirieak, Pornchanok Karnkorn and Virunyupa Keawtong. Evaluation of Environmental Radiation Exposure and Algorithms for Determining the Occupational Effective Dose During $^{99\text{m}}\text{Tc}$ -MDP Bone Scintigraphy: A Comprehensive Analysis. *Appl. Sci.* 2024, 14(23), 11211; <https://doi.org/10.3390/app142311211>
- [17] Marinelli Luigi, Bénard François, Brill Andreas B. Radiation protection considerations in radionuclide therapy. *Health Physics*. 2018;115(5):602–610.
- [18] Traino Caterina, Cremonesi Matteo, Ferrari Marco, Chiesa Carlo. Radiation exposure of family members after ^{131}I therapy. *European Journal of Nuclear Medicine*. 2006;33(7):800–806.
- [19] Larsson CM. Overview of ICRP Committee 5: protection of the environment. *Annals of the ICRP*, 17 Mar 2016, 45(1 Suppl):41–53. <https://doi.org/10.1177/0146645316631210> PMID: 26987889
- [20] Martin Andersson, Lennart Johansson, Keith Eckerman and Sören Mattsson. IDAC-Dose 2.1, an internal dosimetry program for diagnostic nuclear medicine based on the ICRP adult reference voxel phantoms, *EJNMMI Research* (2017) 7:88. DOI 10.1186/s13550-017-0339-3
- [21] Barrington Sarah F., O'Doherty Michael J. Radioiodine therapy in differentiated thyroid cancer. *European Journal of Nuclear Medicine and Molecular Imaging*. 2020;47(5):1041–1052.
- [22] Lassmann Michael, Chiesa Carlo, Flux Glen, Bardies Michel. EANM dosimetry committee guidance on internal dosimetry. *European Journal of Nuclear Medicine and Molecular Imaging*. 2018;45(2):188–205.
- [23] IAEA TECDOC-1608, Release of Patients After Nuclear Medicine Therapy, 2010.
- [24] Ali Abdulhasan Kadhim, Peyman Sheikhzadeh, Saeed Farzanefar, Shima Yavari and Mohammad Reza Ay, radiation dose assessment to family member taking care of non-cancerous thyroid patient treated with ^{131}I therapy in nuclear medicine department. *Radiation Protection Dosimetry* (2020), pp. 1–9. doi:10.1093/rpd/ncaa092
- [25] Shahbazi-Gahrouei D., Ayat S. Determination of organ doses in radioiodine therapy using Monte Carlo simulation. *World Journal of Nuclear Medicine*. 2015;14(1):16–18.
- [26] Kramer R., Zankl M., Williams G., Drexler G. The calculation of dose from external photon exposures using reference human phantoms and Monte Carlo methods. *Physics in Medicine and Biology*. 1997;42(8):1519–1537.
- [27] Kramer R., Vieira J. W., Khoury H. J., Lima V. J. M., Fuelle D. All about MAX: a male adult voxel phantom for Monte Carlo calculations in radiation protection dosimetry. *Physics in Medicine and Biology*. 2003;48(10):1239–1262.
- [28] Iwano S., Ayabe K., Ito S., Kato K. Radiation exposure to family members after outpatient radioiodine therapy. *World Journal of Nuclear Medicine*. 2020;19(2):120–126.
- [29] Mattsson S., Johansson L., Leide-Svegborn S. Radiation exposure of caregivers after radionuclide therapy. *Radiation Protection Dosimetry*. 2016;168(2):238–245.
- [30] Valentin, J. The 2007 recommendations of the international commission on radiological protection. (Oxford: Elsevier) (2007).