



ISSN (O): 2320-5407
ISSN (P): 3107-4928

Journal Homepage: www.journalijar.com

INTERNATIONAL JOURNAL OF ADVANCED RESEARCH (IJAR)

Article DOI:10.21474/IJAR01/24184
DOI URL: <http://dx.doi.org/10.21474/IJAR01/24184>



RESEARCH ARTICLE

ANTI-INFLAMMATORY ACTIVITY OF AQUEOUS EXTRACTS FROM THE FRUITS OF TWO TOMATO VARIETIES CULTIVATED IN COTE DIVOIRE

Dembele Syndoux¹, Nda Kesse Philippe², Koffi Ndri Emmanuel^{2,3} and Anin Atchibri Anin Louise⁴

1. Training and Research Unit for Agriculture, Fisheries Resources and Agro-Industry (ARHAI), University of San Pédro (USP), 01 BP 1800 San Pédro 01, Côte d'Ivoire.
2. Laboratory of Industrial Processes, Synthesis, Environment and New Energies (LAPISEN), National Polytechnic Institute Houphouët Boigny (INP-HB), BP 1093 Yamoussoukro, Côte d'Ivoire.
3. Department of Sciences and Technologies, Advanced Teachers' Training College of Abidjan (ENS), 08 BP 10 Abidjan 08, Côte d'Ivoire.
4. Laboratory of Nutrition and Food Safety (LNSA), Research Training-Unit of Sciences and Food Technologies, University Nangui Abrogoua, Abidjan, Ivory Coast.

Manuscript Info

Manuscript History

Received: 12 July 2026

Final Accepted: 14 August 2026

Published: September 2026

Key words:-

inflammation, Lycopersiconesculentum, anti-inflammatory activity, chronic diseases.

Abstract

Inflammatory diseases constitute a major risk factor for the development of chronic diseases. In this context, the search of natural alternatives about treatments is essential. This study aims to evaluate the anti-inflammatory effect of extracts from two tomato varieties (*Lycopersiconesculentum* var. Cobra F1 and var. Locale Côtelette) cultivated in Côte d'Ivoire. Anti-inflammatory activity was studied in rats (*Rattus norvegicus*) using a paw edema induction model, followed by treatment with the extracts. Edema evolution was monitored after administration. Hematological parameters were also analyzed to evaluate extract safety. Results showed a significant reduction in edema as early as three hours after treatment, with a more pronounced effect over time. No significant changes were observed in leukocyte and platelet counts ($p > 0.05$), while a slight increase in red blood cells and hemoglobin levels was recorded. These findings indicate that both tomato varieties exhibit notable anti-inflammatory activity without apparent immunotoxic effects, suggesting their potential in preventing inflammatory diseases.

"© 2026 by the Author(s). Published by IJAR under CC BY 4.0. Unrestricted use allowed with credit to the author."

Introduction:-

Inflammatory diseases currently constitute one of the major public health problems worldwide (Lutu, 2026; Sarter, 2024). They trigger excessive reactions from the immune system, thereby leading to inflammation throughout the body. These reactions occur in response to various assaults on the organism by antigens of physical, chemical, or biological origin (Joulia, 2022). Inflammatory diseases are divided into four (4) categories: rheumatic diseases (rheumatoid arthritis), auto-immune diseases (lupus, type 1 diabetes), chronic inflammatory bowel diseases (Crohn's disease), and type 2 inflammatory diseases (asthma) (Laurent, 2023). The global prevalence of inflammatory

Corresponding Author:-Dembele Syndoux

Address:-Training and Research Unit for Agriculture, Fisheries Resources and Agro-Industry (ARHAI), University of San Pédro (USP), 01 BP 1800 San Pédro 01, Côte d'Ivoire.

diseases varies considerably depending on the type of disease. Indeed, the overall prevalence of autoimmune diseases is estimated at between 5% and 10% of the global population, with high peaks among women (Nihad, 2023). This predominance among women could be due to several factors, notably hormonal factors (Casimir, 2026; ZenkoufiSawsen and Zinedddine, 2024), the extra chromosome X in women (Lacombe, 2025), and the microbiota (Chloé et al., 2024).

Nowadays, the treatment of inflammatory diseases aims to suppress or modulate the immune response using medications such as non-steroidal anti-inflammatory drugs (NSAIDs), corticosteroids, or immunosuppressants. Although effective, these molecules can cause long-term adverse effects. Might a diet rich in fruits and vegetables, particularly tomatoes offer an interesting alternative for preventing or treating this disease? This study was conducted to address this issue. It aims to evaluate the effect of several tomato varieties (*Lycopersiconesculentum*) grown in Côte d'Ivoire on inflammatory diseases. To this end, inflammation was induced in the animals, which were then treated with extracts from two (2) tomato varieties (Lycopersiconesculentum Mill.). The results obtained following treatment were compared with the corresponding controls.

Materials and Methods:-

Plant material

The tomato (*Lycopersiconesculentum* Mill.) is the plant material used in this study. They were collected on April 24, 2025 from Yamoussoukro's plantations. The collected varieties are the hybrid Lycopersiconesculentum var. Cobra F1 (Figure 1A) and the traditional Lycopersiconesculentum var. Locale Côtelette (Figure 1B).




Figure 1: Different varieties of the *Lycopersiconesculentum* Mill tomato fruit

A= Cobra F1, B= Local cotelette

Animal material:-

This experiments were conducted on Wistar rats of species *Rattus norvegicus*, aged 2 to 3 months (figure 2). The average weights ranging of these animals were from 90 to 200 g. These animals were raised in plastic cages and acclimated to the experimental laboratory conditions. They were fed with pellets provided by the Ivorian Society for the Manufacture of Animal Feed (FACI) and watered with tap water without discontinuity.



Figure 2 : Rat Wistar (*Rattus norvegicus*).

Technicalequipment:-

A digital caliper with 0.01 mm precision was used to measure edema in the animals' paws. (**Figure 3**).



Figure 3 : Pied à coulisse (Précision 0,01).

Methods:-**Samplepreparation :**

Sampling involved firm, ripe fruits, i.e. at the red stage. The tomatoes were washed, cut, and dried in an oven at 55°C for 72 hours before being ground in a porcelain mortar. The tomato extracts were obtained by maceration and decoction according to the method described by Bidiéet al.(2011).

Preparation of aqueous extracts by maceration:

In 1 L of distilled water, 100 g of tomato powder was dissolved. This powder was homogenized in water at room temperature ($30 \pm 2^\circ\text{C}$) under stirring for 24 h. Afterwards, the mixture was centrifuged at 1000 rpm for 10 min. The supernatant obtained was then collected and dried in an oven at 55°C for 48 hours.

Preparation of aqueous extracts by decoction:

This consisted of heating under reflux for 10 min a mixture of 100 g of tomato powder in 1 L of distilled water. After cooling, this mixture was centrifuged at 1000 rpm for 10 min. The supernatant obtained was recovered and dried in an oven at 55°C for 48 hours.

Induction of inflammation by carrageenan:

The rats were fasted for 17 hours before the test. Paw edema (inflammation) was induced according to the method described by Sy et al., (2011). Except for the control group (T0), each rat received an injection under the plantar pad of the right hind paw of a 1% carrageenan solution (40 μL). The volume of the edema caused by this phlogogenic agent was measured with the caliper.

Treatment of animals with the extracts:-

After induction, the rats were divided into groups. A total of fifty (50) rats were used for this study, i.e. 10 groups of 5 rats. One group underwent no procedure (neither induction nor treatment). This group was simply fed and given water. This was the control (T0). The other rats in each group were treated intraperitoneally (ip) with the aqueous extracts and an anti-inflammatory drug used therapeutically (Ibuprofen). Table 1 presents the treatment administered to each group.

Table 1 : Treatment administered to each group after induction of inflammation

		Cobra 26 F1			
Aqueousextract		Maceration		Decoction	
Group	Negative control	Group 1	Group 2	Group 3	Group 4
Number of rats	5	5	5	5	5
Extract dose (mg/kg BW)	water	25	50	25	50
Meanweight (g)	172.33	157.00	173.33	149.67	176.33
Concentration (mg/mL)	-	3.925	8.666	3.742	8.816
Volume to administer (mL)	1	1	1	1	1
		Local côtelette			
Group	Positive control	Group 5	Group 6	Group 7	Group 8
Number of rats	5	5	5	5	5
Extract dose (mg/kg BW)	25 (Ibuprofen)	25	50	25	50
Meanweight (g)	193.33	183.67	193.67	180.33	161.33

Concentration (mg/mL)	4.833	4.592	9.683	4.508	8.066
Volume to administer (mL)	1	1	1	1	1

Measurement of edema size and anti-inflammatory activity:-

The equipment used to measure edema size is the caliper (Figure 3). It involves determining the swelling of the paw by using the external jaws to grip the foot (or any other affected body part) between them. The sliding part is moved until the jaws are in firm contact. The measurement is then displayed on the digital screen.

The percentage increase in paw volume was determined using the following formula:

$$\%AUG = \frac{Pawvolume(Vt) - Initialvolume(V0) \times 100}{Initialvolume(V0)}$$

Vt = Paw volume of the treated animal at time t,

V0 = Paw volume of the T0 control

Anti-inflammatory activity was evaluated by calculating the percentage inhibition of edema (% INH),

$$\%INH = \frac{\%AUG(negativecontrol) - \%AUG(treated) \times 100}{\%AUG(negativecontrol)}$$

%AUG = Percentage increase in paw volume.

Statistical analysis:-

The comparison of mean percentages of increase and inhibition was performed using Student's t-test. A significant difference is represented by $p < 0.05$ and $p < 0.01$; the graph was constructed using the Graph Pad Prism 10 software, $n = 5$ represents the number of experiments per group.

Results and Discussion:-

Results:-

Table II presents the percentage increase in edema at 1 hour, 2 hours, and 3 hours after treatment of rats induced with carrageenan. Three hours after treatment with the extracts, a decrease in edema size was observed. This decrease was more pronounced two hours after treatment. Furthermore, animals treated with the macerate of the Locale cotelette variety at a dose of 25 mg/kg BW showed the lowest percentage increase, dropping from 29.22% one hour after treatment to 9.39% three hours after treatment. These results are significantly different from those of the corresponding control groups ($p < 0.05$). In addition, animals treated with the same extract (macerate of Locale cotelette) at a dose of 50 mg/kg BW showed percentage increases significantly different from those of the corresponding controls after two hours of treatment ($p < 0.01$)

Table II : Percentage increase in paw edema following treatment with tomato extracts.

	Dose (mg/kg BW)	Increase in paw edema (%)		
		1h	2h	3h
Control (water)	-	80.42±17.22	101.72±6.95	59.03±16.45
Ibuprofen (reference)	25	40.37±3.30**	34.37±20.66*	31.50±26.51
Cobra F1 Macerate	25	65.64±8.79	45.78±9.96*	32.90±7.29*
	50	72.90±4.32	44.69±9.51*	31.07±17.61
Cobra F1 Decoction	25	79.98±17.65	53.00±10.55*	38.80±8.55
	50	61.49±20.74	56.03±8.38*	28.15±0.09
Locale cotelette Macerate	25	29.22±11.35*	16.13±9.49*	9.39±6.16**
	50	64.67±13.33	42.59±3.24**	36.79±2.93**
Locale cotelette Decoction	25	82.26±3.90	38.62±13.13*	28.41±22.89
	50	76.14±6.43	50.78±7.37**	25.88±10.10*

*Result significantly different from the control ($p < 0.05$), **Result significantly different from the control ($p < 0.01$), $n=5$.

Figure 4 presents the percentage inhibition of edema in animals treated with the extracts compared to the control groups. The tomato extracts show a percentage inhibition close to that of Ibuprofen (reference anti-inflammatory substance), with the exception of the macerate of the Locale c  telette variety at a dose of 25 mg/kg BW. This extract (macerate of Locale c  telette) shows the highest inhibition rate, exceeding 80% three hours after treatment.




Figure 4 : Percentage inhibition of paw edema following treatment with tomato extracts.

Figures 5, 6, 7, and 8 present the hematological parameters, namely white blood cell count (WBC), hemoglobin (HBG), red blood cell count (RBC), and platelet count (PLT) of rats treated with the different extracts after 3 hours of observation. Comparing these results to those of the respective control rats, white blood cell and platelet counts did not vary considerably relative to controls. Furthermore, animals treated with extracts obtained by decoction of the tomato varieties showed hemoglobin and red blood cell levels higher than those of the respective controls at the significance threshold ($p < 0.05$). Also, animals treated with extracts from the Locale c  telette variety showed higher red blood cell and hemoglobin levels compared to those treated with extracts from the Cobra F1 variety




Figure 5 : White blood cell count of rats three (3) hours after treatment with tomato extracts




Figure 6 : Hemoglobin levels of rats three (3) hours after treatment with tomato extracts




Figure 7 : Red blood cell count of rats three (3) hours after treatment with tomato extracts




Figure 8 : Platelet count of rats three (3) hours after treatment with tomato extracts

*Result significantly different from the control ($p < 0.05$), $n=5$, G/L= Giga per Liter, T/L = Tera per Liter, and g/dL = Gram per deciliter.

Discussion:-

Induction of inflammation in the animals (rats) with carrageenan caused edema in their paws. Following treatment of these animals with tomato extracts, a decrease in edema size was observed, ranging from 29% after 1 hour of observation to 9% after 3 hours. This reduction in edema could be due to the presence of several bioactive

compounds in the tomato extracts (Chanforan, 2010; Seddiki and Tiet Hamida, 2021). These bioactive compounds include notably lycopene and polyphenols.

Lycopene, the flagship compound in tomatoes, is recognized as one of the most powerful antioxidant compounds (Laouicha, 2023; Sadoun and SiTahar, 2023). Its anti-inflammatory activity has been demonstrated by several authors (Boukhari et al., 2023; Chehbi and Fadila, 2018; Landri and Lina, 2020). Indeed, lycopene inhibits the NF-KB pathway (Nuclear Factor kappa-light-chain-enhancer of activated B cells) which is the activation pathway for the synthesis of inflammatory molecules (Boucherrou, 2025; Enchery, 2017). NF-KB allows the cell to respond rapidly to insults (viruses, bacteria, oxidative stress) by activating hundreds of specific genes (Le Roux, 2022; Loyer, 2023). Inhibition of this pathway by lycopene therefore contributes to reducing the synthesis and release of pro-inflammatory messengers such as TNF- α , IL-1 β , IL-6, and IL-8. Furthermore, lycopene promotes the decrease of an enzyme involved in the inflammatory process, namely cyclooxygenase-2 (COX-2). Indeed, lycopene increases the phosphorylation of certain proteins such as AMPK- α 1. This phosphorylation results in reduced expression of the COX-2 enzyme. This reduction leads to a decrease in chronic inflammation, which is the gateway to numerous diseases. These mechanisms have recently been confirmed by a review noting that lycopene also inhibits the JAK/STAT3 and MAPK pathways, further reducing the production of TNF- α , IL-1 β , and IL-6 (Carvalho and Martinez, 2025). This anti-inflammatory activity is not limited to isolated lycopene: a whole tomato aqueous extract recently showed, in lipopolysaccharide-stimulated RAW264.7 macrophages, a significant reduction in PGE-2, TNF- α , and IL-1 β (Adhika and Tih, 2026), while a tomato seed extract inhibits the expression of TNF- α and IL-1 β by blocking the NF-KB transcription factor (Kumar et al., 2021). These observations are consistent with the results of a meta-analysis of randomized clinical trials showing that carotenoid supplementation, including lycopene, significantly reduces circulating inflammatory markers such as CRP and IL-6 (Hajizadeh-Sharafabad et al., 2022).

Several phenolic compounds present in foods and plants have strong anti-inflammatory activity (Fioroni, 2023; Moualek et al., 2018). These phenolic compounds play a crucial role in the prevention and modulation of inflammatory diseases through their antioxidant and anti-inflammatory effects. The polyphenol families responsible for these effects are flavonoids, phenolic acids, and tannins. Flavonoids reduce the production of pro-inflammatory molecules such as prostaglandins and the cytokines TNF- α and IL-6, as well as certain enzymes such as cyclooxygenase (COX) and lipoxygenase (LOX). These enzymes inhibit the metabolism of arachidonic acid, a pro-inflammatory molecule. This inhibition by certain flavonoids could contribute to reducing inflammatory processes. This mechanism has recently been supported by work showing that flavonoids simultaneously inhibit cyclooxygenase-2 (COX-2) and 5-lipoxygenase (5-LOX), the two key enzymes of the arachidonic acid pathway, giving them anti-inflammatory potential comparable to that of classic nonsteroidal anti-inflammatory drugs while limiting the gastrointestinal toxicity associated with the latter (Al-Khayriet al., 2022; Chatzipieriset al., 2026).

Furthermore, certain extracts administered to the rats led to an increase in hematological parameters (hemoglobin and red blood cells) compared to their respective controls. Red blood cells, also called erythrocytes, play a very important role in the body as they are responsible for supplying the body with oxygen. These erythrocytes contain a key component ensuring this function, namely hemoglobin (HBG). Due to its protein nature and iron content, hemoglobin plays a crucial role in the cellular respiration process. A low hemoglobin level in the body can be caused by iron deficiency, which can result in dizziness, shortness of breath, fatigue, and paleness (Naciri, 2014; Silué et al., 2023). The increase in red blood cell and hemoglobin levels following administration of the tomato extracts shows that tomato consumption could help prevent iron deficiency and thus promote the prevention of anemia (WHO, 2011) and combat fatigue (Cheboub, 2025; Gouraud, 2020).

These results are consistent with data from a recent review on iron deficiency, which points out that foods rich in vitamin C, such as tomatoes, promote the reduction of ferric iron (Fe³⁺) to ferrous iron (Fe²⁺) in the intestine, thereby improving the absorption of non-heme iron of plant origin (Kolaršet al., 2025). Regular tomato consumption, in addition to its anti-inflammatory activity, could therefore be a useful dietary supplement in preventing iron deficiency and iron-deficiency anemia, which is particularly prevalent among women of childbearing age and children in developing countries.

Conclusion:-

This work showed that the tomato varieties studied possess strong anti-inflammatory activity. Among the different extracts, the macerate of Locale côtelette appears to be the most effective, as it led to a decrease in inflammation of up to 80%. Moreover, these tomato extracts did not significantly affect hematological parameters. Therefore, regular consumption of these tomatoes could be a promising alternative for the prevention of inflammatory diseases.

References:-

1. Adhika O. A. ;Tih F. (2026) Anti-inflammatory and antioxidant effect of tomato extract (*Solanum lycopersicum* L.) in lipopolysaccharide-induced mice macrophage cells. *Asia-Pacific Journal of Science and Technology*. 31(01), APST-31,
2. Al-Khayri J. M. ;Sahana G. R. ; Nagella P. ; Joseph B. V. ; Alessa F. M. ; Al-Mssallem M. Q. (2022) Flavonoids as potential anti-inflammatory molecules: a review. *Molecules*. 27(9), 2901,
3. Bidié A. P. ; N'guessan B. ; Yapo A. F. ; N'guessan J. D. Djaman A. J. (2011) Antioxidant activities of ten medicinal plants from the Ivorian pharmacopoeia. *Sci. & Nat*. 8, 1-11.
4. Boucherrou Khaled (2025) Study of the alternative NF-KB pathway in T lymphocyte biology: functions and mechanisms in cancer and autoimmunity. Doctoral dissertation, Université Claude Bernard Lyon 1. 250 p,
5. Boukhari Kahina ; Moussa Rahma, Ibtihal Saadoudi (2023) Phytochemical study and biological activities of methanolic extracts from the leaves and fruits of *Pistacia lentiscus* L. . Master's thesis in Biological Sciences (Biochemistry), Université Mohamed El Bachir El Ibrahimi (Algeria). 59 p,
6. Carvalho J. F. de ; Martinez A. T. A. (2025) Lycopene supplementation in rheumatic diseases: a comprehensive review. *Exploration of Musculoskeletal Diseases*. 3, 1007112,
7. Casimir G. (2026) What are the consequences for women of a sex-based inflammatory response? *Bulletin de l'Académie Nationale de Médecine*.
8. Chanforan C (2010) Stability of tomato microconstituents (phenolic compounds, carotenoids, vitamins C and E) during processing: model system studies, development of a stoichio-kinetic model, and validation for the tomato sauce preparation unit step. Doctoral dissertation, Université d'Avignon et des pays de Vaucluse (France). 399 Pages,
9. Chatzipieris F. P. ; Petsas E. ; Lambrinidis G. ; Vassiliou S. ; Chasapis C. T. (2026) Recent advances in dual COX/LOX inhibitor design (2020-2024): establishing “the rule of four for inflammation”. *Life*. 16(1), 163,
10. Cheboub M. (2025) Diagnosis and treatment of iron-deficiency anemia and hemolytic anemia. Doctoral dissertation. Université Ibn Khaldoun-Tiaret. 41 pages,
11. Chehbi Soued et Fadila Chadou (2018) Improving the quality of preserved tomato concentrate using spices: black pepper, cumin, and cubeb pepper. Master's thesis in Product Quality and Food Safety. Université KASDI Merbah-Ouargla (Algeria). 79 p,
12. Chloé M. ; Madjid M. ; Gimenez É. ; Stéphanie H. Jean-Philippe L. (2024) Influence of lifestyle-related factors on vaginal microbiota composition. . *Revue Francophone des Laboratoires*. 563, 61-70.
13. Enchery François (2017) Study of the modulation of the canonical NF-KB activation pathway by Nipah virus non-structural proteins. Doctoral dissertation, Université de Lyon. 167 p,
14. Fioroni Nelly (2023) Evaluation of the antioxidant capacity of African leafy vegetables rich in bioactive compounds and micronutrients, and study of their effect on the modulation of intestinal inflammatory status using an in vitro tri-culture cell model. Thesis in Nutrition and Food Science, Université de Montpellier. 256 p,
15. Gouraud É. (2020) Impact of physical exercise on sickle cell skeletal muscle: characterization of muscle fatigability and the effects of aerobic exercise on the pro-antioxidant balance. Doctoral dissertation, Université de Lyon. 259 pages,
16. Hajizadeh-Sharafabad F. ; Zahabi E. S. ; Malekahmadi M. ; Zarrin R. ; Alizadeh M. (2022) Carotenoids supplementation and inflammation: a systematic review and meta-analysis of randomized clinical trials. *Critical Reviews in Food Science and Nutrition*. 62, 8161-8177,
17. Joulia E. (2022) Roles of the Eomes transcription factor in the function, metabolism, and survival of CD4 T lymphocytes in a murine model of multiple sclerosis. Doctoral dissertation, Université Paul Sabatier-Toulouse III.
18. Kolarš B. ; Mijatović Jovin V. ; Živanović N. ; Minaković I. ; Gvozdenović N. ; Dickov Kokeza I. ; Lesjak M. (2025) Iron deficiency and iron deficiency anemia: a comprehensive overview of established and emerging concepts. *Pharmaceuticals*. 18(8), 1104,
19. Kumar M. ; Tomar M. ; Bhuyan D. J. ; et al. (2021) Tomato (*Solanum lycopersicum* L.) seed: a review on bioactives and biomedical activities. *Biomedicine & Pharmacotherapy*. 142, 112018,
20. Lacombe V. (2025) Immunology (the scientific news you wouldn't have dared read elsewhere): X chromosome, Xist, and female autoimmunity. . *La Revue de Médecine Interne*. 46(6), 348-351.
21. Landri Lycia, Lina Simoud Yasmine (2020) Nutritional, antioxidant, and therapeutic properties of watermelon (*Citrullus lanatus*). Master's thesis in Food Science. Université Mouloud Mammeri de Tizi-Ouzou (Algeria). 76 p,
22. Laouicha S (2023) Antioxidant and anti-inflammatory effects of *Arbutus unedo* L. extracts. . Doctoral dissertation in Biological Sciences (Biochemistry). Université Ferhat Abbas Sétif 1. 134 p,

23. Laurent T. (2023) Development of new approaches for analyzing cellular interactions in immune diseases. Doctoral dissertation, Nantes Université.
24. Le Roux Mélina (2022) Role of IL-20 cytokines during viral exacerbations of COPD and bacterial superinfections. Doctoral dissertation, Université de Lille. 239 p,
25. Loyer Chloé (2023) Involvement of neutrophil granulocytes in the pathophysiology of severe forms of COVID-19. Doctoral dissertation, Sorbonne Université. 221 p,
26. Lutu A., Lavergne, L., Peltzer, J., Banzet, S., Luce, E., & Duclos-Vallée, J. C. (2026) Mesenchymal stromal cells and treatment of liver diseases: current status and perspectives. *Hépatogastro- & Oncologie Digestive*. 33(3), 337-346.
27. Moualek I ; Hadim Katia Aissou Chabha Ben (2018) Effect of heat treatment on the antioxidant, anti-inflammatory, and antimicrobial activity of *Arbutus unedo* L. aqueous extract. Master's degree in Applied Microbiology, Université Mouloud Mammeri de Tizi-Ouzou. 98 p,
28. Naciri B. (2014) Anemia in inflammatory bowel diseases: prevalence and management. Medical thesis, Faculté de médecine et de pharmacie, Rabat (Morocco).
29. Nihad Bouzeraa., & Aida, Mecheri. (2023) Epidemiological study of inflammatory diseases among pregnant women in the Mila region. Doctoral dissertation, University center of Abdalhafid Boussouf-MILA.
30. Sadoun F Sitahar M (2023) Comparative study of the antioxidant activity of citrus fruits Master's thesis in Biological Sciences, Université Mouloud Mammeri de Tizi-Ouzou (Algeria). 69p,
31. Sarter H. (2024) Study of chronic inflammatory bowel diseases using a general population registry: contributions to epidemiology and to the statistical analysis of clinical and omics data. Doctoral dissertation, Université de Lille.
32. Seddiki Dounia Tiet Hamida A. E (2021) Valorization of tomato by-products. Master's degree in Food Science. Université Mohamed El Bachir El Ibrahimi (Algeria). 52 p,
33. Silué D ; Atsé Achi a, P. Diabaté S. (2023) Explanatory and interpretive model of severe anemia (Gbélile) among the Senufo of Korhogo (Côte d'Ivoire). *Revue Africaine des Sciences Sociales et de la Santé Publique*. 5(2), 219-233.
34. Sy G. Y. ; Barbosa F. S. ; Wele A. ; Gueye P. M. ; Gueye C. D. ; Cissé A. B. Faye (2011) Evaluation of the Anti-Inflammatory Activity of *Annona Reticulata* (Annonaceae) Leaves on Acute Carrageenan-Induced Rat Paw Edema. *Pharmacopée et médecine traditionnelle africaine*. 15, 23 - 25.
35. Zenkoufi Sawsen B. M. Zineddine S. (2024) Sex hormones and autoimmune diseases.