Journal Homepage: -www.journalijar.com

INTERNATIONAL JOURNAL OF ADVANCED RESEARCH (IJAR)

Article DOI:10.21474/IJAR01/24476
DOI URL: http://dx.doi.org/10.21474/IJAR01/24476

ISSN (O): 2320-5407
ISSN (P): 3107-4928

*RESEARCH ARTICLE*

# SELF-TOLERANCE, AUTOIMMUNITY AND INDIVIDUALIZED THERAPEUTICS: EXPLORING THE ROLE OF HOMOEOPATHIC INDIVIDUALIZATION IN IMMUNE REGULATION

**Ankita Acharya[1] and Vikrant Tripathi[2]**

1. Assistant Professor, Department of Pathology & Microbiology, University College of Homoeopathy, Dr. S. R. Rajasthan Ayurved University, Jodhpur, Rajasthan – 342037.
2. Associate Professor, Department of Human Anatomy, University College of Homoeopathy, Dr. S. R. Rajasthan Ayurved University, Jodhpur, Rajasthan – 342037.

## *Manuscript Info*

***Manuscript History***
Received: 16 August 2026
Final Accepted: 18 September 2026
Published: October 2026

***Key words:-***
Autoimmunity; self-tolerance; immune homeostasis; individualized homoeopathy; immunomodulation; personalized therapeutics.

## *Abstract*

**Background:** Autoimmunity develops when the mechanisms that maintain immunological self-tolerance become dysregulated, allowing immune responses to be directed against endogenous tissues. Autoimmune diseases are heterogeneous: patients with the same diagnosis differ in symptom pattern, disease activity, progression and treatment response, which raises the question of whether individualized therapeutic approaches have a role in disorders of immune dysregulation. Homoeopathy is based on the principle of individualization, in which treatment is selected according to the totality of the patient's characteristic symptoms rather than the diagnostic label alone.

**Methods:** Narrative review and conceptual analysis of the immunological literature on central and peripheral tolerance, regulatory T cells and the breakdown of self-tolerance, together with the published evidence on individualized homoeopathy, including a systematic review and meta-analysis of individualized homoeopathic trials, a systematic review of homoeopathy in rheumatological diseases and a randomized trial in rheumatoid arthritis.

**Results:** Evidence from randomized trials and systematic reviews suggests that individualized homoeopathy has been associated with clinical effects in some settings, although the quality and consistency of the evidence remain insufficient for definitive conclusions. In rheumatological diseases, available studies provide preliminary indications of symptomatic improvement but are limited in number and methodological quality.

Direct evidence that individualized homoeopathic medicines restore central or peripheral immune tolerance is currently lacking. Restoration of self-tolerance is therefore proposed as a testable biological hypothesis, not an established mechanism, and a research framework with objective immunological end points is outlined.

*"© 2026 by the Author(s). Published by IJAR under CC BY 4.0. Unrestricted use allowed with credit to the author."*

**Corresponding Author:-** Ankita Acharya
**Address:-** Assistant Professor, Department of Pathology & Microbiology, University College of Homoeopathy, Dr. S. R. Rajasthan Ayurved University, Jodhpur, Rajasthan – 342037.

**Conclusion:** Individualized homoeopathic therapeutics may be investigated as a potential approach to modifying immune dysregulation in autoimmune disease. Future studies combining individualized prescribing with assessment of regulatory T cells, autoantibodies, cytokines, disease activity and quality of life may determine whether clinically observed changes are accompanied by measurable immunological changes.

## Introduction:-

The immune system is designed to protect the host against infectious and potentially harmful agents while avoiding destructive responses against normal tissues. This capacity is maintained through a complex network of mechanisms collectively referred to as immunological self-tolerance.[1]Self-tolerance is established centrally during lymphocyte development and maintained peripherally through anergy, apoptosis, regulatory T cells and inhibitory immune pathways. Failure of these mechanisms can permit autoreactive lymphocytes to survive and become activated, ultimately producing autoimmune tissue injury.[1,2]

Autoimmune diseases are heterogeneous. Individuals with the same conventional diagnosis may differ considerably in their presenting symptoms, severity, organ involvement, laboratory findings, disease progression and response to treatment. This heterogeneity provides a rationale for investigating individualized approaches to treatment.Homoeopathy is characterized by the principle of individualization, in which the prescription is selected according to the totality of characteristic symptoms of the individual patient. The clinical question, therefore, is whether an individualized therapeutic approach could have measurable effects in disorders characterized by dysregulated immune responses.

An important hypothesis arising from this concept is that, if autoimmune disease represents a state of disturbed immune regulation, an effective individualized intervention might potentially be associated with restoration of immune homeostasis and, ultimately, improved self-tolerance. This hypothesis requires experimental verification.This article first summarizes the immunological basis of self-tolerance and its breakdown (Section 2), then considers why individualization may be relevant in autoimmune disease and reviews the available clinical evidence (Sections 3 and 4). It then sets out the self-tolerance hypothesis and a framework for testing it (Sections 5 and 6), and discusses the limitations of current evidence and directions for future research (Section 7).

### Immunological Basis of Self-Tolerance:-

### Self-tolerance and immune homeostasis:-

Self-tolerance can be broadly divided into central and peripheral tolerance (Table I). Central tolerance operates during lymphocyte development, through negative selection of T cells in the thymus[3,4] and selection of B cells in the bone marrow,[5] and involves receptor editing and deletion of strongly autoreactive lymphocytes. Peripheral tolerance operates on mature lymphocytes and comprises anergy, apoptosis, regulatory T-cell suppression, inhibitory receptor signalling and controlled antigen presentation.[1,6] A simplified model of how these mechanisms lead to immune homeostasis, and of what follows when they become inadequate, is shown in Figure I.

**Table I. Mechanisms of central and peripheral self-tolerance.**

| Type of tolerance | Mechanism | Principal effect |
|---|---|---|
| Central | T-cell negative selection (thymus) | Removal of strongly autoreactive T cells |
| Central | B-cell selection (bone marrow) | Elimination or control of autoreactive B cells |
| Central | Receptor editing | Modification of autoreactive receptor specificity |
| Central | Deletion of strongly autoreactive lymphocytes | Removal of dangerous autoreactive clones |
| Peripheral | Anergy | Functional inactivation of autoreactive cells |
| Peripheral | Apoptosis | Deletion of autoreactive cells |
| Peripheral | Regulatory T-cell suppression | Active suppression of autoreactive responses |
| Peripheral | Inhibitory receptor signalling | Dampening of lymphocyte activation |
| Peripheral | Controlled antigen presentation | Prevention of inappropriate activation |

**Sources: refs 1, 3–6. The "principal effect" column is a summary descriptor added for this table.**





**Figure I. Simplified model of self-tolerance. (A) Normal development of central and peripheral tolerance leading to immune homeostasis. (B) Sequence that follows when these mechanisms become inadequate. Treg, regulatory T cell.**

**Breakdown of self-tolerance:-**

The development of autoimmunity is multifactorial. Genetic susceptibility and environmental influences lead to altered antigen presentation and failure of immune regulation, with activation of autoreactive lymphocytes, expansion of B and T cells, production of autoantibodies and inflammatory cytokines, and eventually tissue injury and clinical disease (Figure II).[2,7,8] Important mechanisms include molecular mimicry, epitope spreading, abnormal apoptosis, altered antigen presentation and defective regulatory mechanisms.





**Figure II. Multifactorial pathway from genetic and environmental factors to clinical autoimmune disease.**

**Regulatory T cells and restoration of immune balance:-**

Regulatory T cells (Tregs), particularly FOXP3-expressing T cells, are critical components of peripheral tolerance. Their functions include suppression of autoreactive immune responses and maintenance of immune homeostasis.[6] Conceptually, adequate Treg activity suppresses autoreactive lymphocytes and maintains self-tolerance, whereas impaired regulatory mechanisms permit autoreactive immune activation, chronic inflammation and autoimmune disease (Figure III).[9]





**Figure III. Conceptual role of regulatory T cells (Tregs). (A) Adequate Treg activity maintains self-tolerance. (B) Impaired regulatory mechanisms lead to chronic inflammation and autoimmune disease.**

Restoration of immune regulatory mechanisms therefore represents an important therapeutic objective in autoimmune disease. This provides a potential biological framework for investigating any intervention proposed to influence immune homeostasis.

**Individualization in Autoimmune Disease:-**

**Why individualization may be relevant:-**

Autoimmune disease is not a uniform biological state. Patients with the same diagnosis may show different symptom patterns, organ involvement, inflammatory profiles, autoantibody patterns, disease activity and treatment responses. Thus, the same diagnosis does not imply an identical biological phenotype. This is consistent with the contemporary movement toward individualized and precision medicine.Homoeopathy approaches the individual from a different theoretical framework, emphasizing the totality of characteristic symptoms. Conventional diagnosis identifies the disease category, whereas homoeopathic individualization attempts to characterize the individual patient's symptom pattern (Table II). The potential research question is whether this individualized therapeutic selection produces reproducible differences in clinical and biological outcomes.

**Table II. Conventional diagnosis compared with homoeopathic individualization.**

| Aspect | Conventional diagnosis | Homoeopathic individualization |
|---|---|---|
| What it identifies | The disease category | The individual patient's characteristic symptom pattern |
| Basis of treatment selection | Disease-based treatment | Individualized treatment |
| Two patients with the same autoimmune diagnosis | Share the same diagnostic label | May receive different prescriptions |

**Homoeopathic individualization:-**
Individualization is one of the defining principles of classical homoeopathic practice.

**The prescription is traditionally selected after considering the totality of symptoms, including:-**
- characteristic symptoms;
- modalities;
- concomitant symptoms;
- temporal patterns;
- individual reactions;
- constitutional characteristics; and
- other features considered relevant during case-taking.

Consequently, two patients with the same autoimmune diagnosis may receive different homoeopathic prescriptions. This creates an important methodological distinction between disease-based treatment and individualized treatment, and evidence specifically examining individualized homoeopathy is therefore particularly relevant to this discussion.A systematic review and meta-analysis published in 2014 evaluated 32 randomized controlled trials involving individualized homoeopathic treatment across 24 medical conditions. The meta-analysis of 22 trials reported an overall effect favouring individualized homoeopathy, although the authors emphasized that the overall evidence quality was low or unclear and recommended further high-quality trials.[10] This finding provides evidence for a research signal associated with individualized treatment, but it does not establish a mechanism involving self-tolerance.

**Evidence in Rheumatological and Autoimmune-Related Conditions:-**
Rheumatological disorders provide an important clinical area for investigating individualized approaches because several are characterized by immune dysregulation and chronic inflammation. The principal sources of evidence discussed in this article are summarized in Table III.A 2024 systematic review of homoeopathy in rheumatological diseases identified 15 studies involving 811 patients. The conditions included rheumatoid arthritis, osteoarthritis, ankylosing spondylitis, hyperuricemia and tendinopathy. The authors reported that most included studies showed improvement, while also emphasizing the limited number of studies and the need for larger future investigations.[11]

An earlier randomized, double-blind, placebo-controlled trial investigated homoeopathy in 112 patients with rheumatoid arthritis. Among those completing the study, reductions were reported in pain, articular index and ESR during the treatment period. However, the study had substantial attrition, with only 58 participants completing the trial, and the trial authors did not find clear evidence that these improvements were attributable to active homoeopathic treatment as opposed to placebo; the findings therefore cannot be interpreted as proof that homoeopathy restored immunological tolerance.[12]These findings are relevant because they support investigation of individualized approaches in immune-mediated disorders, but they do not establish the proposed biological mechanism.

**Table III. Summary of the clinical evidence discussed.**

| Source | Design | Population | Main findings and limitations |
|---|---|---|---|
| Ref. 10 (2014) | Systematic review and meta-analysis of randomized placebo-controlled trials of individualized homoeopathy | 32 trials, 24 conditions; meta-analysis of 22 trials | Overall effect favouring individualized homoeopathy; evidence quality low or unclear; further high-quality trials recommended |
| Ref. 11 (2024) | Systematic review of homoeopathy in rheumatological diseases | 15 studies, 811 patients (rheumatoid arthritis, osteoarthritis, ankylosing spondylitis, hyperuricemia, tendinopathy) | Most studies showed improvement; limited number of studies; larger studies needed |

| Source | Design | Population | Main findings and limitations |
|---|---|---|---|
| Ref. 12 (2001) | Randomized, double-blind, placebo-controlled trial | 112 patients with rheumatoid arthritis; 58 completed | Reductions in pain, articular index and ESR among completers; substantial attrition; improvements not clearly attributable to active treatment over placebo |

**The Self-Tolerance Hypothesis:-**

**From clinical individualization to the self-tolerance hypothesis:-**

The central hypothesis proposed in this article is represented in Figure IV. Individual patient characteristics inform homoeopathic case individualization and the selection of an individualized intervention; a potential clinical response and modification of disease activity may follow, and it is hypothesized that this is accompanied by restoration of immune homeostasis, improved regulation of autoreactive immune responses and improved self-tolerance. This sequence should currently be regarded as a research hypothesis. It should not be presented as a proven pharmacological or immunological mechanism.





**Figure IV. Proposed sequence from homoeopathic case individualization to improved self-tolerance. Solid boxes indicate observable clinical steps; dashed boxes indicate hypothesized immunological steps that require experimental verification.**

**Proposed immunological model:-**

If individualized homoeopathic treatment produces genuine disease-modifying effects in autoimmune disease, one possible area for investigation would be whether such effects are accompanied by measurable changes in immune regulation. A hypothetical model is shown in Figure V: autoimmune dysregulation is characterized by increased autoreactive immune activity, reduced regulatory control and increased inflammatory signalling; an individualized intervention is hypothesized to restore immune balance, with increased regulatory mechanisms and reduced pathological inflammatory activity, leading to reduced tissue injury and improved clinical disease activity. This model requires experimental testing.





**Figure V. Hypothetical immunological model of the effect of an individualized therapeutic intervention in autoimmune disease. The dashed box indicates a hypothesized step. ↑, increased; ↓, decreased.**

**Clinical effect versus mechanism:-**

A fundamental distinction must be maintained between clinical effect and mechanism. Three levels of inference can be separated (Table IV). At present, evidence supporting the first level is more developed than evidence establishing the third level specifically for self-tolerance. Therefore, "homoeopathy restores self-tolerance" should be presented as a hypothesis requiring investigation, not as an established conclusion.

**Table IV. Levels of inference distinguishing clinical effect from mechanism.**

| Level | Example | Interpretation |
|---|---|---|
| 1. Clinical observation | A patient receiving individualized homoeopathic treatment reports improvement | Can be studied as a clinical outcome |
| 2. Biological association | The patient also demonstrates changes in inflammatory or immune biomarkers | Suggests a possible biological association |
| 3. Mechanistic evidence | Controlled experimental studies demonstrate that the intervention directly modifies a defined immune pathway | Provides evidence for a mechanism |

**Testing the Hypothesis:-**

**Potential biological markers:-**

To investigate the hypothesis objectively, future clinical studies could measure the cellular, humoral, inflammatory, disease-specific and clinical markers listed in Table V. This combination would allow researchers to determine whether clinical improvement is accompanied by measurable immunological changes.

**Table V. Potential biological and clinical markers for testing the self-tolerance hypothesis.**

| Category | Markers |
|---|---|
| Cellular markers | Regulatory T-cell frequency; CD4/CD8 populations; B-cell subsets; activated lymphocyte populations |

| Category | Markers |
|---|---|
| Humoral markers | Disease-specific autoantibodies; total immunoglobulin levels where relevant; complement components |
| Inflammatory markers | C-reactive protein; ESR; cytokine profiles |
| Disease-specific markers (depending on the disease) | Rheumatoid factor; anti-CCP antibodies; ANA and disease-specific antibodies; thyroid antibodies; disease-specific inflammatory biomarkers |
| Clinical outcomes | Validated disease activity scores; symptom severity; functional status; quality of life; frequency of disease exacerbations |

**A proposed research framework:-**

The design shown in Figure VI could be used to investigate the central hypothesis. Patients with confirmed autoimmune disease undergo baseline assessment of clinical phenotype, disease activity, autoantibody profile, inflammatory biomarkers, Treg/B-cell profile and quality of life; receive an individualized homoeopathic prescription; are followed up in a standardized manner with clinical and laboratory assessment; and are compared with an appropriate control, followed by statistical analysis. This approach is substantially stronger scientifically than simply reporting symptomatic improvement.





**Figure VI. Proposed research framework to determine whether clinical improvement after individualized homoeopathic prescribing is accompanied by immunological change.**

**A testable hypothesis:-**

The hypothesis can be formulated as follows: in patients with autoimmune disease, individualized homoeopathic treatment, when compared with an appropriate control, is associated with improvement in validated clinical disease activity and measurable changes in biological markers of immune regulation. Possible primary and secondary outcomes are listed in Table VI. This transforms a theoretical concept into a testable scientific question.

**Table VI. Possible primary and secondary outcomes for a test of the hypothesis.**

| Outcome type | Outcomes |
|---|---|
| Primary | Disease activity score; validated clinical response; quality of life |
| Secondary | Autoantibody titres; Treg frequency/function; inflammatory cytokines; complement levels; CRP/ESR; treatment requirements |

## Discussion:-

**Limitations of current evidence:-**

The current literature has several limitations. First, individualized homoeopathic trials cover heterogeneous medical conditions, making direct extrapolation to autoimmune disease difficult.[10] Second, the systematic review of individualized homoeopathy found considerable variation in study quality, with many trials having unclear or high risk of bias.[10] Third, the 2024 rheumatological systematic review included relatively few studies and emphasized the need for larger studies and replication.[11] Fourth, clinical improvement does not demonstrate restoration of self-tolerance. Finally, no sufficiently robust evidence currently establishes that individualized homoeopathic medicines directly increase Treg function, eliminate autoreactive lymphocytes or reverse the immunological basis of autoimmune disease. These limitations identify important opportunities for future research rather than invalidating the research question itself.

**Future research directions:-**

The next generation of studies should move beyond purely symptomatic outcomes. A particularly useful research model would combine homoeopathic individualization, validated autoimmune disease criteria, objective immunological biomarkers and long-term clinical follow-up.

**This could answer three sequential questions:-**

1. Does individualized homoeopathic treatment produce reproducible clinical improvement?
2. Is clinical improvement accompanied by measurable changes in immune regulation?
3. Are those immunological changes consistent with improved regulation of autoreactive immune responses?

Only if these questions are answered through appropriately controlled research could the proposed relationship between individualization and restoration of self-tolerance be supported.

## Conclusion:-

Autoimmune disease results from a complex failure of immunological tolerance involving genetic, environmental and immune-regulatory factors. Restoration of immune homeostasis and control of autoreactive immune responses are important goals of modern autoimmune disease research.Homoeopathy is fundamentally characterized by individualization, in which treatment is selected according to the individual patient's characteristic symptom profile rather than the diagnostic label alone. Clinical research has produced some evidence suggesting potential effects of individualized homoeopathic treatment, while systematic reviews of rheumatological conditions have identified preliminary clinical signals. However, the existing evidence remains limited and heterogeneous.

The central proposition of this review is therefore that individualized homoeopathic therapeutics may be investigated as a potential approach to modifying immune dysregulation and restoring immune homeostasis in autoimmune disease. The specific proposition that such treatment restores self-tolerance remains a biological hypothesis requiring direct investigation.Future studies incorporating disease activity scores, autoantibody profiles, regulatory T-cell measurements, cytokine analysis and other validated biomarkers could determine whether clinical improvement is associated with measurable restoration of immune regulation. Such an approach could provide a scientifically testable bridge between the classical homoeopathic principle of individualization and contemporary understanding of immune tolerance.

## Acknowledgments:-

I acknowledge the contribution of my colleague, Dr. Vikrant Tripathi for collection of the research database for this review article. He had also contributed in writing the main manuscript.

**Ethics Statement:-**
Not applicable. This article is a narrative review and hypothesis paper and involved no human or animal participants.

**Conflict of Interest:-**
There is no conflict of interest for this review article.

## References:-


1. Goodnow CC, Sprent J, Fazekas de St Groth B, Vinuesa CG. Cellular and genetic mechanisms of self tolerance and autoimmunity. Nature. 2005;435(7042):590-597.
2. Rosenblum MD, Remedios KA, Abbas AK. Mechanisms of human autoimmunity. J Clin Invest. 2015;125(6):2228-2233.
3. Hogquist KA, Baldwin TA, Jameson SC. Central tolerance: learning self-control in the thymus. Nat Rev Immunol. 2005;5(10):772-782.
4. Klein L, Robey EA, Hsieh CS. Central CD4+ T cell tolerance: deletion versus regulatory T cell differentiation. Nat Rev Immunol. 2019;19(1):7-18.
5. Nemazee D. Mechanisms of central tolerance for B cells. Nat Rev Immunol. 2017;17(5):281-294.
6. Sakaguchi S, Yamaguchi T, Nomura T, Ono M. Regulatory T cells and immune tolerance. Cell. 2008;133(5):775-787.
7. Theofilopoulos AN, Kono DH, Baccala R. The multiple pathways to autoimmunity. Nat Immunol. 2017;18(7):716-724.
8. Wahren-Herlenius M, Dörner T. Immunopathogenic mechanisms of systemic autoimmune disease. Lancet. 2013;382(9894):819-831.
9. Sumida TS, Cheru NT, Hafler DA. The regulation and differentiation of regulatory T cells and their dysfunction in autoimmune diseases. Nat Rev Immunol. 2024;24(7):503-517.
10. Mathie RT, Lloyd SM, Legg LA, Clausen J, Moss S, Davidson JRT, et al. Randomised placebo-controlled trials of individualised homeopathic treatment: systematic review and meta-analysis. Syst Rev. 2014;3:142. doi:10.1186/2046-4053-3-142.
11. Freire de Carvalho J, Lerner A, Benzvi C. Homeopathy for rheumatological diseases: a systematic review. Eur J Rheumatol. 2024;11(3):378-384. doi:10.5152/eurjrheum.2024.23123.
12. Fisher P, Scott DL. A randomized controlled trial of homeopathy in rheumatoid arthritis. Rheumatology (Oxford). 2001;40(9):1052-1055. doi:10.1093/rheumatology/40.9.1052.