Journal Homepage: -www.journalijar.com

INTERNATIONAL JOURNAL OF ADVANCED RESEARCH (IJAR)

Article DOI:10.21474/IJAR01/24509
DOI URL: http://dx.doi.org/10.21474/IJAR01/24509

ISSN (O): 2320-5407
ISSN (P): 3107-4928

RESEARCH ARTICLE

QUANTITATIVE ASSESSMENT OF GLENOHUMERAL CARTILAGE USING MRI T2 RELAXOMETRY FOR EARLY DEGENERATION DETECTION: A PROSPECTIVE STUDY

D Rajitha and Sheila Elangovan

Manuscript Info

Manuscript History
Received: 04 September 2026
Final Accepted: 08 October 2026
Published: November 2026

Key words:-
MRI, T2 relaxometry, Glenohumeral joint, cartilage degeneration, quantitative imaging

Abstract

Background: Early detection of cartilage degeneration is essential for preventing the progression of osteoarthritis. Conventional magnetic resonance imaging (MRI) has limitations in detecting subtle biochemic al changes in cartilage.T2 relaxometry provides a quantitative approach for evaluating cartilage integrity.

Objective: To assess the utility of MRI T2 relaxometry in detecting early degenerative changes in the glenohumeral cartilage

Materials and Methods : A prospective study was conducted on 18 patients below 50 years of age presenting with shoulder pain without trauma .MRI examinations were performed using 1.5 tesla scanners from two vendors with standardized imaging protocols .T2 maps were generated using vendor –specific software and analyzed using region of interest based measurements .

Results: The mean T2 relaxation value was 43.62+/- 10.35 ms with a range of 27.32 to 60.29ms. elevated T2 values were observed in 44.4% of patients .T2 mapping demonstrated improved sensitivity in detecting cartilage abnormalities compared to conventional MRI.

Conclusion: MRI T2 relaxometry is a reliable quantitative imaging technique for early detection of cartilage degeneration and provides additional diagnostic value beyond routine MRI.

"© 2026 by the Author(s). Published by IJAR under CC BY 4.0. Unrestricted use allowed with credit to the author."

Introduction:-

Cartilage degeneration plays a signification role in the development of osteoarthritis, particularly in highly mobile joints such as the glenohumeral joint (1).early identification of degenerative changes is essential for preventing disease progression and preserving joint function (2).however, conventional magnetic resonance imaging (MRI) primarily evaluates structural abnormalities and may not detect early biochemical change sin cartilage.(3) Quantitative MRI techniques have emerged as valuable tools for assessing cartilage composition(4–6). T2 relaxometry reflects collagen matrix integrity and water content within cartilage ((7,8). Increased T2 relaxation times are associated with disruption of collagen architecture and increased hydration, which are early indicators of degeneration. ((9,10)Although extensively studied in weight-bearing joints such as the knee , the application of T2 relaxometry in shoulder cartilage remains limited (11,12). Given the clinical importance of early diagnosis, this study aims to evaluate the role of T2 relaxometry in detecting early degenerative changes in glenohumeral cartilage.

Corresponding Author:- D Rajitha

Materials and Methods:-

Study design and setting:-

A prospective study was conducted over a period of three months (January 2020 to March 2020) in the department of radiology and imaging sciences at Sri Ramachandra institute of higher education and research, Chennai, India.

Study population: a total of 18 patients below the age of 50 years presenting with shoulder pain without a history of trauma were included

Inclusion criteria:-

Patients aged <50 years
Shoulder pain without trauma

Exclusion criteria:-

History of shoulder surgery, traumatic injury, known shoulder pathology

MRI acquisition protocol :-

MRI examinations were performed using 1.5 tesla scanners from two vendors (Siemens magneto avanto and GE signa series) with dedicated shoulder coils.Routine imaging included T1 –weighted, T2 –weighted and proton density fat –saturated sequences.T2 relaxometry was performed using a multi-echo-spin-echo sequence with comparable parameters across both systems. imaging parameters were standardized to ensure consistency of quantitative measurements (table 3)

Table 3. MRI acquisition parameters across different vendors

PARAMETER	Siemens (Magnetom avanto)	GE (signa series)
Field strength	**1.5 tesla**	**1.5tesla**
Scan plane	Oblique coronal	Oblique coronal
Mode	2D	2D
Pulse sequence family	Spin echo	Spin echo
Pulse sequence	FSE-T2	FSE-T2
TR (ms)	1020	1020
TE (ms)	▪ 13.9 ▪ 27.8 ▪ 41.7 ▪ 55.6 ▪ 69.5	▪ 13.9 ▪ 27.8 ▪ 41.7 ▪ 55.6 ▪ 69.5
FOV read (mm)	160	160
FOV phase (%)	100	100
Slice thickness (mm)	3-4mm	3-4mm
Spacing (mm)	1	1
Flip angle (degree)	180	180
Matrix/phase resolution	224 x 320	224 x 320
NEX	1	1
Scan time	3:20 min	3:20 min

Post processing and T2 analysis :-

T2 maps were generated using vendor–specific platforms (syngo MR workplace for Siemens and GE AW workstation for GE systems). Despite differences in workflow, both systems employed mono-exponential fitting models.Regions of interest (ROI) were manually placed over the glenohumeral cartilage with care to avoid partial volume effects. Consistency in ROI placement was maintained across all cases.

Statistical analysis:-
Descriptive statistics were used. Mean, standard deviation, and range were calculated. The proportion of patients with elevated T2 values was determined.

Results:-

A total of 18 patients were included in the analysis .The mean T2 relaxation value was 43.62 +/- 10.35ms with a range of 27.32 to 60.29ms (table-1).Among the study population ,8 patients (44.4%) demonstrated elevated T2 values .10 patients (55.6%) had normal values (table 2).T2 relaxometry enabled detection of early cartilage abnormalities through increased relaxation values improved visualization using color-coded maps (fig.1).In several cases, conventional MRI sequences did not demonstrate abnormalities, whereas T2 mapping revealed elevated values (fig.2).The distribution of T2 relaxation values is illustrated in (fig.3)




Figure.1. T2 relaxometry color map showing increased T2 values in glenohumeral cartilage indicating early degeneration




Figure.2. conventional MRI image with corresponding T2 map demonstrating elevated relaxation values not appreciable on routine sequences




Figure.3.Graph showing distribution of T2 relaxation values among study population

Table1.summary of T2 Relaxometry values

Parameter	Value
Sample size	18
Mean t2 value	43.62ms
Standard deviation	+/-10.35ms
Range	27.32 - 60.29ms

Table 2. Distribution of patients based on T2 values

Category	Number of patients	Percentage
Elevated T2 values	8	44.4%
Normal T2 values	10	55.6%

Discussion:-

This study demonstrates the utility of MRI T2 relaxometry in detecting early degenerative changes in glenohumeral cartilage. Conventional MRI is primarily limited to morphological evaluation and may not identify early biochemical alterations The mean T2 value observed in this study (43.62+/- 10.35 ms) and the proportion of patients with elevated values (44.4%) indicate the presence of early cartilage degeneration. increased t2 values reflect disruption of collagen architecture and increased water content. T2 mapping showed improved sensitivity compared to routine MRI, particularly in cases where conventional imaging appeared normal. This highlights its importance as a complementary diagnostic tool The use of scanners from different vendors and variation of sin post-processing platforms may introduce minor variability in measurements. However, standardized acquisition parameters and consistent ROI-based analysis were employed to minimize this effect. Limitations include small sample size and lack of longitudinal follow-up. Future studies with larger populations are required to establish standardized reference values.

Conclusion:-

MRI T2 relaxometry is a reliable and non-invasive technique for detecting early degenerative changes in glenohumeral cartilage. It enhances diagnostic sensitivity compared to conventional MRI and has potential clinical value in early disease detection.

References:-

1. Thomas M, Bidwai A, Rangan A, Rees JL, Brownson P, Tennent D, et al. Glenohumeral osteoarthritis. Shoulder Elbow. 2016 Jul 1;8(3):203–14. doi:10.1177/1758573216644183 ar3845.
2. Choi JA, Gold GE. MR imaging of articular cartilage physiology. Magnetic Resonance Imaging Clinics of North America. 2011. p. 249–82. doi:10.1016/j.mric.2011.02.010 PubMed PMID: 21665090.
3. Matzat SJ, van Tiel J, Gold GE, Oei EHG. Quantitative MRI techniques of cartilage composition. Quant Imaging Med Surg. 2013 Jun;3(3):162–74. doi:10.3978/j.issn.2223-4292.2013.06.04 PubMed PMID: 23833729.
4. Casula V, Kajabi AW. Quantitative MRI methods for the assessment of structure, composition, and function of musculoskeletal tissues in basic research and preclinical applications. Magnetic Resonance Materials in Physics, Biology and Medicine. Springer Science and Business Media Deutschland GmbH; 2024. p. 949–67. doi:10.1007/s10334-024-01174-7 PubMed PMID: 38904746.
5. Lavalle S, Scapaticci R, Masiello E, Salerno VM, Cuocolo R, Cannella R, et al. Beyond the Surface: Nutritional Interventions Integrated with Diagnostic Imaging Tools to Target and Preserve Cartilage Integrity: A Narrative Review. Biomedicines. Multidisciplinary Digital Publishing Institute (MDPI); 2025. doi:10.3390/biomedicines13030570
6. Shao H, Pauli C, Li S, Ma Y, Tadros AS, Kavanaugh A, et al. Magic angle effect plays a major role in both T1rho and T2 relaxation in articular cartilage. Osteoarthritis Cartilage. 2017 Dec 1;25(12):2022–30. doi:10.1016/j.joca.2017.01.013 PubMed PMID: 28161394.
7. Lavalle S, Scapaticci R, Masiello E, Salerno VM, Cuocolo R, Cannella R, et al. Beyond the Surface: Nutritional Interventions Integrated with Diagnostic Imaging Tools to Target and Preserve Cartilage Integrity: A Narrative Review. Biomedicines. Multidisciplinary Digital Publishing Institute (MDPI); 2025. doi:10.3390/biomedicines13030570
8. Dunn TC, Lu Y, Jin H, Ries MD, Majumdar S. T2 relaxation time of cartilage at MR imaging: Comparison with severity of knee osteoarthritis. Radiology. 2004 Aug;232(2):592–8. doi:10.1148/radiol.2322030976 PubMed PMID: 15215540.
9. Baum T, Joseph GB, Karampinos DC, Jungmann PM, Link TM, Bauer JS. Cartilage and meniscal T2 relaxation time as non-invasive biomarker for knee osteoarthritis and cartilage repair procedures. Osteoarthritis and Cartilage. 2013. p. 1474–84. doi:10.1016/j.joca.2013.07.012 PubMed PMID: 23896316.
10. Wuennemann F, Kintzelé L, Braun A, Zeifang F, Maier MW, Burkholder I, et al. 3-T T2 mapping magnetic resonance imaging for biochemical assessment of normal and damaged glenoid cartilage: a prospective arthroscopy-controlled study. Sci Rep. 2020 Dec 1;10(1). doi:10.1038/s41598-020-71311-4 PubMed PMID: 32873848.
11. Verschueren J, Van Langeveld SJ, Dragoo JL, Bierma-Zeinstra SMA, Reijman M, Gold GE, et al. T2 relaxation times of knee cartilage in 109 patients with knee pain and its association with disease characteristics. Acta Orthop. 2021;92(3):335–40. doi:10.1080/17453674.2021.1882131 PubMed PMID: 33538221.