



CASE REPORT

Sirenomelia: A Case Report

Bichitrananda Roul¹, Mamita Nayak², Sagar Prusti³, Sadananda Rath⁴, Bijay Kumar Dutta⁵

1. Assistant Professor ,Dept of Anatomy,Pt.J.N.M.Medical College,Raipur
2. Senior Resident,Dept of Pathology,AIIMS,Bhubaneswar
3. Associate professor, Dept. of Anatomy,M.K.C.G.Medical College
4. Professor & H.O.D., Dept. of Anatomy,M.K.C.G.Medical College
5. Professor & H.O.D., Dept. of Anatomy,V.S.S..Medical College

Manuscript Info

Manuscript History:

Received: 15 March 2015
Final Accepted: 22 April 2015
Published Online: May 2015

Key words:

Sirenomelia, mermaid syndrome, agenesis, diabetes, vascular steal phenomenon

**Corresponding Author*

Bichitrananda Roul

Abstract

Sirenomelia, which is also known as mermaid syndrome, is an extremely rare congenital developmental disorder characterized by anomalies of the lower spine and the lower limbs. Affected infants are born with partial or complete fusion of the legs. Additional malformations may also occur including genitourinary abnormalities, gastrointestinal abnormalities, anomalies of the lumbarsacral spine and pelvis and absence or underdevelopment (agenesis) of one or both kidneys. Affected infants may have one foot, no feet or both feet, which may be rotated externally. The tailbone is usually absent and the sacrum is partially or completely absent as well. Additional conditions may occur with sirenomelia including imperforate anus, spina bifida, and heart (cardiac) malformations. Sirenomelia is often fatal during the newborn period. : The case was reported at Zenana hospital,Berhampur ,during routine delivery. There was H/O diabetes, but no H/O of epilepsy,hypertension of the mother,after the baby was born live & it was transferred to NICU & within few hours it died . The morphology was studied & seen that the lower limbs are fused up to distal end with toes frinzizing out. There were six toes found, three in each leg. Great toe being present laterally instead of medially. Popliteal fossa was present in anterior aspect . Absense of external genitalia (sex cant be determined) . A Potter's face (ocular hypertelorism, low-set ears, receding chin and flattening of the nose)Imperforate anus . No deformity found on upperlimb, trunk, abdomen, face & head morphologically.

Copy Right, IJAR, 2015.. All rights reserved

INTRODUCTION

Sirenomelia, which is also known as mermaid syndrome, is an extremely rare congenital developmental disorder characterized by anomalies of the lower spine and the lower limbs. Affected infants are born with partial or complete fusion of the legs. Additional malformations may also occur including genitourinary abnormalities, gastrointestinal abnormalities, anomalies of the lumbarsacral spine and pelvis and absence or underdevelopment (agenesis) of one or both kidneys. Affected infants may have one foot, no feet or both feet, which may be rotated externally. The tailbone is usually absent and the sacrum is partially or completely absent as well. Additional conditions may occur with sirenomelia including imperforate anus, spina bifida, and heart (cardiac) malformations. The condition is thought to be part of a spectrum including imperforate anus, sacral agenesis and sirenomelia³. Sirenomelia is often fatal during the newborn period. The exact cause of sirenomelia is unknown, most cases occur randomly for no apparent reason (sporadically). It is a lethal type of birth defect in which the two lower limbs of

the new born are attached to or fused together as one. It is otherwise called as mermaid syndrome as the two fused lower limbs look like mermaid's tail. Affected newborns do not survive more than 24 hours after birth. Only four cases of a surviving infant with sirenomelia have been reported¹⁰.

This is an uncommon congenital deformity and occurs in 1 out of every 100,000 live births in the US. The condition occurs predominantly in males, with a sex ratio of 2.7: 1²⁴. Infants suffering from this disorder die within a day or two due to the abnormal development of kidney and urinary bladder. Sirenomelia has been reported among different ethnic groups of the world and is common among infants born to pregnant women suffering from Diabetes mellitus.

Case Report :

The case was reported at Zenana hospital, Berhampur, during routine delivery. There was H/O diabetes, but no H/O of epilepsy, hypertension of the mother, after the baby was born live & it was transferred to ICU & within few hours it died. Then with the permission of the ethical committee of medical college, the case was transferred to department of Anatomy, M.K.C.G Medical college for academic purpose, with prior permission from H.O.D., Anatomy, MKCGMC & Director, Zenana hospital & with consent of the parents,

The morphology was studied (fig-1,2 &3) & seen that the lower limbs are fused up to distal end with toes fringing out. There are six toes found, three in each leg. Great toe being present laterally instead of medially. Popliteal fossa was present in anterior aspect. Absence of external genitalia (sex can't be determined). A Potter's face (ocular hypertelorism, low-set ears, receding chin and flattening of the nose) Imperforate anus. No deformity found on upper limb, trunk, abdomen, face & head morphologically.

Internal abnormalities : normal liver, renal agenesis, absence bladder found

For further study radiological study was done. X-ray of the body including the lower limb was done. X-ray findings (fig-4): both the femurs present, both tibia present but single fibula present. Partial sacral agenesis present.

Discussion :

This is an uncommon congenital deformity and occurs in 1 out of every 100,000 live births in the US. Infants suffering from this disorder die within a day or two due to the abnormal development of kidney and urinary bladder. Sirenomelia has been reported among different ethnic groups of the world and is common among infants born to pregnant women suffering from Diabetes mellitus.

The abnormalities of embryologic development leading to the sirenomelia sequence occur about the fourth week of gestation, and concern the medio-posterior mesodermic axis and the caudal blastem.¹⁶ At this gestational age, the cloaca (cloaca) is formed, the kidneys are located in the pelvis, while the gonads are intra-abdominal. It seems logical that any damage to the caudal extremity of the embryo would affect the development of the external and internal genital organs (except the gonads, which are intra-abdominal), the terminal bowel, the bladder, the kidneys and the pelvic bones⁴.

In humans, fusion of the legs in sirenomelia occurs in a spectrum of morphologies, ranging from the mildest cases in which all bones of the two fused lower limbs are discernible, to the most severe cases in which there is no indication that the single rudimentary lower limb that is present derives from the fusion of two

Infants of diabetic mothers have two to three times the average incidence of congenital anomalies. These include neural tube defects, cardiac defects (transposition of the great vessels, coarctation of the aorta, VSD, ASD, cardiomyopathy), situs inversus, renal anomalies (hydronephrosis, renal agenesis, multicystic dysplastic kidney, duplication of the renal tracts), intestinal atresia, and forms of caudal regression including sacral agenesis^{17,34}.

Lynch and Wright¹⁸ reported a case where the mother had diabetes and the infant had sirenomelia with renal agenesis and an absent right radius. There was also a hypertrophic cardiomyopathy and a bicuspid pulmonary valve. Sirenomelia is thought to occur in about 1 in 65,000 live births and may be more common in one of monozygotic twins. It bears resemblance to the mermaid Greek mythology. Affected individuals exhibit a variable range of

defects, including hypoplasia and or fusion of lower limbs, vertebral abnormalities, renal agenesis, imperforate anus, and anomalies of the genital organs²⁷

Several etiologic factors have been suggested; however none of them explains all the anomalies encountered in the sirenomelia sequence⁴:

Maternal diabetes is the only maternal disease known to be associated with sirenomelia (2% of cases), although this is associated more frequently with CRS (22% of cases). Teratogens: retinoic acid, cadmium and cyclophosphamide have been implicated in the genesis of sirenomelia in mice and hamsters. Genetic factors: several family cases of CRS have been reported, as well as some cases of minor vertebral abnormalities in the parents, suggesting a possible genetic transmission. Several modalities of transmission have been suggested: dominant sex-linked transmission, multifactor polygenetic transmission, and dominant autosomic transmission with a variable expression and an attenuated penetrance. The vascular steal theory: this theory was initially suggested in 1927 by Kampmeier and was called "nutritional deficit". It was reintroduced in 1986 by Stevenson³⁶ under the term "vascular steal". Stevenson relied on observations taken from the autopsies of 11 cases of sirenomelia, He observed a mega-artery ensures the function of both umbilical arteries and diverts the blood flow on the embryo's caudal extremity toward the placenta, causing a nutritional deficit and a lack of development of the caudal extremity. Sirenomelia is quite often accompanied by a unique umbilical artery.³⁷ Our observation of a single umbilical artery, renal agenesis and the absence of renal arteries might support this theory,. The gonads are vascularized, in principle, by collateral arteries of the abdominal aorta starting from the top of the renal arteries.. The vascular steal theory also fails to explain the frequent association of sirenomelia with other abnormalities such as cranial, cardiac and esophageal defects.³⁵

Previous reports described the extent of the malformation complex in association with sirenomelia. M Al-Haggat et al¹⁹ reported one case of a fetus of sirenomelia sequence with Potters syndrome which showed oligohydramnios and symelia apus. The infant showed absent urinary tract and external genitalia, the legs were fused by skin and had separate bones associated with Potter's syndrome. The mother had a history of gestational diabetes mellitus.. Anis Fadhlaoui et al⁴ report a case of sirenomelia sequence observed in an incident of preterm labor during the 29th gestational week. According to some authors, this syndrome should be classified separately from caudal regression syndrome and is likely to be the result of an abnormality taking place during the fourth gestational week, causing developmental abnormalities in the lower extremities, pelvis, genitalia, urinary tract and digestive organs. McCoy et al²³ reported an unusual case with two normal kidneys. Perez-Aytes et al,²⁵ reported an unusual case where there appeared to be partial fusion of the lower limbs associated with absent feet, but other features of caudal regression and a single aberrant umbilical artery. Akbiyik et al.² reported male twins, one of which had sirenomelia and the other just an imperforate anus. Unfortunately zygosity was not established. Shonubi et al³¹ reported sirenomelia in 1 of 2 monozygotic twins. Duncan et al.,¹¹ reviewed 445 cases ascertained for sacrococcygeal malformations and found that 12% had sirenomelia, 27% VATER association and 34% what they called the sacrococcygeal dysgenesis association. Guidera et al¹³ reported a case of orthopaedic abnormalities and pelvic dysplasia, fused lower extremities, and osseous fusion of the calcanei, but with normal femurs, tibias, and fibulas. This patient also had significant cardiac and renal abnormalities. Stevenson *et al*³³ reported in 1986 that the aetiology of sirenomelia was explained by vascular steal theory. They documented the common feature of a single large artery arising from high in the abdominal cavity, which they thought diverted nutrients from the caudal end of the embryo in a vascular 'steal' phenomenon. The associated anomalies are variable and involved multiple organ systems. The most common anomalies include a single umbilical artery, malformation of urinary tract, lower gastrointestinal tract and external genitalia³⁴.

Acknowledgement :

We are very much thankful to CMO, & other faculties of Zenana Hospital for their kind support .We are also very much thankful Dr. Giri of Sahara Diagnostics for his wholehearted support.we are also very much thankful to the authors from which we have taken references.

References

- [1] Adra A, Cordero D, Mejides A, et al. Caudal regression syndrome: Etio-pathogenesis, prenatal diagnosis and perinatal management. *Obstet Gynecol Surv.* 1994;49(7):508–76. [[PubMed](#)]
- [2]. Akbiyik F, Balci S, Akkoyun I, Aktas D, Cakmak O. Type 1 sirenomelia in one of male twins, with imperforate anus in the other male twin. *Clin Dysmorphol.* 2000;9:227–229. [PubMed](#). [[PubMed](#)]
- [3]. Murphy JJ, Fraser GC, Blair GK. Sirenomelia: The case of the surviving mermaid. *J Pediatr Surg.* 1992;27:1265–1268. [[PubMed](#)]
- [4]. [Anis Fadhlaoui](#) et al **The Sirenomelia Sequence: A Case History** *Clin Med Insights Case Rep.* 2010; 3: 41–49. Published online 2010 Aug 13.
- [5]. . Banerjee A, Faridi MM, Banerjee TK, Mandal RN, Aggarwal A. Sirenomelia. *Indian J Pediatr.* 2003;70(7):589–91. [[PubMed](#)]
- [6]. Buekens P. Invited commentary: prenatal glucose screening and gestational diabetes. *Am J Epidemiol.* 2000;152:1015–1016.
- [7]. Chen CP, Shih S-L, Liu FF, Jan SW. Cebocephaly, alobar holoprosencephaly, spina bifida, and sirenomelia in a stillbirth. *J Med Genet.* 1997;34:252–255. [PubMed](#). [[PMC free article](#)] [[PubMed](#)]
- [8]. Clarke LA, Stringer DA, Fraser GC, Yong SL. Long term survival of an infant with sirenomelia. *Am J Med Genet.* 1993 Feb 1;45(3):292–6. [[PubMed](#)]
- [9]. Dilorenzo M, Brandt ML, Veilleux A. Sirenomelia in an identical twin. A case report. *J Pediatr Surg.* 1991;26:1334. [[PubMed](#)]
- [10]. Duhamel B. From the mermaid to anal imperforation: The syndrome of caudal regression. *Arch Dis Child.* 1961;36:153–155. [[PMC free article](#)] [[PubMed](#)]
- [11]. Duncan PA, Shapiro LR. Sirenomelia and VATER association: possible interrelated disorders with common embryologic pathogenesis. *Dysmorph Clin Genet.* 1988;2:96–103. [PubMed](#).
- [12]. Gardner NJ, Breuer AC. Anomalies of heart, spleen, kidneys, gut and limbs may result from an overdistended neural tube. A hypothesis. *Pediatrics.* 1980;65:508. [[PubMed](#)]
- [13]. Guidera KJ, Raney E, Ogden JA, et al. Caudal regression: A review of seven cases, including the mermaid syndrome. *J Pediatr Orthop.* 1991;11:743–747. [[PubMed](#)]
- [14]. . Harika G, Gabriel R, Bory JP, et al. La sirenomelie. *J Gynecol Obstet Biol Reprod.* 1995;24:49–51. [[PubMed](#)]
- [15]. Jarrett RJ. Gestational diabetes: anon-entity? *Br Med J.* 1993;306:37–38. [[PMC free article](#)] [[PubMed](#)]
- [16]. Jones KL. *Smith's Recognizable Pattern of Human Malformation.* 4th edition. Philadelphia: Saunders; 1988. p. 574.
- [17]. Loffredo CA, Wilson PD, Ferencz C. Maternal diabetes: an independent risk factor for major cardiovascular malformations with increased mortality of affected infants. *Teratology.* 2001;64:98–106. [PubMed](#). [[PubMed](#)]
- [18]. Lynch SA, Wright C. Sirenomelia, limb reduction defects, cardiovascular malformation, renal agenesis in an infant born to a diabetic mother. *Clin Dysmorphol.* 1997;6:75–80. [PubMed](#). [[PubMed](#)]
- [19]. [M Al-Haggag](#) et al. **Sirenomelia (symelia apus) with Potter's syndrome in connection with gestational diabetes mellitus: a case report and literature review,** *Afr Health Sci.* 2010 Dec; 10(4): 395–399
- [20]. Managoli S, Chaturvedi P, Vilhekar KY, Iyenger J. Mermaid syndrome with amniotic band disruption. *Indian J Pediatr.* 2003;70:105–107. [PubMed](#). [[PubMed](#)]
- [21]. . Maroteaux P, Le Merrer M. *Maladies osseuses de l'enfant.* 4th edn. Paris: Medicine-Science, Flammarion; 2002. pp. 266–268
- [22]. Martinez-Frias ML, Garcia A, Bermejo EM. Cyclopia and Sirenomelia in a live born infant. *J Med Genet.* 1998;35:263. [[PMC free article](#)] [[PubMed](#)]
- [23]. . McCoy MC, Chescheir NC, Kuller JA, et al. A fetus with sirenomelia, omphalocele, and meningomyelocele, but normal kidneys. *Teratology.* 1994;50:168–171. [PubMed](#). [[PubMed](#)]
- [24]. . Murphy JJ, Fraser GC, Blair GK. Sirenomelia: The case of the surviving mermaid. *J Pediatr Surg.* 1992;27:1265–1268. [[PubMed](#)]
- [25]. Perez-Aytes A, Montero L, Gomez J, Paya A. Single aberrant umbilical artery in a fetus with severe caudal defects: Sirenomelia or Caudal dysgenesis. *Am J Med Genet.* 1997;69:409–412. [PubMed](#). [[PubMed](#)]
- [26]. Rodriguez JI, Palacios J. Craniorachischisis totalis and sirenomelia. *Am J Med Genet.* 1992;43:732–736. [PubMed](#). [[PubMed](#)]

- [27]. Sadler T W. Langman's Medical Embryology. 9th edition. Philadelphia: Lippincott Williams and Wilkins; 2004. p. 79.
- [28]. Sarpong S, Headings V. Sirenomelia accompanying exposure of the embryo to cocaine. South Med J. 1992;85:545-547. [[PubMed](#)]
- [29]. Selig AM, Benacerraf B, Greene MF, et al. Renal dysplasia, megalocystis, and sirenomelia in four siblings. Teratology. 1993;47:65-71. PubMed. [[PubMed](#)]
- [30]. Sells CJ, Robinson NM, Brown Z, Knopp RH. Long-term developmental follow-up of infants of diabetic mothers. J Pediatr. 1994;125:s9-s17. PubMed. [[PubMed](#)]
- [31]. Shonubi AM, Salami BA, Falade AG, Ejiwunmi AB. Sirenomelia in one of Nigerian mono chorionic twins. Afr J Med Med Sci. 2002;31:377-379. PubMed. [[PubMed](#)]
- [32] Stanton MP, Penington EC, Huston JM. A surviving infant with Sirenomelia (Mermaid syndrome) associated with absent bladder. J Pediatr Surg. 2003;38(8):1266-8. [[PubMed](#)] [33].Stevenson RE, Phelan MC, Saul RA. Defects of the abdominal wall: association with vascular steal. Proc Gr Genet Center. 1989;8:15-21. PubMed.
- [34]. Stevenson RE, Jones KL, Phelan MC, et al. Vascular steal: the pathogenic mechanism producing sirenomelia and associated defects of viscera and soft tissues. Pediatrics. 1986;78:451-7. [[PubMed](#)]
- [35].Stocker JT, Heifetz SA. Sirenomelia: A morphological study of 33 cases and review of the literature. Perspect Pediatr Pathol. 1987;210:7-50. [[PubMed](#)]
- [36] Valenzano M, Paoletti R, Rossi A, et al. Sirenomelia. Pathological features, antenatal ultrasonographic clues and a review of current embryogenic theories. Hum Reprod Update. 1999;5(1):82-6. [[PubMed](#)].
- [37]. Stockers JT, Heifetz SA. Sirenomelia : a morphological study of 33 cases and review of the literature. Perspect Pediatr Pathol. 1987;10:7-50. [[PubMed](#)]

FIG-1



FIG-2



FIG -3



FIG-4



