

 <p>ISSN NO. 2320-5407</p>	<p>Journal Homepage: -www.journalijar.com</p> <p>INTERNATIONAL JOURNAL OF ADVANCED RESEARCH (IJAR)</p> <p>Article DOI:10.21474/IJAR01/14256 DOI URL: http://dx.doi.org/10.21474/IJAR01/14256</p>	 <p>INTERNATIONAL JOURNAL OF ADVANCED RESEARCH (IJAR) ISSN 2320-5407</p> <p>Journal Homepage: http://www.journalijar.com Article DOI:10.21474/IJAR01/14256</p>
---	--	--

RESEARCH ARTICLE

ANALYSIS OF THE SECONDARY METABOLITES OF VERNONIA AMYGDALINA AND THE ANTIMICROBIAL EFFECTS ON PATHOGENIC ORGANISMS

Rose I. Nwabueze

Federal College of Education, Technical, Umuze.

Manuscript Info

Manuscript History

Received: 18 December 2021

Final Accepted: 20 January 2022

Published: February 2022

Key words: -

Bitter Leaf, Phytochemical, Ethanol, Acetone, Antimicrobial

Abstract

Medicinal plants have assumed the basis of traditional treatment and have proven effective in human health care worldwide. There is an increasing interest in complementary and alternative therapies to treat various illnesses. The continued consumption of antibiotics with little or no impact on bacteria has negatively affected health care delivery. Bitter leaf (*Vernonia amygdalina*) is a common shrub belonging to the family Asteraceae and a perennial shrub usually cultivated as a homestead vegetable and fodder tree in various parts of Nigeria. The current study examined the effect of bitter leaf's phytochemical constituents and antimicrobial activity on *Candida albicans*, *Pseudomonas aeruginosa*, and *Staphylococcus aureus*. The result of the phytochemical analysis performed on the root and stem bark of *V. amygdalina* revealed the presence of alkaloids, steroids, tannin, flavonoids, saponins, and phenol. However, the investigation revealed that tannins and phenol were mainly deposited in the plant's stem bark. Also, the antimicrobial analysis conducted showed that *S. aureus* was more vulnerable to the ethanol, acetone, and hot aqueous extracts of *V. amygdalina*. At the same time, *C. Albicans* and *P. aeruginosa* showed sensitivity on exposure to ethanol and acetone but did not react to the hot aqueous extracts. The study concludes that the observed biological reactions indicated in the aqueous extracts of *V. amygdalina* validate the traditional application of this plant as an alternative antibiotic.

Copy Right, IJAR, 2022, All rights reserved.

Introduction:-

Over the years, medicinal plants have played a substantial role in Nigeria's health sector. Medicinal plants have presumed the basis of traditional medicine and have proven active in human health care worldwide (Asif, 2013; Hamayun et al., 2006; Johnson et al., 2015; Kumar et al., 2013; Tripathi & Pandey, 2017). There is an increasing interest in the role of alternative medicines to treat various illnesses (Albejo et al., 2015; Ali et al., 2020; Baars et al., 2019; James et al., 2018; Mordeniz, 2019). Plant's secondary metabolites and phytochemicals are the essential elements in the curative success of plants. However, the mechanisms underlying most medicinal plant-related cure is still unclear. The success of complementary and alternative medicines in treating infectious diseases indicates the effectiveness of plants in treating different types of fungal, bacterial, and other infections (el Hajj & Holst, 2020; Kamatenesi-Mugisha et al., 2008; Kayanja, 2008; Martin & Ernst, 2003; Wang et al., 2014). Complementary and alternative medicine is increasingly employed to curb the infectious disease in Nigeria. The trend has continued to

attract wide patronage following most conventional antibiotics' ineffectiveness due to the antimicrobial resistance by bacteria.

The growing occurrence of antimicrobial drug resistance by most bacteria has triggered a public health concern (Bennani et al., 2020; Buchy et al., 2020; Christaki et al., 2020; Dadgostar, 2019; Didier et al., 2019; Lowy, 2003; Taneja & Sharma, 2019; Vidovic & Vidovic, 2020; WHO, 2014). The continued consumption of antibiotics with little or no impact on bacteria has negatively affected health care delivery worldwide. Perhaps, extensive research has been dedicated to searching for an alternative remedy to bacterial infections. (Alsheikh et al., 2020; Chandra et al., 2017; Esmael et al., 2020; Nawab et al., 2020; Santos et al., 2019; Septama et al., 2020; Shin et al., 2018; Singh et al., 2020; Srivastava et al., 2014; Yang et al., 2018). The research in alternative to the chemically powered antibiotics has focused on plant bioactive components. Phytochemical contents of many plants such as alkaloids, essential oils, resins, lactose, and saponins have proved relevant in curbing bacterial activities (Igbinauwa et al., 2012; Mujeib et al., 2014; Muthu et al., 2006; Steenkamp et al., 2004; Umaru et al., 2019). Thus, this study is aimed to ascertain the aqueous constituents and the antimicrobial properties of a bitter leaf on microorganisms.

Bitter leaf (*Vernonia amygdalina*) is a popular shrub belonging to the family Asteraceae (Okwuzu et al., 2017). *V. amygdalina* is a perennial shrub commonly cultivated as a homestead vegetable and fodder tree (Ndaeyo, 2007). It is widely found in various parts of Nigeria. Bitter leaf is one of the leafy vegetables crucial in curbing micronutrient malnutrition (Tonukari et al., 2015). The plant is an essential protective vegetable and a popular ingredient used in making soup (Onabanjo & Oguntola, 2003; Shokunbi et al., 2011; Uchechukwu, 2011). The leaves are widely considered appetizers and used to aid digestion (Oyeyemi et al., 2018). It is beneficial for human health care and the treatment of various diseases. Bitter leaf is widely recognized due to the vast bioactive compounds obtained across multiple plant parts (Oyeyemi et al., 2018).

Research on the pharmacological potentials of *V. amygdalina* indicates that the plant possesses immunomodulatory activities (Setiawan et al., 2018), antimicrobial properties (Ijeh & Ejike, 2011; Salawu et al., 2011), antibacterial activity (Habtamu & Melaku, 2018), insecticidal properties (Green et al., 2017; Ileke Kayode & Olabimi Isaac, 2019), Antidiabetic (Owen et al., 2011), Anthelmintic (Nalule et al., 2011), Anticancer (Izevbigie et al., 2008; Yedjou et al., 2015), Anthelmintic and Antimalarial properties (Obboh, 2006), anxiolytic, sedative and hypothermic effects (Olorunfemi et al., 2014). Although *V. amygdalina* has been widely studied, its bioactive components and antimicrobial effects are well documented. This study aimed to further evaluate existing evidence from the literature on the antimicrobial effects of *V. amygdalina* on bacteria. Thus, the current study's primary purpose is to determine the phytochemical contents and the antimicrobial activity of *V. amygdalina* on *Staphylococcus aureus*, *Pseudomonas aeruginosa*, and *Candida albicans*.

Materials and Methods:-

Collection and Preparation of Plant Materials

Fresh samples of *V. amygdalina* plant were collected from the open market in Awka, Anambra State, Nigeria, and were taken to the lab for proper identification and authentication. The samples were washed and prepared according to the method described in (Alara et al., 2019).

Sample Extraction

Aqueous Extract

Ten grams of the ground sample of the plant leaf and stem bark were extracted was added to 100ml of sterile distilled. The plant's aqueous components were removed following the method adopted by (Abdulmalik et al., 2016; Ali et al., 2017).

Phytochemical Screening of *V. amygdalina*

Phytochemical screening of the *V. amygdalina* stem bark and root extracts was conducted to ascertain bioactive components such as alkaloids, saponins, tannins, steroids, phenols, and flavonoids using the standard qualitative method as previously described by Trease and Evans (1989).

Sterility Test of the Plant Extracts

After sterilization, the plant's extracted constituents were tested for sterility by inoculating 1 mL of each extract on sterilized nutrient agar incubated at 37°C for 24 hours. Indeed, the plates were carefully observed for growth.

Standardization of the Bacterial Cell Suspension

McFarland standard was adopted to test for the standardization of the bacteriological suspension. Colonies of the tested organism were picked into a disinfected test tube containing sterile nutrient broth and incubated for one day.

Determination of Antimicrobial Activities

The crude extracts' antimicrobial activity was conducted using the agar well diffusion method described by (Gashe & Zeleke 2017) with few modifications.

Table 1:- Table showing the phytochemical contents of the root and stem bark of *V. amygdalina*.

Phytochemical compounds	Root	Stem bark
Alkaloids	++	++
Saponins	++	+++
Phenolic	-	+++
Steroid	+++	+
Tannin	-	+++
Flavonoids	++	+

Key: + = Positive, ++ = Moderate, +++ = High, - = Negative

The above table shows the outcome of the aqueous screening conducted on the root and stem bark of *V. amygdalina*, indicating that the plant contains a significant number of alkaloids, tannin, steroids, flavonoids, phenol, and saponins. However, the screening revealed tannins and phenols in the stem bark and not the plant's root.

Table 2:- Table showing the sensitivity parameter of the organisms to ethanol extract.

Organisms	Stem bark	%	Root	%
<i>C. Albicans</i>	8mm	36.89	2mm	12.45
<i>P. aeruginosa</i>	5mm	18.21	1mm	6.78
<i>S. aureus</i>	8mm	37.10	7mm	36.26

Table showing the observed zone of inhibition of ethanol extract on *V. amygdalina* stem bark and root against some pathogenic organisms. For sensitivity, exposing the microorganisms to the aqueous extract of stem bark revealed the same 8mm diameter of inhibition zone (36.89% and 37.10%), respectively, for *C. Albicans* and *S. aureus*. However, *P. aeruginosa* possessed a minimal 5mm (6.78%) diameter inhibition zone than *C. Albicans* and *S. aureus*. The test on an ethanol extract of the root shows that *S. aureus* produced a high diameter of inhibition zone of 7mm (36.26%) while *C. Albicans* had lower with 3mm (12.45%) inhibition zone diameter.

Table 3:- Table showing the sensitivity parameter of an organism to an acetone extract of the plant.

Organisms	Stem bark	%	Root	%
<i>C. Albicans</i>	8mm	21.38	7mm	41.19
<i>P. aeruginosa</i>	15mm	31.36	3mm	8.82
<i>S. aureus</i>	15mm	36.10	5mm	26.16

The table above shows the test organisms' sensitivity parameter when exposed to an acetone extract of *V. amygdalina* stem bark and root. The result indicated a high diameter of inhibition zones of 15mm for *P. aeruginosa* and *S. aureus* (31.36% and 36.10%), respectively, when tested with acetone stem bark extract of *V. amygdalina*. However, *C. Albicans* produced a reduced diameter of inhibition zones of 8mm (21.38%). Furthermore, it was revealed that *C. Albicans* had an increased diameter of inhibition zones of 7mm (41.19%) when subjected to acetone root extract of *V. amygdalina*.

Table 4:- Table showing the sensitivity position of the test organisms to hot aqueous extract.

Organisms	Stem bark	%	Root	%
-----------	-----------	---	------	---

<i>C. Albicans</i>	r	-	r	-
<i>P. aeruginosa</i>	r	-	r	-
<i>S. aureus</i>	8mm	96.10	r	-

The organisms were exposed to hot aqueous extract of the plant. Only one pathogenic organism (*S. aureus*) exhibited a reaction with the 8mm diameter of inhibition zone. Thus, other organisms maintained a resistant position towards the extract.

Discussion:-

The current study was conducted to assess bitter leaf's phytochemical constituents and antimicrobial activity on *Candida albicans*, *Pseudomonas aeruginosa*, and *Staphylococcus aureus*. The result of the phytochemical analysis performed on the root and stem bark of *V. amygdalina* revealed the presence of alkaloids, tannin, phenol, steroids, flavonoids, and saponins. However, it was found that phenol and tannins were mainly deposited in the plant's stem bark. Thus, the study is consistent with previous studies (Alara et al., 2019; Asfere et al., 2018; Oyeyemi et al., 2018). The plant's bioactive content has been implicated in the antimicrobial potentials of *V. amygdalina* (Jin et al., 2017; Nenaah, 2013).

Furthermore, the antimicrobial analysis revealed that *S. aureus* was more vulnerable to the ethanol, acetone, and hot aqueous extracts of *V. amygdalina*. As shown in the tables, this is indicated in the increased size of the inhibition zones' diameter. However, *P. aeruginosa* and *C. Albicans* showed sensitivity to ethanol and acetone exposure. However, they failed to react to the hot aqueous extracts. Consistent with (Adetutu et al., 2011; Moreno et al., 2006), the findings affirmed ethanol and acetone extracts' antimicrobial potentials compared to aqueous extracts.

Conclusion:-

The present study assessed the antimicrobial potentials of *V. amygdalina* extracts on *Candida albicans*, *Pseudomonas aeruginosa*, and *Staphylococcus aureus*. The result confirmed that the plant's stem bark and root extracts possess antimicrobial tendencies. Thus, the observed biological reactions shown by the aqueous extracts of *V. amygdalina* validate the traditional adaptation of this plant as an alternative antibiotic.

References:-

1. Abdulmalik, O., Oladapo, O. O., & Bolaji, M. O. (2016). Effect of aqueous extract of *Vernonia amygdalina* on atherosclerosis in rabbits. *ARYA Atherosclerosis*, 12(1).
2. Adetutu, A., Morgan, W. A., & Corcoran, O. (2011). Ethnopharmacological survey and in vitro evaluation of wound-healing plants used in South-western Nigeria. *Journal of Ethnopharmacology*, 137(1). <https://doi.org/10.1016/j.jep.2011.03.073>
3. Alara, O. R., Abdurahman, N. H., Ukaegbu, C. I., & Kabbashi, N. A. (2019). Extraction and characterization of bioactive compounds in *Vernonia amygdalina* leaf ethanolic extract comparing Soxhlet and microwave-assisted extraction techniques. *Journal of Taibah University for Science*, 13(1). <https://doi.org/10.1080/16583655.2019.1582460>
4. Albejo, B., Endale, M., Kibret, B., & Anza, M. (2015). Phytochemical investigation and antimicrobial activity of leaves extract of *Vernonia Auricularia* Hiern. *Journal of Pharmacy and Pharmacognosy Research*, 3(6).
5. Ali, M., Yahaya, A., Zage, A., & Yusuf, Z. (2017). In-vitro antibacterial activity and phytochemical screening of *Psidium guajava* on some enteric bacterial isolates of public health importance. *Journal of Advances in Medical and Pharmaceutical Sciences*, 12(3). <https://doi.org/10.9734/jamps/2017/31126>
6. Ali, S., Ullah, S., Paudyal, V., Ali, M., Khalid Tipu, M., & Ur-Rehman, T. (2020). Complementary and Alternative Medicines for the Treatment of Hepatitis C: Perspectives of Users and CAM Practitioners. *Evidence-Based Complementary and Alternative Medicine*, 2020. <https://doi.org/10.1155/2020/3932690>
7. Alsheikh, H. M. al, Sultan, I., Kumar, V., Rather, I. A., Al-sheikh, H., Jan, A. T., & Haq, Q. M. R. (2020). Plant-based phytochemicals as a possible alternative to antibiotics in combating bacterial drug resistance. In *Antibiotics* (Vol. 9, Issue 8). <https://doi.org/10.3390/antibiotics9080480>
8. Asfere, Y., Kebede, A., & Muthuswamy, M. (2018). In-vitro antimicrobial activities and phytochemical screening of *Calotropis Procera* (Ait.) and *Vernonia amygdalina* (Del.) Extract Against Some Medically Important Pathogenic Bacteria. *American Journal of Bioscience and Bioengineering*, 6(6).
9. Asif, M. (2013). The anticonvulsant potential of some medicinal plants and their beneficial properties. *Tang Humanitas Medicine*, 3(4). <https://doi.org/10.5667/tang.2013.0012>

10. Baars, E. W., Zoen, E. B. van, Breitzkreuz, T., Martin, D., Matthes, H., Schoen-Angerer, T. von, Soldner, G., Vagedes, J., Wietmarschen, H. van, Patijn, O., Willcox, M., Flotow, P. von, Teut, M., Ammon, K. von, Thangavelu, M., Wolf, U., Hummelsberger, J., Nicolai, T., Hartemann, P., ... Huber, R. (2019). The contribution of complementary and alternative medicine to reduce antibiotic use: A narrative review of health concepts, prevention, and treatment strategies. In *Evidence-based Complementary and Alternative Medicine* (Vol. 2019). <https://doi.org/10.1155/2019/5365608>
11. Bennani, H., Mateus, A., Mays, N., Eastmure, E., Stärk, K. D. C., & Häslar, B. (2020). Overview of evidence of antimicrobial use and antimicrobial resistance in the food chain. In *Antibiotics* (Vol. 9, Issue 2). <https://doi.org/10.3390/antibiotics9020049>
12. Buchy, P., Ascioğlu, S., Buisson, Y., Datta, S., Nissen, M., Tambyah, P. A., & Vong, S. (2020). Impact of vaccines on antimicrobial resistance. In *International Journal of Infectious Diseases* (Vol. 90). <https://doi.org/10.1016/j.ijid.2019.10.005>
13. Chandra, H., Bishnoi, P., Yadav, A., Patni, B., Mishra, A. P., & Nautiyal, A. R. (2017). Antimicrobial resistance and the alternative resources emphasize plant-based antimicrobials - A review. In *Plants* (Vol. 6, Issue 2). <https://doi.org/10.3390/plants6020016>
14. Christaki, E., Marcou, M., & Tofarides, A. (2020). Antimicrobial resistance in bacteria: Mechanisms, evolution, and persistence. In *Journal of Molecular Evolution* (Vol. 88, Issue 1). <https://doi.org/10.1007/s00239-019-09914-3>
15. Dadgostar, P. (2019). Antimicrobial resistance: implications and costs. In *Infection and Drug Resistance* (Vol. 12). <https://doi.org/10.2147/IDR.S234610>
16. Didier, K., Salomé, K., Joanne, B., Pierre, A., Fernand, G., Bénédicte, K.-K., Brice, S., Michel, F., & Joëlle, Q.-L. (2019). Chemical composition, in vitro antioxidant and antiparasitic properties of the essential oils of three plants used in traditional medicine in Benin. *Journal of Medicinal Plants Research*, 13(16). <https://doi.org/10.5897/jmpr2019.6791>
17. el Hajj, M., & Holst, L. (2020). Herbal Medicine Use During Pregnancy: A Review of the Literature with a Special Focus on Sub-Saharan Africa. In *Frontiers in Pharmacology* (Vol. 11). <https://doi.org/10.3389/fphar.2020.00866>
18. Esmael, A., Hassan, M. G., Amer, M. M., Abdelrahman, S., Hamed, A. M., Abd-raboh, H. A., & Foda, M. F. (2020). Antimicrobial activity of certain natural-based plant oils against antibiotic-resistant acne bacteria. *Saudi Journal of Biological Sciences*, 27(1). <https://doi.org/10.1016/j.sjbs.2019.11.006>
19. Green, P. W. C., Belmain, S. R., Ndakidemi, P. A., Farrell, I. W., & Stevenson, P. C. (2017). Insecticidal activity of *Tithonia diversifolia* and *Vernonia amygdalina*. *Industrial Crops and Products*, 110. <https://doi.org/10.1016/j.indcrop.2017.08.021>
20. Habtamu, A., & Melaku, Y. (2018). Antibacterial and antioxidant compounds from the flower extracts of *Vernonia amygdalina*. *Advances in Pharmacological Sciences*, 2018. <https://doi.org/10.1155/2018/4083736>
21. Hamayun, M., Khan, S. A., Kim, H. Y., Chae, I. N., & Lee, I. J. (2006). Traditional knowledge and ex situ conservation of Swat Kohistan, Pakistan, threatened medicinal plants. *International Journal of Botany*, 2(2). <https://doi.org/10.3923/ijb.2006.205.209>
22. Igbinuwa, P., Usifoh, C., & Ugwu, C. (2012). Phytochemical analysis and toxicological evaluation of the methanolic extract of *Jatropha tanjorensis* leaf. *Journal of Pharmacy & Bioresources*, 8(2). <https://doi.org/10.4314/jpb.v8i2.4>
23. Ijeh, I. I., & Ejike, C. E. C. (2011). Current perspectives on the medicinal potentials of *Vernonia amygdalina* Del. In *Journal of Medicinal Plants Research* (Vol. 5, Issue 7).
24. Ileke Kayode, D., & Olabimi Isaac, O. (2019). Insecticidal Activities of *Chromolaena odorata* and *Vernonia amygdalina* leaf extracts against *Anopheles gambiae* [Diptera: Culicidae]. *International Journal of Tropical Diseases*, 2(1). <https://doi.org/10.23937/ijtd-2017/1710018>
25. Izevbigie, E., Howard, C., & Lee, K. (2008). *V. Amygdalina*: Folk Medicine, Analysis, and Potential Application for Cancer Treatment. *Current Pharmaceutical Analysis*, 4(1). <https://doi.org/10.2174/157341208783497579>
26. James, P. B., Wardle, J., Steel, A., & Adams, J. (2018). Traditional, complementary and alternative medicine use in Sub-Saharan Africa: A systematic review. *BMJ Global Health*, 3(5). <https://doi.org/10.1136/bmjgh-2018-000895>
27. Jin, Z., Gao, L., Zhang, L., Liu, T., Yu, F., Zhang, Z., Guo, Q., & Wang, B. (2017). Antimicrobial activity of saponins produced by two novel endophytic fungi from *Panax notoginseng*. *Natural Product Research*, 31(22). <https://doi.org/10.1080/14786419.2017.1292265>
28. Johnson, M., Kolawole, O. S., & Olufunmilayo, L. A. (2015). Phytochemical analysis, in vitro evaluation of the antioxidant and antimicrobial activity of methanolic leaf extract of *Vernonia amygdalina* (bitter leaf) against

- Staphylococcus aureus* and *Pseudomonas aeruginosa*. *International Journal of Current Microbiology and Applied Sciences*, 4(5).
29. Kamatenesi-Mugisha, M., Oryem-Origa, H., Odyek, O., & Makawiti, D. W. (2008). Medicinal plants treat fungal and bacterial infections in and around Queen Elizabeth Biosphere Reserve, western Uganda. *African Journal of Ecology*, 46(SUPPL. 1). <https://doi.org/10.1111/j.1365-2028.2008.00935.x>
 30. Kayanja, F. I. B. (2008). Biodiversity in inhabited areas of Eastern Africa. *African Journal of Ecology*, 46(1).
 31. Kumar, G., Jalaluddin, M., Rout, P., Mohanty, R., & Dileep, C. L. (2013). Emerging trends of herbal care in dentistry. In *Journal of Clinical and Diagnostic Research* (Vol. 7, Issue 8). <https://doi.org/10.7860/JCDR/2013/6339.3282>
 32. Lowy, F. D. (2003). Antimicrobial resistance: The example of *Staphylococcus aureus*. In *Journal of Clinical Investigation* (Vol. 111, Issue 9). <https://doi.org/10.1172/JCI18535>
 33. Martin, K. W., & Ernst, E. (2003). Herbal medicines for the treatment of bacterial infections: A review of controlled clinical trials. In *Journal of Antimicrobial Chemotherapy* (Vol. 51, Issue 2). <https://doi.org/10.1093/jac/dkg087>
 34. Mordeniz, C. (2019). Introductory Chapter: Traditional and Complementary Medicine. In *Traditional and Complementary Medicine*. <https://doi.org/10.5772/intechopen.86373>
 35. Moreno, S., Scheyer, T., Romano, C. S., & Vojnov, A. A. (2006). Antioxidant and antimicrobial activities of rosemary extracts linked to their polyphenol composition. *Free Radical Research*, 40(2). <https://doi.org/10.1080/10715760500473834>
 36. Mujeeb, F., Bajpai, P., & Pathak, N. (2014). Phytochemical evaluation, antimicrobial activity, and determination of bioactive components from leaves of *Aegle marmelos*. *BioMed Research International*, 2014. <https://doi.org/10.1155/2014/497606>
 37. Muthu, C., Ayyanar, M., Raja, N., & Ignacimuthu, S. (2006). Medicinal plants used by traditional healers in Kancheepuram District of Tamil Nadu, India. *Journal of Ethnobiology and Ethnomedicine*, 2. <https://doi.org/10.1186/1746-4269-2-43>
 38. Nalule, A. S., Karue, C. N., & Katunguka-Rwakishaya, E. (2011). Anthelmintic activity of *Phytolacca dodecandria* and *Vernonia amygdalina* leaf extracts in naturally infected miniature East African goats. *Livestock Research for Rural Development*, 23(12).
 39. Nawab, A., Li, G., An, L., Nawab, Y., Zhao, Y., Xiao, M., Tang, S., & Sun, C. (2020). The potential effect of dietary tannins on enteric methane emission and ruminant production, as an alternative to antibiotic feed additives- A review. *Annals of Animal Science*, 20(2). <https://doi.org/10.2478/aoas-2020-0005>
 40. Ndaeyo, N. U. (2007). Assessing the contributions of homestead farming to food security in a developing economy: a case study of southeastern Nigeria. *Journal of Agriculture and Social Sciences*, 3(1).
 41. Nenaah, G. (2013). Antimicrobial activity of *Calotropis Procera* Ait. (Asclepiadaceae) and isolation of four flavonoid glycosides as the active constituents. *World Journal of Microbiology and Biotechnology*, 29(7). <https://doi.org/10.1007/s11274-013-1288-2>
 42. Oboh, G. (2006). Nutritive value and hemolytic properties (in vitro) of the leaves of *Vernonia amygdalina* on human erythrocyte. *Nutrition and Health*, 18(2). <https://doi.org/10.1177/026010600601800207>
 43. Okwuzu, J. O., Odeiga, P., Adetoro-tubanojo, O., & Ezechi, O. C. (2017). Cytotoxicity testing of aqueous extract of bitter leaf (*Vernonia amygdalina* Del) and sniper 1000EC (2,3 dichlorovinyl dimethyl phosphate) using the *Alium cepa* test. *African Health Sciences*, 17(1). <https://doi.org/10.4314/ahs.v17i1.19>
 44. Oloruntobi, I., Ajayi, O., & Rufus, I. (2014). Anxiolytic, Sedative and Hypothermic Effects of Aqueous Leaf Extract of *Vernonia amygdalina* Del. (Asteraceae) in Albino Mice. *British Journal of Pharmaceutical Research*, 4(18). <https://doi.org/10.9734/bjpr/2014/12529>
 45. Onabanjo, O. O., & Oguntola, C. R. B. (2003). Iron, zinc, copper, and phytate content of standardized Nigerian dishes. *Journal of Food Composition and Analysis*, 16(6). [https://doi.org/10.1016/S0889-1575\(03\)00063-2](https://doi.org/10.1016/S0889-1575(03)00063-2)
 46. Owen, O. J., Amakiri, A. O., & Karibi-Botoye, T. A. (2011). Sugar-lowering effects of bitter leaf (*Vernonia amygdalina*) in experimental broiler finisher chickens. *Asian Journal of Pharmaceutical and Clinical Research*, 4(SUPPL. 1).
 47. Oyeyemi, I. T., Akinlabi, A. A., Adewumi, A., Aleshinloye, A. O., & Oyeyemi, O. T. (2018). *Vernonia amygdalina*: A folkloric herb with anthelmintic properties. *Beni-Suef University Journal of Basic and Applied Sciences*, 7(1), 43–49. <https://doi.org/10.1016/j.bjbas.2017.07.007>
 48. Salawu, S. O., Ogundare, A. O., Ola-Salawu, B. B., & Akindahunsi, A. A. (2011). Antimicrobial activities of phenolic-containing extracts of some tropical vegetables. *African Journal of Pharmacy and Pharmacology*, 5(4). <https://doi.org/10.5897/AJPP10.317>

49. Santos, S. A. O., Martins, C., Pereira, C., Silvestre, A. J. D., & Rocha, S. M. (2019). Current challenges and perspectives for using aqueous plant extracts in managing bacterial infections: The case-study of *Salmonella enterica* serovars. In *International Journal of Molecular Sciences* (Vol. 20, Issue 4). <https://doi.org/10.3390/ijms20040940>
50. Septama, A. W., Simbak, N., & Rahmi, E. P. (2020). Prospect of plant-based flavonoids to overcome antibacterial resistance: A mini-review. In *Walailak Journal of Science and Technology* (Vol. 17, Issue 5). <https://doi.org/10.48048/wjst.2020.5583>
51. Setiawan, L. T. K., Nugraha, J., Tamayanti, W. D., & Utomo, D. H. (2018). The computational study reveals the immunomodulatory and antimicrobial effects of *vernonia amygdalina* extract. *Asian Journal of Microbiology, Biotechnology, and Environmental Sciences*, 20(December).
52. Shin, J., Prabhakaran, V. S., & Kim, K. sun. (2018). The multi-faceted potential of plant-derived metabolites as antimicrobial agents against multidrug-resistant pathogens. In *Microbial Pathogenesis* (Vol. 116). <https://doi.org/10.1016/j.micpath.2018.01.043>
53. Shokunbi, O. S., Anionwu, O. A., Sonuga, O. S., & Tayo, G. O. (2011). Effect of post-harvest processing on the nutrient and anti-nutrient compositions of *Vernonia amygdalina* leaf. *African Journal of Biotechnology*, 10(53). <https://doi.org/10.5897/ajb11.1532>
54. Singh, A., Gautam, P. K., Verma, A., Singh, V., Shivapriya, P. M., Shivalkar, S., Sahoo, A. K., & Samanta, S. K. (2020). Green synthesis of metallic nanoparticles as effective alternatives to treat antibiotics resistant bacterial infections: A review. *Biotechnology Reports*, 25. <https://doi.org/10.1016/j.btre.2020.e00427>
55. Srivastava, J., Chandra, H., Nautiyal, A. R., & Kalra, S. J. S. (2014). Antimicrobial resistance (AMR) and plant-derived antimicrobials (PDAs) as an alternative drug line to control infections. In *3 Biotech* (Vol. 4, Issue 5). <https://doi.org/10.1007/s13205-013-0180-y>
56. Steenkamp, V., Mathivha, E., Gouws, M. C., & van Rensburg, C. E. J. (2004). Studies on antibacterial, antioxidant, and fibroblast growth stimulation of wound healing remedies from South Africa. *Journal of Ethnopharmacology*, 95(2–3). <https://doi.org/10.1016/j.jep.2004.08.020>
57. Taneja, N., & Sharma, M. (2019). Antimicrobial resistance in the environment: The Indian scenario. In *Indian Journal of Medical Research* (Vol. 149, Issue 2). https://doi.org/10.4103/ijmr.IJMR_331_18
58. Tonukari, N. J., Avwioroko, O. J., Ezedom, T., & Anigboro, A. A. (2015). Effect of Preservation on Two Different Varieties of *Vernonia amygdalina*; Del. (Bitter) Leaves. *Food and Nutrition Sciences*, 06(07). <https://doi.org/10.4236/fns.2015.67067>
59. Tripathi, Y. C., & Pandey, A. K. (2017). Bioprospecting of Phylodiversity for New Therapeutic Products: Trends, Potential, and Challenges. *Organic and Medicinal Chemistry International Journal*, 2(1).
60. Uchechukwu Anastasia, U. N. (2011). The body calming and body vitalizing effects of *Vernonia amygdalina* leaf extract and leaf powder and their mechanisms. *American Journal of Agricultural and Biological Science*, 6(3). <https://doi.org/10.3844/ajabssp.2011.429.432>
61. Umaru, I. J., Badruddin, F. A., & Umaru, H. A. (2019). Phytochemical Screening of Essential Oils and Antibacterial Activity and Antioxidant Properties of *Barringtonia Asiatica* (L) Leaf Extract. *Biochemistry Research International*, 2019. <https://doi.org/10.1155/2019/7143989>
62. Vidovic, N., & Vidovic, S. (2020). Antimicrobial resistance and food animals: Influence of livestock environment on the emergence and dissemination of antimicrobial resistance. In *Antibiotics* (Vol. 9, Issue 2). <https://doi.org/10.3390/antibiotics9020052>
63. Wang, Y. F., Que, H. F., Wang, Y. J., & Cui, X. J. (2014). Chinese herbal medicines for treating skin and soft-tissue infections. In *Cochrane Database of Systematic Reviews* (Vol. 2014, Issue 7). <https://doi.org/10.1002/14651858.CD010619.pub2>
64. WHO. (2014). Antimicrobial resistance. Global report on surveillance. World Health Organization, 61(3). <https://doi.org/10.1007/s13312-014-0374-3>
65. Yang, S. K., Low, L. Y., Yap, P. S. X., Yusoff, K., Mai, C. W., Lai, K. S., & Lim, S. H. E. (2018). Plant-derived antimicrobials: Insights into mitigation of antimicrobial resistance. In *Records of Natural Products* (Vol. 12, Issue 4). <https://doi.org/10.25135/rnp.41.17.09.058>
66. Yedjou, C. G., Tchounwou, P. B., Miele, L., Ogungbe, I. v., Brown, R. J., & Payton, M. (2015). Abstract 5313: The anticancer effect of *Vernonia amygdalina* in the MMTV-PyVT transgenic mouse model. <https://doi.org/10.1158/1538-7445.am2015-5313>.