



Journal Homepage: -www.journalijar.com

INTERNATIONAL JOURNAL OF ADVANCED RESEARCH (IJAR)

Article DOI:10.21474/IJAR01/14400

DOI URL: <http://dx.doi.org/10.21474/IJAR01/14400>



RESEARCH ARTICLE

DOPPLER EFFECT ON THE WAVE ASSOCIATED WITH A PARTICLE

Alfredo Olmos Hernández and Reyna Romyna Olmos Hernández

College of Bachelors of the State of Hidalgo.

Manuscript Info

Manuscript History

Received: 10 January 2022

Final Accepted: 15 February 2022

Published: March 2022

Key words:-

Duality, Corpuscle-Wave, Doppler Effect, Wave

Abstract

It is known that a particle presents the duality corpuscle-wave. In such a way that a particle possesses an associated electromagnetic wave, the Doppler Effect appears in all the waves; in such a way that it also affects the wave associated with the particle. In this article we study the Doppler Effect on the wave associated with the particle.

Copy Right, IJAR, 2022,. All rights reserved.

Introduction:-

Wave-particle duality is a quantum phenomenon, by which particles exhibit wave behavior.

This is an experimentally proven fact on multiple occasions. It was introduced by Louis-Victor de Broglie, a French physicist of the early twentieth century. In 1924, in his doctoral thesis, inspired by experiments on electron diffraction, he proposed the existence of matter waves, meaning that all matter had a wave associated with it. This revolutionary idea, based on the analogy that the radiation had an associated particle, property already demonstrated then, did not arouse great interest, despite the correctness of its approaches, since it had no evidence of occurrence. However, Einstein recognized its importance and five years later, in 1929, De Broglie received the Nobel Prize in Physics for his work [1, 2, 5, 7 and 10]

The wavelength λ , of the wave associated with a particle is.

$$\lambda = \frac{h}{p}$$

Where h is the Planck constant and p the linear momentum of the particle

Where $p = mv\gamma$, with m the mass of the particle, v its velocity and γ the Lorentz factor.

The relativistic Doppler Effect is the observed change in the frequency of light from a source moving relative to the observer.

The change in frequency observed when the source moves away is given by the following expression [3, 4, 6, 8 and 9]

$$f_o = f_s \sqrt{\frac{1-v/c}{1+v/c}}$$

Where

Corresponding Author:- Alfredo Olmos Hernández

Address: alfredooh16@gmail.com College of Bachelors of the State of Hidalgo. Pachuca, Hidalgo, Mexico.

f_o is frequency observed.

f_s is frequency issued.

v is relative speed, positive when the emitter and the observer move away from each other.

C is the speed of light

Doppler effect present in the wave associated with a particle.

To obtain the wavelength of a given electromagnetic wave, its frequency f is.

$$\lambda = \frac{c}{f} \quad (3)$$

Substituting 3) in 2)

$$\frac{c}{\lambda_o} = \frac{c}{\lambda_s} \sqrt{\frac{1-v/c}{1+v/c}} \quad (4)$$

$$\lambda_o = \lambda_s \sqrt{\frac{1+v/c}{1-v/c}} \quad (5)$$

Substituting 1) in 5)

$$\frac{h}{p_o} = \frac{h}{p_s} \sqrt{\frac{1+v/c}{1-v/c}} \quad (6)$$

$$p_o = p_s \sqrt{\frac{1-v/c}{1+v/c}} \quad (7)$$

Where

p_o is the amount of movement measured by the observer.

p_s is the amount of movement measured in the particle.

Relative motion

By the expression 7) we have that the amount of movement that is measured of a particle, depends on the observer.

There are 2 cases:

1. Case 1

The particle moves away from the observer.

In this case the relative speed is positive, then.

$$p_o = p_s \sqrt{\frac{1-v/c}{1+v/c}} \quad (8)$$

$$p_o < p_s$$

2. Case 2

The particle approaches the observer.

In this case, the relative velocity is negative, then.

$$p_o = p_s \sqrt{\frac{1+v/c}{1-v/c}} \quad (9)$$

$$p_o > p_s$$

Conclusions:-

In this article, we obtained the expression that allows us to observe the variation in the measurement of the momentum of a particle, depending on its relative movement with respect to the observer. In such a way that if the particle approaches, the observer measures a quantity of movement greater than the measure in the particle; and if

the particle moves away from the observer, the amount of movement measured by the observer is less than if measured directly in the particle.

The expression is obtained from the study of the Doppler Effect to the wave associated with a particle.

References:-

- [1] Angarita Posada, D.S., & Pino Sánchez, J.D. (2016). About the Wave-Particle duality, a conceptual re-signification from the approaches of Louis de Broglie.
- [2] Baixauli, J.G. (2017). Wave-Particle Duality as a Classical Phenomenon.
- [3] Giraldo-Tobón, E. (2014). Wave Physics Lesson# 7c Doppler effect.
- [4] Giraldo-Tobón, E. (2014). Wave Physics Lesson# 10d Electromagnetic Waves.
- [5] Gomez, J.G. (1976). WAVE-PARTICLE EQUATION. Colombian Journal of Chemistry, 6(1), 27-32.
- [6] Londoño Rodríguez, Y. J. (2016). Of the Doppler effect and its implications, a reflection with pedagogical intentionality
- [7] Mayorga, A. (2001). Einstein, De Broglie, Schridinger (1923-1925): the wave-particle duality and the birth of wave mechanics. Technology on the Move, 14(1), 69-89.
- [8] Roatta, A., &Wolti, R. (2009). Doppler Effect for pulses and its representation in the (x, t) plane. Brazilian Journal of Physics Teaching, 31(1), 1304-1.
- [9] Sears, F. W., Zemansky, M. W., Young, H. D., Vara, R. H., García, M. G., Gümes, E. R.,&Benites, F. G. (1986). University Physics (No. 530.076 530.076 S4F5 1986 S43F5 1986 QC23 S45 1986). Inter-American Educational Fund.
- [10] Segura, A., Nieto, V., & Segura, E. (2012). An in-depth analysis of the wave-particle duality phenomenon for understanding the quantum world. Latin-American Journal of Physics Education, 6(1).