

RESEARCH ARTICLE

ARMOUR IS HEAVY, YET IT IS A PROUD BURDEN- CAP SPLINT: A CASE SERIES

Dr. Nidha Farooq, Dr. Shajaratul Yakeen Nabi, Dr. Nazia Lone and Dr. Mohsin Sidiq

Manuscript Info	Abstract
Manuscript History Received: 27 June 2023 Final Accepted: 31 July 2023 Published: August 2023	Traumatic injuries during childhood can have a considerable influence on the minds of growing children. For the purpose of supporting bony framework to pre-trauma condition, minimal manipulation of facial skeleton is necessary. In comparison to adults, the mandible in children have embedded tooth buds that hold the mandibular fragments together like a glue. Open reduction with rigid fixation is not recommended in fractures of the mandible in children. The goal of the treatment is restoration of the bony architecture in a stable position with a minimum malfunction and aesthetic disability. The purpose of the present case series is to demonstrate cap splint as a conservative and effective treatment modality for the management of pediatric mandibular fractures.

Copy Right, IJAR, 2023,. All rights reserved.

Introduction:-

Facial trauma in children can often be challenging to manage with long-term consequences involved and the psychological impact.¹ Maxillofacial fractures in pediatric population consist of less than 15% of all facial fractures with Mandibular fracture being the most common.² The incidence of mandibular fracture in children ranges from 0.6% to 1.2% with the most typical causes of fracture in children being falls (64%), traffic incidents (22%), and sports-related accidents (9%).³ The incidence of facial fractures is higher in boys than in girls with a male preponderance ranges from 1.1:1 to 8.5:1.Closed reduction techniques with maxillomandibular fixation in very young children can pose several concerns, including cooperation, compliance and adequate nutritional intake.⁴

The displaced mandibular fractures are managed by cap splints with circum-mandibular wiring, cap splint cemented onto the arch, and Erich arch-bar fixation.⁵Cap splint is a simple, easy to fabricate and more reliable method. It has other advantages like ease of application and removal, less time consumption, cost-effectiveness, good stability during healing period and minimal trauma to surrounding tissues.³

Case 1:

A7-year-old boy reported to the Department of Pediatric Dentistry with the chief complaint of pain in lower jaw with associated tenderness in chin region and difficulty in opening mouth. Patient had a history of fall from the height 1 day back. There was no history of loss of consciousness or vomiting. Intraoral examination revealed tenderness along the lower border of mandible on right side of canine region. Preoperative OPG showed right mandibular parasymphyseal fracture between 41 and 82. The primary impressions were taken with alginate and two sets of casts were subsequently poured. Mock Surgery and Articulation of casts was done and a modified closed acrylic cap splint was fabricated. Displaced Fractured fragments were reduced under local anaesthesia and a closed cap splint was cemented to the reduced fractured jaw with help of luting GIC. Oral hygiene instructions were given

to the patient and analgesics were prescribed. After 3 weeks the splint was removed and the patient was asymptomatic. At follow-up, an OPG revealed healing of the fracture site after 6 months of follow up.



Preoperative Intraoral photograph



Preoperative OPG showing right mandibular parasymphysis fracture



Closed cap splint cemented with luting GIC



Post-operative Intraoral photograph



Post-operative OPG at 6 months follow up.

Case 2:

A 2-year-old girl reported to the Department of Pediatric Dentistry with the chief complaint of Trauma due to fall 1 day back. There was no history of loss of consciousness or vomiting. There was associated soft tissue laceration on lower lip and chin with Positive Coleman's Sign. There was a step deformity in mandible with associated fracture of right mandibular parasymphyseal region. The primary impressions were taken with alginate and two sets of casts were subsequently poured. Mock Surgery and Articulation of casts was done and a modified closed acrylic cap splint was fabricated. Displaced Fractured fragments were reduced under local anaesthesia and closed cap splint was cemented to the reduced fractured jaw with help of luting GIC. Oral hygiene instructions were given to the patient and analgesics were prescribed. After 3 weeks the splint was removed and the patient was asymptomatic. At follow up there was correction of step deformity.



Preoperative Photograph



Closed cap splint cemented with luting GIC



Post- operative Photograph

Case 3:

A 3-year-old girl reported to the Department of Pediatric Dentistry with the chief complaint of pain in lower jaw with associated tenderness in chin region and difficulty in opening mouth. Patient had a history of fall on the same day. There was no history of loss of consciousness or vomiting. Intraoral examination revealed tenderness along the lower border of mandible on right side of canine region. The primary impressions were taken with alginate and two sets of casts were subsequently poured. Mock Surgery and Articulation of casts was done and a modified closed acrylic cap splint was fabricated. Displaced Fractured fragments were reduced under local anaesthesia and a closed cap splint was cemented to the reduced fractured jaw with help of luting GIC. Oral hygiene instructions were given to the patient and analgesics were prescribed. After 3 weeks the splint was removed and the patient was asymptomatic.



Closed cap splint cemented with luting GIC



Preoperative Photograph



Post- operative Photograph

Discussion:-

In Pediatric age group, Facial fractures account for about 5% of all facial fractures with the most common cause being fall from height, road traffic accident, sports injuries, etc.⁶The high tooth to bone ratio predisposes the mandible to fracture compared to the midface.⁷Mandibular fracture sites include Condyle (55%), Parasymphysis (27%), Body (9%), and Angle (8%).⁸ The various symptoms of mandibular fractures include Pain, Swelling, Trismus, Occlusal derangement, Sublingual Hematoma, Step- deformity, Deviation, Loss of sensation, Open bite.⁹Pediatric patients presents a unique challenge to oral surgeons because of anatomic variation and changes in the mandibular growth.⁶Therefore, the management of the pediatric patients with maxillofacial injury should take into consideration the differences in anatomy and physiology, particular stage in growth and development, degree of compliance, complexity and any concomitant injury, anatomic sites injured, and the time elapsed since injury.⁹

The primary goal in the treatment of mandibular fractures is Reestablishment of Occlusion.Various techniques utilized in management of paediatric fractures include Tape muzzles, Circumferential wiring, Acrylic splint, Percutaneous skeletal fixation, Open reduction, Resorbable plates, Modified orthodontic brackets, Rubber elastics in combination with orthodontics brackets, Nickel titanium staples. ⁶In the present case series, cap splint has been used for the stabilization of mandibular fractures as Cap splint is a promising fixation technique and provides for Occlusion guided fracture reduction, Maximum stability during healing period, Ease of application and removal, Reduced operation time, Minimal trauma for adjacent anatomic structures, Ease of maintenance of oral hygiene, cost-effectiveness and Comfort for young patients. ⁵Therefore, the closed cap splint should be a reliable method in minimally displaced mandibular fractures in pediatric patients.

Conclusion:-

The present case series shows that the conservative management of fractured pediatric mandible is cost-effective, safe and a minimally invasive procedure with good prognosis and almost complication free outcome and the role of a pediatric dentist is vital in managing children with such fractures.

References:-

- 1. Rogan DT, Fang A. Pediatric Facial Trauma. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2023. cited 2023 Aug 31.
- 2. Sharma A, Patidar DC, Gandhi G, Soodan KS, Patidar D. Mandibular Fracture in Children: A New Approach for Management and Review of Literature. Int J Clin Pediatr Dent. 2019;12(4):356–9.
- 3. Abbood AA. Treatment of Pediatric Mandibular Fractures Utilizing the Open Cap Splint: A Clinical Study. J Res Med Dent Sci. 2021;9(2).
- 4. Sezer B, Koyuncu B, Akay C, et al. Splint fixation with circummandibular wires for pediatric mandibular fracture: Two case reports. J Ege University School Dent 2010; 3:113-117 -
- 5. Nezam S, Kumar A, Shukla JN, Khan SA. Management of mandibular fracture in pediatric patient. Natl J Maxillofac Surg. 2018;9(1):106–9.
- 6. Kumar N, Richa, Gauba K. Modified closed cap splint: Conservative method for minimally displaced pediatric mandibular fracture. Saudi Dent J. 2018 Jan;30(1):85-88.
- 7. Singh MM, Rajpal S, Priya N, Ali MG, Akhtar S. CAP splint: An armour to safeguard developing dentition in paediatric mandibular fractures- A case series. IP Indian J Orthod Dentofac Res. 2021 Apr 15;7(1):77–81.
- 8. Sharma S, Vashistha A, Chugh A, Kumar D, Bihani U, Trehan M, et al. Pediatric Mandibular Fractures: A Review. Int J Clin Pediatr Dent. 2009;2(2):1–5.
- 9. Singh J, Khadka R, Chaturvedi PC, Singh V, Rai P. Circummandibular wiring made easy: A case report. Rev Esp Cir Oral Maxilofac. 2014 Oct 1;36(4):191–5.