



Journal Homepage: -www.journalijar.com

INTERNATIONAL JOURNAL OF ADVANCED RESEARCH (IJAR)

Article DOI:10.21474/IJAR01/18525
DOI URL: <http://dx.doi.org/10.21474/IJAR01/18525>



RESEARCH ARTICLE

ECCRINESQUAMOussyRINGOMETAPLASIA IN A PATIENT WITH ROSACEA

Ghita Sqalli Houssini¹, Sara Elloudi¹, Zakia Douhi¹, Meryem Soughi¹, Hanane Baybay¹, Mounia Rimani² and Fatima Zahara Mernissi¹

1. Department of Dermatology, University Hospital Hassan II, Faculty of Medicine and Pharmacy, Sidi Mohamed Ben Abdellah University, Fes, Morocco.
2. Hassan Anatomy and Pathology Laboratory, Rabat, Morocco.

Manuscript Info

Manuscript History

Received: 10 February 2024
Final Accepted: 14 March 2024
Published: April 2024

Key words:-

Eccrine, Metaplasia, Rosacea, Syringometaplasia

Abstract

Background: Eccrine squamous syringometaplasia is a benign adaptive metaplasia that occurs in response to various physiological and pathological stimuli. It is associated with chemotherapy or other dermatological disorders, particularly inflammatory conditions. We present a case of an inflammatory rosacea associated with squamous metaplasia in a female patient.

Observation: A 68-year-old female presented with a chronic skin eruption consisting of translucent and angioma-like papules, accompanied by a flushing sensation on the face. Dermoscopy revealed milium grains and an angioma-like background. Biopsy confirmed the diagnosis of erythematous papular rosacea associated with malpighian squamous syringometaplasia.

Conclusion: Squamous syringometaplasia can be linked to inflammatory dermatoses, such as rosacea. The inflammatory infiltrate is implicated in eccrine metaplasia, which, in turn, may contribute to an uncommon presentation of rosacea.

Copy Right, IJAR, 2024., All rights reserved.

Introduction:-

Syringometaplasia can be defined as a reactive process occurring around eccrine glands in response to various physiological or pathological stimuli, leading to a benign and adaptive metaplasia of the duct and eccrine gland [1]. There are three types of syringometaplasia: squamous, mucinous, and adenoid. Squamous syringometaplasia is the most common subtype, classified into two types: the first associated with chemotherapy and the second linked to dermatological conditions, including infectious, neoplastic, or inflammatory disorders [2,3]. In this report, we present the case of a patient who presented with a chronic papular facial eruption, accompanied by a sensation of flushing and burning. Histological examination confirmed the diagnosis of papulopustular rosacea associated with syringometaplasia.

Observation:-

A 68-year-old female, under observation for an unspecified rheumatic condition, presented during the summer seeking management for a facial eruption that had been progressing over the past 20 years. She complained of a burning sensation, accompanied by flushing, particularly during sun exposure, exposure to heat, and the consumption of

Corresponding Author:- Ghita Sqalli Houssini

Address:- Department of Dermatology, University Hospital Hassan II, Faculty of Medicine and Pharmacy, Sidi Mohamed Ben Abdellah University, Fes, Morocco.

spicy foods. The patient reported a pattern of episodic exacerbation during the summer months and remission in the winter. No history of tuberculosis or systemic symptoms was reported.

Dermatological examination revealed the presence of multiple firm, translucent papules in some areas and erythematous-violet papules in others. These lesions were situated on an erythematous background, predominantly on the cheeks, chin, and forehead, forming a hemifacial distribution that extended to the neck and scalp (Fig. 1a, b). Dermoscopy identified several milium grains with an angioma-like appearance, with suspected lipid features on the cheeks (Fig. 2a, b, c).

Considering this clinical presentation, potential diagnoses included granulomatous rosacea, small-nodule sarcoidosis, crystalline miliaria, and lupus miliaris tuberculosus. An eccrine gland anomaly, specifically eccrine syringometaplasia, was also considered due to the predominance of the eruption on the face, the translucent nature of the papules, and the aggravation with heat.

A 5mm punch biopsy was performed, revealing erythematopapular rosacea associated with milium grains and focal malpighian syringometaplasia (Fig. 3a, b, c). Post-biopsy, the patient was educated on hygienic-dietary measures. She was prescribed an anti-redness cream and a three-month course of tetracycline, resulting in slight improvement, albeit hindered by the patient's poor adherence to the therapeutic regimen.

Discussion:-

Squamoussyringometaplasia is a metaplastic process that replaces the normal cuboidal epithelium of eccrine sweat glands with a keratinizing malpighian epithelium similar to the epidermal spinous layer [4]. Squamoussyringometaplasia is categorized into two types. The first type is associated with chemotherapy drugs such as carmustine, cytarabine, cyclophosphamide, daunorubicin, cisplatin, 5-fluorouracil, doxorubicin, etoposide, methotrexate, and others [2,5]. Clinically, in such cases, it presents as a skin eruption consisting of macules, papules, and vesicles that appear between 2 to 39 days after chemotherapy administration. The second type has been observed in various dermatological conditions, including neoplastic pathologies like keratoacanthoma and Squamous Cell Carcinoma [2], mycosis fungoides [6], infections by cytomegalovirus and herpes, especially in HIV patients [7,8], as well as inflammatory conditions, such as panniculitis, scleroderma, and lupus.

In the latter cases, the inflammatory infiltrate (granulomatous, neutrophilic, or lymphocytic) leads to gland and eccrine duct necrosis, resulting in the regeneration of the epithelium in a keratinizing form. It is important to note that this type of syringometaplasia usually does not cause a significant change in the clinical features of the associated condition. It is often discovered incidentally through histopathological examination [1].

Squamoussyringometaplasia associated with papular erythematous rosacea has not been previously documented. The clinical presentation of our patient is notably atypical compared to the classic inflammatory rosacea. This prompts us to consider whether the metaplastic process within the eccrine glands themselves could be the root cause of the distinctive clinical and dermoscopic features observed in our patient. Further investigation and exploration of this unique case may provide valuable insights into the complex interplay between metaplastic processes and dermatological conditions.

Conclusion:-

For the first time, rosacea is described in association with squamoussyringometaplasia. The inflammation induced by rosacea appears to be a key factor in the metaplastic changes occurring in the eccrine glands and ducts. However, intriguingly, this metaplasia may, in turn, contribute to the clinical presentation of rosacea, characterized by a papular, angioma-like, and translucent appearance. This novel observation underscores the intricate relationship between inflammatory processes and metaplastic changes within eccrine structures, providing a unique perspective on the varied manifestations of dermatological conditions. Further research and exploration are warranted to elucidate the underlying mechanisms and potential therapeutic implications of this distinctive association.

Consent

The examination of the patient was conducted according to the Declaration of Helsinki principles.



Fig1.a:-



Fig1.b:-




Fig2.a:-




Fig2.b:-




Fig2.c:-




Fig3.a:-




Fig3.b:-




Fig3.c:-

References:-

- [1] Abbas O, Bhawan J. Syringometaplasia: variants and underlying mechanisms. *Int J Dermatol.* 2016;55(2):142-8.
- [2] Serrano T, Saez A, Moreno A. Eccrine squamous syringometaplasia. A prospective clinicopathologic study. *J Cutan Pathol.* 1993; 20: 61–65.
- [3] Grynspan D, Meir K, Senger C, et al. Cutaneous changes in fibrous hamartoma of infancy. *J Cutan Pathol.* 2007; 34: 39–43.

- [4] García-Río I, Pérez-Gala S, Fraga J, García-Díez A, Fernández-Herrera J. Eccrine squamous syringometaplasia in a patient with systemic lupus erythematosus. *J Eur Acad Dermatol Venereol.* 2006;20(5):608-10.
- [5] King DT, Barr RJ. Syringometaplasia: mucinous and squamous variants. *J Cutan Pathol.* 1979; 6: 284–291.
- [6] Pileri A, Facchetti F, Rütten A, Zumiani G, Boi S, Fink-Puches R et al C. Syringotropic mycosis fungoides: a rare variant of the disease with peculiar clinicopathologic features. *Am J Surg Pathol.* 2011;35(1):100-9.
- [7] Muñoz E, Valks R, Fernández-Herrera J, Fraga J. Herpetic syringitis associated with eccrine squamous syringometaplasia in HIV-positive patients. *J Cutan Pathol.* 1997;24(7):425-8.
- [8] Dauden E, Porras JI, Buezo GF, et al. Eccrine squamous syringometaplasia and cytomegalovirus. *Am J Dermatopathol.* 2000; 22: 559–561

Figure's Legends:

Figure 1: Erythematous papular eruption with angioma-like features on the face, displaying a hemifacial pattern on the forehead (a, b).

Figure 2: Dermoscopy reveals angioma-like features (a), milium grains (b), and suspicion of a lupoid aspect (c).

Figure 3: Histopathological examination at 100x magnification demonstrates telangiectasias, perivascular, and perifollicular lymphoplasmacytic inflammatory infiltrates, along with milium grains (a). Squamous metaplasia of a sweat duct near dilated sweat glands at 100x magnification (b). Detailed view of the duct with malpighian metaplasia of the lining, highlighted by an eosinophilic cuticle at 400x magnification (c).