



Journal Homepage: - www.journalijar.com

INTERNATIONAL JOURNAL OF ADVANCED RESEARCH (IJAR)

Article DOI: 10.21474/IJAR01/20393

DOI URL: <http://dx.doi.org/10.21474/IJAR01/20393>



RESEARCH ARTICLE

EFFECT OF COMPREHENSIVE NURSING INTERVENTION ON SELF-CARE PRACTICES OF PATIENTS UNDERGOING HEMODIALYSIS

Minu Pargaie¹, Kamli Prakash² and Sanchita Pugazhendi³

1. Research Scholar, Department of Nursing, Swami Rama Himalayan University, Jolly Grant, Dehradun.
2. Professor, Department of Nursing, Swami Rama Himalayan University, Jolly Grant, Dehradun.
3. Professor, Department of Nursing, Swami Rama Himalayan University, Jolly Grant, Dehradun.

Manuscript Info

Manuscript History

Received: 08 December 2024

Final Accepted: 12 January 2025

Published: February 2025

Key words:-

Comprehensive Nursing Intervention, Hemodialysis, Self-Care Practice, Fistula Care, Physical Activity, Dietary Modification, Fluid Restriction

Abstract

Background: Chronic kidney disease is a prolonged medical condition that impacts the self-care routines of patients receiving hemodialysis. Self-care practices significantly influence the treatment outcomes of patients, and it is essential for nurses to provide appropriate guidance to address the challenges that may arise during hemodialysis sessions.

Objectives: This study aimed to invent comprehensive nursing intervention and evaluate its efficacy on self-care practices.

Design: This study was experimental with a time series design.

Participants: Total enumerative sampling method was used to recruit 106 patients newly recruited for hemodialysis. The participants were randomly assigned to one of two groups: the intervention group given comprehensive nursing intervention, while the control group received routine care.

Measurements: This study was experimental with a time series design. The measurement instrument included structured self-care practice checklist which involves questions related to fistula care, physical activity, dietary modification and fluid restriction. The analysis involves use of chi square and independent t test.

Results: The participants mean age was 47.5 years. Application of comprehensive nursing interventions significantly ($p < 0.05$) improved self-care practice of patients in experimental group as compared to control group.

Conclusion: Comprehensive Nursing Intervention can improve self-care practice of patients undergoing hemodialysis.

Copyright, IJAR, 2025.. All rights reserved.

Introduction:-

Chronic Kidney Disease (CKD) is characterized by a gradual decline in kidney function, which may eventually necessitate renal replacement therapies, including dialysis or transplantation. (Vaidya SR, Aeddula NR. 2024). Rapid urbanization, which leads to increased obesity and reduced physical activity, is contributing to shifts in the incidence of risk factors associated with chronic kidney disease (CKD) on a global scale. (Neuen LB 2017). Hemodialysis is the predominant treatment option, with renal transplantation following closely behind, while peritoneal dialysis ranks significantly lower. Similar to other developing nations, India faces distinct circumstances and challenges that

Corresponding Author:- Minu Pargaie

Address:- Research Scholar, Department of Nursing, Swami Rama Himalayan University, Jolly Grant, Dehradun.

affect the management of chronic kidney disease. An overview of preventive strategies for chronic kidney disease (CKD) presented by Li PKT et al. (2020) highlights the importance of prevention at three distinct levels. Primary prevention targets the identification of individuals at high risk, the management of obesity, the enhancement of glycaemic control, and the regulation of blood pressure. It also advocates for the reduction of salt intake and the promotion of healthy dietary and lifestyle choices. Secondary prevention focuses on the early detection of CKD, the management of blood pressure, and the reduction of protein consumption. Tertiary prevention encompasses fluid management, nutritional therapy, and the use of pharmacological treatments. The availability of facilities and expertise varies considerably across different regions of the country. (Varughese S, Abraham G. 2018). An integrative review concerning patients undergoing hemodialysis has highlighted the importance of self-care management factors, emphasizing that the patients themselves play a crucial role in formulating effective strategies aimed at improving both physical and psychological outcomes.

A concept paper highlighted the importance of self-care practices among hemodialysis patients, focusing on the critical aspects of vascular access maintenance, fluid management, dietary compliance, and adherence to treatment during the hemodialysis process. (Pargaian M, Prakash DK. 2025).

A pilot study indicated that a dietary intake monitoring application assisted patients in adhering to their prescribed diets and enabled them to review their previous meals through a reflective mechanism. (Connelly K et.al 2012).

Patients undergoing hemodialysis often lack sufficient self-care, which negatively impacts their self-care efficacy and, consequently, their overall quality of life. (Rayyani M, et.al. 2014). Therefore, this study aimed to develop a comprehensive nursing intervention for patients receiving hemodialysis and evaluate its effectiveness in improving self-care practices.

Methods:-

A quantitative methodology employing a Randomized Controlled Trial with a Time Series design was utilized to evaluate the efficacy of a comprehensive nursing intervention and the participants were chosen through the total enumerative sampling technique.

Participants And Setting

The study recruited adult patients receiving hemodialysis from dialysis centre Uttarakhand, India. New patients planned for hemodialysis during fistula formation were enrolled in the study. Exclusion criteria included: patients having serious cognitive impairment, unstable medical conditions and patients not able to perform activities of daily living.

Based on the literature previously published (Mansouri et al., 2020), a calculation for sample size was performed. The mean \pm standard deviation was extracted from the beforementioned article to attain an 80% power (B) at a significance level of 5% (α). The formula utilized for the sample size calculation is expressed as follows: $n = 2SD^2 \times (Z_{\alpha/2} + Z_{\beta})^2 / d^2$. The determined sample size amounted to 90. Taking into account a 10% dropout rate, the researcher subsequently enrolled 106 patients across both the experimental and control groups.

Intervention

Comprehensive Nursing Intervention

The intervention for this study was formulated by the researcher following an evaluation of patients' needs through the implementation of focused group discussions. The component of Comprehensive nursing intervention encompassed educating patients about the anatomy and function of the kidneys, as well as the operation and significance of hemodialysis. Instruction was provided on the hygiene practices related to the fistula, care for the fistula arm, and the essential Do's and Don'ts associated with it. Additionally, a demonstration was conducted on how to assess the functionality of the fistula. The importance of dietary management was emphasized, including guidance on food items that should be consumed or avoided by patients undergoing hemodialysis, along with a demonstration on potassium leaching. Strategies for managing thirst were also taught, alongside the significance of exercise and sleep hygiene. Furthermore, demonstrations were provided on appropriate exercises, including asanas and breathing techniques, tailored for patients receiving hemodialysis.

Outcome Measure Instrument

Self-Care Practice Checklist: To assess the self-care practices of patients undergoing haemodialysis, a self-care practice checklist was created, comprising twenty questions based on a six-point Likert scale. This checklist focused on two primary areas: fistula care and physical activity. The Likert scale included the following response options: Six to seven days a week (5), Four to five days a week (4), Two days a week (3), Three days a week (2), Once a week (1), and Never (0). The tool contained a total of twenty questions, with fourteen items framed positively and six items framed negatively, necessitating reverse scoring for the latter. Additionally, a dichotomous checklist with twenty-eight questions was utilized to evaluate dietary modifications and fluid restrictions. This dichotomous response format offered options of yes (1) and no (0), with nine items framed positively and nineteen items requiring reverse scoring. A higher score indicated better self-care practices, with the scoring range spanning from a minimum of 0 to a maximum of 148.

Study Procedure

Participant recruitment

New patients enrolled for hemodialysis having new fistula formation within three weeks of fistula formation and meeting inclusion criteria were enrolled in the study. Total 116 patients were eligible for the study eight patient did not meet inclusion criteria and two patients denied to participate in study.

Random assignment and homogeneity analysis

This study was a Randomised Controlled Trial with time series design. The participants were assigned to control and experimental groups at random through the Sequentially Numbered Opaque Sealed (SNOOSE) method. To evaluate the homogeneity of demographic and clinical characteristics between the intervention and control groups, an independent t-test and a chi-square test were employed (refer to Table 1). This approach was taken to confirm that any outcomes observed could be directly linked to the implementation of the comprehensive nursing intervention, rather than to any pre-existing disparities between the two groups.

Participant instruction

Once baseline data for the study were obtained, the intervention group received six session of comprehensive nursing intervention. Participants were encouraged to practice fistula care, use of reuseable ice cubes, potassium leaching of food items and practice of coordination/ strengthening exercise and yoga daily. Follow up of all participants were made for a period of six months.

Data collection

A researcher gathered in-person data from 106 participants at a hemodialysis centre within an Indian hospital while the patients were waiting for their hemodialysis treatment. The investigator provided a questionnaire focused on self-care practices, which the participants could complete on their own or have read to them by the researcher.

Table 1:- Baseline participant profiles between the intervention and control group.

Sociodemographic Characteristics	Experimental group (52) n (%)	Control group (54) n (%)	χ^2/ t value	p value
Age Mean \pm SD	47.5 \pm 12.86	48.5 \pm 14.60	0.67	0.50
Gender				
Male	28 (54)	32(60)	0.31	0.57
Female	24 (46)	22 (40)		
Marital status				
Married	41 (79)	44(81)	1.28	0.52
Unmarried	07 (13)	11(10)		
Divorced/ Widow/Widower	04 (8)	10(09)		
Education qualification				
No formal Education	12 (23)	07 (13)	3.78	0.70
Primary Education	06 (11)	07 (13)		
Middle School	09 (17)	07 (13)		
High School	11 (21)	12 (22)		
Intermediate	08 (15)	15 (28)		

Graduate	02 (04)	02 (04)		
Post Graduate	04(08)	04 (07)		
Residency				
Rural	35 (67)	38 (70)		
Urban	15 (29)	15 (28)	0.41	0.81
Semi urban	02 (04)	01 (02)		
Family Income per month in Rupees.				
Less than 10,000	36 (69)	33 (61)		
10,001- 20,000	03 (06)	10 (19)	8.78	0.10
20,001 – 30,000	09 (17)	04 (07)		
30,001 – 40,000	0	03 (06)		
40,001-50,000	02 (04)	02 (03)		
More than 50,000	02 (04)	02 (04)		
Employment/ job status				
Private job	05 (10)	08 (15)		
Government job	00 (00)	01 (02)	1.70	0.88
Self employed	06 (12)	06 (11)		
Homemaker	15 (29)	14 (26)		
Unemployed	24 (46)	23 (43)		
Retired	02 (04)	02 (04)		
Receiving health benefits from				
Ayushman Card	50 (96.15)	49 (90.74)	3.77	0.28
Health insurance	01 (1.92)	04 (7.41)		
Personal money	01 (1.92)	01 (1.85)		
Years since diagnosed with CKD	0.73 ± 1.32	0.70 ±1.07	0.68	0.49
Mean±SD				
Hemodialysis Session per week				
1 times	04 (7.69)	04 (7.41)		
2 times	35 (67.31)	36(66.67)	3.86	0.42
3 times	13(25.00)	14 (25.92)		
Diabetes mellitus				
Yes	18(34.62)	17(31.48)	0.11	0.73
No	34(65.38)	37(68.52)		
Hypertension				
Yes	30 (57.69)	36 (66.67)	0.90	0.34
No	22 (42.31)	18 (33.33)		
BMI				
Underweight	15 (28.85)	18 (33.33)		
Normal Weight	34 (65.38)	30 (55.56)	0.50	0.61
Overweight	03 (5.77)	06 (11.11)		

Data analysis

Demographic and clinical characteristics were evaluated through descriptive statistics. The homogeneity of the groups at the baseline of the study was examined utilizing the chi-square test, Fisher's exact test, and two-tailed independent t-tests. The impact of the intervention was subsequently assessed at 12 and 24 weeks.

Table 2:- Comparison of Self-care practice scores between experimental and control group at baseline.

Domains of self-care practice	Experimental group (52) Mean±SD	Control group (54) Mean±SD	t- test	p-value
Fistula Care	24.15 ±4.4	22.48 ± 3.4	2.23	0.025
Physical Activity	18.71 ±4.5	17.85 ± 4.8	1.35	0.175
Dietary Modification	14.73 ± 2.2	13.57 ± 2.7	1.99	0.046
Fluid restriction	3.73 ± 1.1	3.75 ± 1.3	0.08	0.930

Total Self-care Practice	61.32 ± 08.13	57.60 ± 07.47	2.41	0.017
---------------------------------	---------------	---------------	------	-------

*Independent t test, p< 0.05

Table no 3:- Changes in self-care practice of patients undergoing haemodialysis from Baseline to 12 weeks and Baseline to 24 weeks.

Variable		Experimental group (52) Mean±SD Median (min,max)	Control group (54) Mean±SD Median (min,max)	p-value (between group)
Fistula Care	Baseline to 12 weeks	9.96 ± 4.07 0 (-3 ,10)	-0.43 ± 3.60 0 (-8, 15)	< 0.001
	Baseline to 24 weeks	9.11 ± 5.24 10 (-3, 20)	-1.34 ± 9.11 -0.5 (-16, 20)	< 0.001
Activity	Baseline to 12 weeks	29.64 ± 9.46 31 (-5, 42)	-1.18 ± 6.99 -2 (-11, 44)	< 0.001
	Baseline to 24 weeks	32.07 ± 9.49 34 (-6, 46)	2.21 ± 6.67 -2.5 (-12, 38)	< 0.001
Dietary Modification	Baseline to 12 weeks	5.19 ± 3.31 5 (-2 ,16)	-2.69 ± 3.30 -3 (-10, 8)	< 0.001
	Baseline to 24 weeks	5.41 ± 2.69 5 (-1, 10)	-3.46 ± 3.93 -3.5(-14,11)	< 0.001
Fluid restriction	Baseline to 12 weeks	2.96 ± 1.21 3 (0 ,5)	-0.20 ± 1.26 0(-3 , 5)	< 0.001
	Baseline to 24 weeks	3.09 ± 1.23 3(-1, 5)	-0.48 ± 1.30 0 (- 4, 3)	< 0.001

Ethical considerations

To maintain confidentiality, participant records, which were designated by numerical identifiers, were securely stored apart from the measurement data in a locked file drawer. Electronic files were safeguarded on an encrypted computer. Access to the research information was restricted solely to the study researcher.

Results:-

Baseline personal profiles

As shown in Table 1, 106 patients receiving haemodialysis were enrolled in the study. Participants' mean age was 47.5 (SD = 12.86). They had received haemodialysis for a mean of 0.73 (SD = 1.32) months. Most participants were married, lived with family members, reported no formal education, residing in rural area, having monthly family income of less than 10,000 rupees per month and were unemployed. No significant differences in demographic and clinical characteristics were found between intervention and control groups at baseline (all p >0.05)

Changes in self-care practice

The mean baseline fistula care score was 24.15 (SD = 04.40), physical activity score was 18.71 (SD = 4.50), dietary modification score was 14.73 (SD = 2.20), fluid restriction score was 3.73 (SD = 1.10) indicating moderate self-care practice. For all the variables (Fistula Care, Activity, Dietary Modification, and Fluid Restriction), the experimental group showed significant improvements compared to the control group at both 12 and 24 weeks. The p-values indicate that these differences are statistically significant (p < **0.001**) for all measures.

Discussion:-

Study's findings align with existing literature demonstrating the positive impact of educational interventions on self-care practices among hemodialysis patients. Several studies have reported significant improvements in patients' adherence to treatment regimens, including fluid and dietary restrictions, following structured educational programs.

For instance, a study published in the International Journal of Health Sciences evaluated the effect of an educational intervention on hemodialysis patients' knowledge and adherence levels. The results indicated that the intervention group experienced a significant increase in knowledge regarding disease management, fluid adherence, and dietary

adherence compared to the control group. Notably, adherence improved across all domains, with statistically significant enhancements in fluid and dietary restrictions (Dsouza Bet al. 2023)

Similarly, research featured in the Journal of Education and Health Promotion assessed the impact of education based on the PRECEDE model on self-care behaviours in hemodialysis patients. The findings revealed that the educational intervention effectively elevated the level of self-care behaviours among participants, emphasizing the importance of tailored educational programs in promoting better health practices.(Mohammed AW .et.al. 2023)

Furthermore, a study in the Journal of Renal Care examined the effects of a hemodialysis patient education program on adherence to fluid and dietary restrictions. The study concluded that the training provided to hemodialysis patients positively influenced their compliance with diet and fluid restrictions, leading to improved adherence (Başer E, Mollaoğlu M. 2019)

These studies, among others, corroborate findings that structured educational interventions can significantly enhance self-care practices in hemodialysis patients, particularly concerning fluid restriction and dietary modifications. Implementing such programs can empower patients to manage their condition more effectively, potentially leading to improved health outcomes.

Limitations

The results of the study were influenced by the participants' adherence to the comprehensive care intervention, which encompassed multiple domains. This complexity contributed to lower adherence rates and inconsistent application of the intervention. Additionally, the participants were aware of their group assignments, which may have introduced bias into the self-reported outcomes.

Implications for clinical practice and future research

Enhancing the value of future investigations involves exploring the adherence to self-care practice. Particular emphasis should be placed on qualitative responses to contribute to the development and refinement of the intervention. Face-to-face interviews instead of written self-reports can further enrich the qualitative data. This expanded approach aims to foster a comprehensive understanding of the intervention's potential benefits.

Conclusions:-

In summary, this study provides compelling evidence to support the efficacy of comprehensive nursing intervention on self-care practice among patients receiving haemodialysis. User-friendly and cost-effective supplementary modalities

Author Contributions

Minu Pargaian: Principal project leader who conceived the study, invented comprehensive nursing intervention, recruited the participants and drafted and revised the manuscript.

Dr. Kamli Prakash: Helped to select the study, helped in formulating the intervention and guided during the study period.

Dr. Sanchita Pugazhendi: Guided during the study period. and approved the final manuscript.

Acknowledgements:-

The authors would like to thank the nurse in charge and nurses for helping them recruit the study participants at the haemodialysis departments in the Nephroplus dialysis centre.

Conflict Of Interest Statement

The authors declare no conflict of interest.

Data Availability Statement

Data sharing is not applicable to this article as no new data were created in this study.

Referances:-

1. Başer E, Mollaoğlu M. (2019) The effect of a hemodialysis patient education program on fluid control and dietary compliance. *Hemodial Int.* 23(3):392–401. Available from: <https://onlinelibrary.wiley.com/doi/10.1111/hdi.12744>
2. Connelly K, Siek KA, Chaudry B, Jones J, Astroth K, Welch JL. (2012) An offline mobile nutrition monitoring intervention for varying-literacy patients receiving hemodialysis: a pilot study examining usage and usability. *Journal of the American Medical Informatics Association.* 19(5):705–712. Available from: <https://doi.org/10.1136/amiajnl-2011-000732>
3. Dsouza B, Prabhu R, Unnikrishnan B, Ballal S, Mundkur SC, Chandra Sekaran V, et al. (2022) Effect of Educational Intervention on Knowledge and Level of Adherence among Hemodialysis Patients: A Randomized Controlled Trial. *Glob Health Epidemiol Genomics.* 2023(1), 4295613. Available from: <https://doi.org/10.1155/2023/4295613>
4. Lukman NA, Leibing A, Merry L. (2020) Self-Care Experiences of Adults with Chronic Disease in Indonesia: An Integrative Review. *Int J Chronic Dis.* 1379547. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7468663/>
5. Li PKT, Garcia-Garcia G, Lui SF, Andreoli S, Fung WWS, Hradsky A, et al (2020). Kidney Health for Everyone Everywhere – From Prevention to Detection and Equitable Access to Care. *KDD.* 6(3):136–43. Available from: <https://www.karger.com/Article/FullText/506528>
6. Mansouri S, Jalali A, Rahmati M, Salari N. (2020) Educational supportive group therapy and the quality of life of hemodialysis patients. *BioPsychoSocial Medicine.* 15;14. Available from: <https://doi.org/10.1186/s13030-020-00200-z>
7. Mohammed A W , Abdel- Mordy AM , Mohamoud AA.(2023).Effect of Educational Intervention on Self- Care Behaviours of Hemodialysis Patients: Based on PRECEDE Model. *Egyptian Journal of Health Care.* 14 (3) 264 – 284. Available from: https://ejhc.journals.ekb.eg/article_316046_3e06ad49801582b5b05c063e838d5e03.pdf?utm_source=chatgpt.com
8. Neuen BL, Chadban SJ, Demaio AR, Johnson DW, Perkovic V. (2017) Chronic kidney disease and the global NCDs agenda. *BMJ Glob Health.* 2 (2). 1-4 Available from: <https://gh.bmj.com/content/2/2/e000380>
9. Pargaian M, Prakash DK. (2025) Significance of self - care practices adopted by patients undergoing Hemodialysis: Concept Article. *Journal of Emerging Technology and Innovative Research .* 12 (1). 426-430. Available from: <https://www.jetir.org/view?paper=JETIR2501175>
10. Rayyani M, Malekyan L, Forouzi M, Razban F. (2014) Self-care Self-efficacy and Quality of Life among Patients Receiving Hemodialysis in South-East of Iran. *Asian Journal of Nursing Education and Research.* 4 (2). 165-171
11. Vaidya SR, Aeddula NR. (2024) Chronic Kidney Disease. In: *StatPearls.* Treasure Island (FL): StatPearls Publishing; Available from: <http://www.ncbi.nlm.nih.gov/books/NBK535404/>
12. Varughese S, Abraham G. (2018) Chronic Kidney Disease in India: A Clarion Call for Change. *Clinical Journal of American Society of Nephrology.* 13(5):802–804. Available from: <https://cjasn.asnjournals.org/content/13/5/802>
13. Yasin F, Khraim F, Santos M, Forgrave D, Hamad A. (2024) Factors influencing self-care management in adult hemodialysis patients: An integrative review. *Qatar Med Journal.* (1):12.1-15 Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC11037095/>