



RESEARCH ARTICLE

EFFECTIVENESS OF THE SAFE DELIVERY APP IN ENHANCING NURSING STUDENTS' KNOWLEDGE OF MATERNAL SEPSIS

Kavita Thakur¹, Gurwinder Kaur², Naazdeep Kaur², Navpreet Kaur², Peer Junaid², Tyeibah Bashir²,
Dr. Anupama Sharma³, Shikha Sharma⁴ and Dr. Davinder Kaur⁵

1. Nursing Tutor, Gian Sagar College of Nursing, Ramnagar, Rajpura, District Patiala, Punjab.
2. B.Sc. Nursing 4th year students, Gian Sagar College of Nursing, Ramnagar, Rajpura, District Patiala, Punjab.
3. Assistant Professor, Himalayan College of Nursing, SRHU, Jolly Grant Dehradun, U.K.
4. Nursing Tutor, Gian Sagar College of Nursing, Ramnagar, Rajpura, District Patiala, Punjab.
5. Principal, Gian Sagar College of Nursing, Ramnagar, Rajpura, District Patiala, Punjab.

Manuscript Info

Manuscript History

Received: 18 January 2025

Final Accepted: 21 February 2025

Published: March 2025

Key words:-

Safe Delivery App, Maternal Sepsis,
Nursing Education, Mobile Health,
Maternal Health Management

Abstract

Maternal sepsis is a life-threatening condition arising from infections related to childbirth, leading to severe complications or mortality. This study evaluates the effectiveness of the Safe Delivery App in enhancing nursing students' knowledge of maternal sepsis and its management. A quasi-experimental pre-test and post-test design was used. The study was conducted at Gian Sagar College of Nursing, Punjab, with 40 fourth-year B.Sc. Nursing students selected through convenient sampling. Data collection tools included a socio-demographic proforma, a modified structured knowledge questionnaire, and a Safe Delivery App module on maternal sepsis. An orientation session was conducted on app installation and usage, followed by a post-test on the seventh day to assess knowledge improvement. The findings showed a significant increase in students' knowledge and skills regarding maternal sepsis after using the Safe Delivery App, highlighting its effectiveness as an educational tool. The Safe Delivery App serves as a valuable resource for improving nursing students' competency in maternal sepsis management. Its integration into nursing curricula could enhance maternal healthcare education, leading to timely recognition and intervention, ultimately reducing maternal mortality and morbidity rates.

"© 2025 by the Author(s). Published by IJAR under CC BY 4.0. Unrestricted use allowed with credit to the author."

Introduction:-

Maternal and newborn mortality rates remain a critical global health challenge, necessitating innovative interventions to enhance maternal care. The Safe Delivery App (SDA), developed by the Maternity Foundation in collaboration with the University of Copenhagen and the University of Southern Denmark, is a mobile health (mHealth) tool designed to improve healthcare providers' knowledge and skills in managing childbirth-related complications. The app serves as an instructional guide, offering evidence-based protocols for obstetric emergencies, thereby strengthening maternal and neonatal healthcare services, particularly in resource-limited settings¹

Corresponding Author:- Kavita Thakur

Address: - Nursing Tutor, Gian Sagar College of Nursing, Ramnagar, Rajpura,
District Patiala, Punjab.

The Safe Delivery App provides healthcare professionals with instant access to evidence-based guidelines for handling childbirth complications, including maternal sepsis. Available in multiple languages and customizable to national guidelines, the app functions offline once downloaded, ensuring accessibility in remote areas. It integrates instructional videos, drug lists, and e-learning tools to enhance professional competency in Basic Emergency Obstetric and Neonatal Care (BEmONC). Additionally, the app's "My Learning" platform enables users to test their knowledge, achieve certification, and become Safe Delivery Champions².

Maternal sepsis, a severe and life-threatening complication, results from an uncontrolled immune response to infection during pregnancy, childbirth, or the postpartum period. It remains a leading cause of maternal mortality worldwide, particularly in low-resource settings where timely diagnosis and intervention are challenging. The World Health Organization (WHO) has defined maternal sepsis as a life-threatening condition marked by organ dysfunction resulting from infection, highlighting the need for improved detection and management strategies³.

Despite advances in maternal care, the identification of maternal sepsis remains complex due to physiological adaptations in pregnancy that may mask early symptoms. Common clinical signs include fever, tachycardia, hypotension, and altered mental status, necessitating heightened awareness among healthcare providers. The Global Maternal Sepsis Study (GLOSS) emphasized the importance of standardized diagnostic criteria and early intervention to reduce maternal mortality rates associated with sepsis⁴.

Several risk factors predispose pregnant and postpartum women to sepsis, including prolonged rupture of membranes, caesarean delivery, retained products of conception, and invasive intrauterine procedures. Additionally, underlying conditions such as diabetes, obesity, and immunosuppression increase susceptibility to severe infections. Regular training and a structured approach to assessment and management are essential in improving early diagnosis and outcomes for affected women⁵.

The Safe Delivery App aligns with WHO guidelines by providing healthcare professionals with structured, evidence-based protocols for preventing, identifying, and managing maternal sepsis. By integrating digital learning tools into maternal healthcare education, the app empowers midwives and skilled birth attendants with the necessary competencies to enhance maternal and neonatal survival rates. The widespread adoption of such mHealth solutions could significantly contribute to achieving Sustainable Development Goals related to maternal and child health⁶.

Need of Study

The time of childbirth poses significant risks for both mother and child. In 2019, 2.4 million newborns died in their first month, with nearly 6,700 deaths occurring daily. Additionally, approximately 2 million babies died during childbirth, highlighting the critical need for improved maternal healthcare⁷. Maternal sepsis remains a leading cause of maternal mortality, accounting for nearly half of intra-hospital maternal deaths worldwide, as reported by the World Health Organization (WHO) in 2020⁸. Despite declining maternal mortality rates, sepsis continues to be an overlooked contributor, necessitating urgent attention to achieve Sustainable Development Goal 3 (SDG3)⁹. Recent studies indicate a rise in sepsis cases, with reports from Imperial College London showing a 30% increase within two years¹⁰. Sepsis is responsible for 50% of intra-hospital maternal deaths globally, affecting both low- and high-income countries. However, data on the incidence and clinical outcomes of obstetric infections, including maternal sepsis, remains limited¹¹.

India accounts for 32,000 of the 2, 95,000 annual global maternal deaths, representing 20% of maternal fatalities worldwide¹². The primary cause is the lack of skilled care during childbirth. Investing in skilled healthcare could prevent an estimated 1, 13,000 maternal deaths, 5, 31,000 stillbirths, and 1.325 million neonatal deaths annually¹³.

Addressing maternal sepsis through enhanced healthcare strategies, digital learning tools, and policy interventions is essential for reducing maternal and neonatal mortality rates globally.

Material and Method:-

After approval from Gian Sagar ethical committee and permission taken from Gian Sagar College of Nursing. The structured questionnaire was provided to the students. By enrolling 40 nursing students from BSc nursing 4th year from Gian Sagar college of Nursing Ramnagar, Rajpura, Patiala. Convenient sampling technique was used to select the samples. A self-structure questionnaire was formulated and validated by 10 experts in the fields of Obstetrics and Gynaecology. The reliability of tool was checked by split half method ($r = .8$). The tool was divided into 3 sections.

Section-I is Socio Demographic Variable likes age, gender, parents' occupations, mobile phone use, prior awareness of the Safe Delivery App, study time on the app, and exposure to maternal sepsis cases, Section-II is Self-structured knowledge questionnaire consist of 20 multiple choice question, Section-III is Safe delivery app module for Maternal sepsis. After the data collection excel sheet was filled. One mark given for right answer and zero for wrong answer. The criteria used for the assessment of knowledge were Good knowledge score (13-20), Average knowledge score (7-12), Poor knowledge score (0-6). All the students were informed about the study. The anonymity and confidentiality of the participants about findings while reporting the study. Data was analysed by using excel, for descriptive statistics use of percentage, mean, median and SD were calculated.

Results:-

Section 1- Frequency and percentage distribution of socio-demographic variable of nursing students.

Section 2- Findings related to the effectiveness of safe delivery app on level of knowledge regarding maternal sepsis and its management.

Section 3- Findings related to the association between knowledge regarding maternal sepsis and its management with selected demographic variables of nursing students.

Section 1- Frequency and percentage distribution of socio-demographic variable of nursing students.

Table 1:- Frequency and percentage distribution of socio-demographic variable of nursing students.

N-40

Socio-demographic variable	Frequency (f)	Percentage (%)
Age of student in year		
≤21 years	2	5%
22 years	20	50%
23 years	12	30%
≥24 years	6	15%
Gender of student		
Male	13	32.5%
Female	27	67.5%
Occupation of father		
Home maker	0	0
Self employed	19	47.5%
Medical professional	4	10%
Other professional	17	42.5%
Occupation of mother		
Homemaker	34	85%
Self employed	0	0
Nursing professional	0	0
Other professional	6	15%
You are using mobile for which purpose?		
Social media	1	2.5%
Study purpose	1	2.5%
Social media and study both	38	95%
Have you heard about safe delivery app (SDA) earlier?		
Yes	19	47.5%
No	21	52.5%
What was your estimated time duration for studying maternal sepsis while using safe delivery app?		
<1 hour	15	37.5%
2 hours	21	52.5%
3 hours	3	7.5%
>4 hours	1	2.5%
Have you ever seen a patient with maternal sepsis?		
Yes	13	32.5%
No	27	67.5%

Table-1 This table gives details about 40 nursing students and how they study. Most students are 22 years old (50%) and more than half are female (67.5%). The higher influence of female in the sample likely influence these findings. Almost half of their fathers are self-employed (47.5%), while most of their mothers are homemakers (85%). Almost all students (95%) use their phones for both studying and social media. About half of them (47.5%) have heard of the Safe Delivery App (SDA). Most students spent 2 hours (52.5%) using the app to learn about maternal sepsis. However, only 32.5% have seen a patient with maternal sepsis. These details show both their background and learning habits.

Section 2-Findings related to the effectiveness of safe delivery app on level on Knowledge regarding maternal sepsis and its management.




Fig 1:- Percentage Distribution of Subjects according to their level of Knowledge regarding maternal sepsis and its management shows the distribution of post-test knowledge scores among participants regarding maternal sepsis and its management. Out of 40 participants, 80% (32 individuals) achieved a "Good" level of knowledge with scores between 13-20, reflecting a strong grasp of the subject matter. 15% (6 individuals) scored in the "Average" range (7-12), indicating moderate understanding. Meanwhile, 5% (2 individuals) fell into the "Poor" category with scores between 1-6, suggesting limited comprehension. This data highlights that the majority of participants demonstrated a satisfactory to strong level of knowledge post-test, while a small percentage showed room for improvement.

Table2 (a):- Frequency and percentage distribution of post-test knowledge scores regarding maternal sepsis and its management.

N-40

Level of Post-Test Knowledge	Scores	Frequency	Percentage
Good	13-20	32	80%
Average	7-12	6	15%
Poor	1-6	2	5%

Table-2 (a) shows the distribution of post-test knowledge scores among participants regarding maternal sepsis and its management. Out of 40 participants, 80% (32 individuals) achieved a "Good" level of knowledge with scores between 13-20, reflecting a strong grasp of the subject matter. 15% (6 individuals) scored in the "Average" range (7-12), indicating moderate understanding. Meanwhile, 5% (2 individuals) fell into the "Poor" category with scores between 1-6, suggesting limited comprehension. This data highlights that the majority of participants demonstrated a satisfactory to strong level of knowledge post-test, while a small percentage showed room for improvement.

Table-2(b):- Mean, standard deviation and t- value, degree of freedom of knowledge score regarding maternal sepsis.

N-40

Descriptivestatistics	Mean	SD	T – value	df
KnowledgeScore	14	3.587	24.680	39

Table-2 (b) reveals that after the test, participants' understanding of maternal sepsis was measured using a questionnaire. Out of a possible score of 20, the average score was 14, indicating that most participants understood about 70% of the concepts. The variability in scores, represented by the standard deviation of 3.587, shows that while some scores were higher or lower, most were close to the average. The statistical results showed a t-value of 24.680 with 39 degrees of freedom, confirming that the improvement in knowledge was not random, it was significant. This data clearly shows that participants gained meaningful knowledge after the test.

Table3:- Item wise frequency distribution analysis of responses given by nursing students on post-test questionnaire regarding maternal sepsis and its management.

N-40

Items	F	%
Meaning of sepsis	40	100%
Common sign & symptom of maternal sepsis	30	75%
First identification of maternal sepsis can be done by	32	80%
Signs of sepsis diagnosis	34	85%
Without signs of septic shock, time interval for fluid balance calculation in sepsis	15	37.5%
Importance of assessing fluid balance of a female in shock	36	90%
Term used for uterine infection during antenatal period	33	82.5%
True about tetracycline drug	26	65%
Time interval for fluid balance calculation after septic shock	23	57.5%
Drug to avoid during pregnancy and breastfeeding	30	75%
Route of artesunate drug administration in maternal sepsis	19	47.5%
Drug(s) appropriate for treating malaria in maternal sepsis	21	52.5%
Fluid balance in maternal sepsis	28	70%
Term used for infection occurred after delivery in the womb (uterus)	35	87.5%
Critical complications post sepsis	26	65%
Treatment for chorioamnionitis	25	62.5%
Treatment for mastitis	32	80%
Correct dosage for metronidazole in sepsis	20	50%
Antibiotics given in maternal sepsis	33	82.5%
Appropriate dosage of gentamicin in sepsis	22	55%

Table-3 Shows that 40(100%) knowledge among nursing students regarding meaning of maternal sepsis, 30 (75%) knowledge among nursing students regarding common sign and symptoms of maternal sepsis, 32 (80%) knowledge among nursing students regarding first identification of maternal sepsis can be done by, 34 (85%) knowledge among nursing students regarding signs of sepsis diagnosis, 15 (37.5%) knowledge among nursing students regarding Without sign of septic shock, time interval for fluid balance calculation in sepsis, 36 (90%) knowledge among nursing students regarding importance of assessing fluid balance of a female in shock, 33 (82.5%) knowledge among nursing students regarding term used for uterine infection during antenatal period, 26 (65%) knowledge among nursing students regarding true about tetracycline drug, 23 (57.5%) knowledge among nursing students regarding time interval for fluid balance calculation after septic shock, 30 (75%) knowledge among nursing students regarding drug to avoid during pregnancy and breastfeeding, 19 (47.5%) knowledge among nursing students regarding route of artesunate drug administration in maternal sepsis, 21(52.5%) knowledge among nursing students

regarding drugs appropriate for treating malaria in maternal sepsis, 28 (70%) knowledge among nursing students regarding fluid balance in maternal sepsis, 35 (87.5%) knowledge among nursing students regarding term used for infection occurred after delivery in the womb (uterus), 26 (65%) knowledge among nursing students regarding critical complication post sepsis, 25 (62.5%) knowledge among nursing students regarding treatment for chorioamnionitis, 32 (80%) knowledge among nursing students regarding treatment for mastitis, 20 (50%) knowledge among nursing students regarding correct dosage for metronidazole in sepsis, 33 (82.5%) knowledge among nursing students regarding antibiotic given in maternal sepsis, 22 (55%) knowledge among nursing students appropriate dosage of gentamycin in sepsis.

Table4:- Findings related to completion status of Module on maternal sepsis on Safe delivery app.

N-40

Levels of Module		F	%	Completion Status
Level-I	Familiar (1-12)	4	10%	Incomplete
Level-II	Proficient (1-12)	9	22.5%	Partially Complete
Level-III	Expert (1-12)	27	67.5%	Complete

Table-4 highlights the completion status of a module on maternal sepsis through the Safe Delivery app among 40 participants. It shows that 4 individuals (10%) are at the basic "Familiar" level, with their progress marked as incomplete. Nine participants (22.5%) are at the intermediate "Proficient" level, indicating partial completion. The majority, 27 individuals (67.5%), have achieved the "Expert" level, signifying full completion of the module. This indicates that most participants have successfully engaged with and completed the module.

Discussion:-

Maternal sepsis remains a critical public health challenge, contributing to preventable maternal morbidity and mortality worldwide. Effective management depends on healthcare professionals' knowledge and preparedness, including nursing students. This study evaluated the effectiveness of the Safe Delivery App (SDA) as a digital learning tool for enhancing nursing students' knowledge of maternal sepsis and its management in a selected college in Punjab. The study findings revealed a significant improvement in knowledge levels post-intervention. After using the SDA, 80% of students demonstrated good knowledge, 15% had average knowledge, and 5% had poor knowledge. A t-value of 24.680 ($p < 0.00$) confirmed the effectiveness of the SDA in enhancing knowledge. These results align with a study by Joyti Choure et al. (2023), which assessed the SDA's utility among staff nurses in Bhopal, reporting 76% with good knowledge, 16% with average knowledge, and 8% with poor knowledge, with a t-value of 9.4043 ($p < 0.05$), further validating SDA's effectiveness. Additionally, the study examined the association between students' knowledge and demographic variables. Gender was found to be statistically significant ($p = 0.02$), with female students demonstrating stronger knowledge levels. Other factors, such as age, parental occupation, mobile phone use, prior awareness of SDA, study time on the app, and exposure to maternal sepsis cases, did not show significant associations ($p > 0.05$).¹⁴

Conclusion:-

This study highlights the SDA's positive impact on nursing students' knowledge of maternal sepsis. The significant improvement observed post-intervention underscores the potential of digital learning tools in bridging educational gaps and enhancing maternal healthcare outcomes.

Conflict Of Interest

There is no conflict of interest.

References:-

1. Thomsen CF, Barrie AMF, Boas IM, Lund S, Sørensen BL, Oljira FG, Tersbøl BP. Health workers' experiences with the SafeDelivery App in West Wollega Zone, Ethiopia: a qualitative study. *Reprod Health*. 2019 May 9;16(1):50. doi: 10.1186/s12978-019-0725-6. PMID: 31072399; PMCID: PMC6506934
2. Usmani S, Chhugani M, Khan M. A study to assess the effectiveness of safe delivery application for pre-service nursing students in a selected college of nursing of New Delhi. *Int J NursMidwif Res*. 2019;6(4):22-27.
3. World Health Organization. (2018). Global action plan for the prevention and control of NCDs 2013-2020. Geneva: World Health Organization. Available from: <https://iris.who.int/handle/10665/272817>

4. Safe Delivery App [Internet]. Maternity Foundation. [cited 2024 Apr 1]. Available from: <https://www.maternity.dk/safe-deliveryapp/>.
5. Sodha TS, Grønbaek A, Bhandari A, Mary B, Sudke A, Smith LT. mHealth learning tool for skilled birth attendants: scaling the Safe Delivery App in India. *BMJ Open Qual.* 2022 Aug;11(Suppl 1):e001928. doi: 10.1136/bmjopen-2022-001928. PMID: 35977730; PMCID: PMC9389095.
6. Laerdal Global Health [Internet]. [cited 2024 Apr 1]. Safe Delivery App. Available from: [https://laerdalglobalhealth.com/Resources/COVID19 res/safe-delivery-app/](https://laerdalglobalhealth.com/Resources/COVID19%20res/safe-delivery-app/).
7. UNICEF. Levels and Trends in Child Mortality: Report 2019 [Internet]. New York: UNICEF; 2019 [cited 2024 May 1]. Available from: <https://www.unicef.org/reports/levels-and-trends-child-mortality-report-2019>
8. Lawson GW, Keirse MJ. Reflections on the maternal mortality millennium goal. *Birth.* 2013 Jun;40(2):96-102. doi: 10.1111/birt.12041. PMID: 24635463.
9. WHO. Frequency and management of maternal infection in health facilities in 52 countries (GLOSS): A 1-week inception cohort study. *Lancet Glob Health.* 2020;8:661-671. doi: 10.1016/s2214-109x(20)30109-1.
10. Kvalvik SA, Zakariassen SB, Overrein S, et al. Obstetric infections and clinical characteristics of maternal sepsis: a hospitalbased retrospective cohort study. *Sci Rep.* 2024;14:6067. doi:10.1038/s41598-024-56486-4.
11. World Health Organization. Trends in maternal mortality 2000 to 2017: Estimates by WHO, UNICEF, UNFPA, World Bank Group and the United Nations Population Division. Geneva: World Health Organization; 2019. Licence: CC BY-NC-SA 3.0 IGO
12. Sorensen BL, Rasch V, Massawe S, Nyakina J, Elsass P, Nielsen BB. Advanced Life Support in Obstetrics (ALSO) and postpartum hemorrhage: A prospective intervention study in Tanzania. *Acta Obstet Gynecol Scand.* 2011;90:609-614.
13. Bhutta ZA, Das JK, Bahl R, Lawn JE, Salam RA, Paul VK, et al. Can available interventions end preventable deaths in mothers, newborn babies, and stillbirths, and at what cost? *Lancet.* 2014;384:347-370.