

RESEARCH ARTICLE

MAGNITUDE OF STUNTING AND UNDERWEIGHT AMONG HIV INFECTED CHILDREN IN A TERTIARY HOSPITAL SOUTH EAST NIGERIA: A CROSS-SECTIONAL ASSESSMENT BETWEEN URBAN AND RURAL INHABITANTS

Chinelo Vivian Okeke^{1,2} Victor Ikechukwu Ogoke², Maureen Anikpe Okumoko³, Ann Oluchi Ikeabbah⁴, Stella Ijeoma Umeh⁵, Abbas Mohammed Abdulsalam⁶ and Oluwafunke Morounkeji Olugbade⁷

.....

- 1. National Obstetrics Fistula Center Abakaliki Ebonyi State South East Nigeria.
- 2. Imo State University Teaching Hospital Orlu South East Nigeria.
- 3. Niger Delta University Teaching Hospital, Okolobiri Yenagoa Bayelsa State South South Nigeria.
- 4. Nnamdi Azikiwe University Teaching Hospital Nnewi Anambra State South East Nigeria.
- 5. Abia State University Teaching Hospital Aba South East Nigeria.
- 6. Abubakar Tafewa Belewa University Teaching Hospital Bauchi Northern Nigeria.
- 7. General Hospital, Ijede, Lagos State South West Nigeria.

Manuscript Info

Manuscript History Received: 27 March 2025 Final Accepted: 30 April 2025 Published:May 2025

*Key words:-*Nutrition, HIV, Children, Malnutrition Abstract

Background: Nutrition plays an important role in infections. Human Imunodeficiency virus infection and Acquired Immune Deficiency Syndrome, is commonly complicated by malnutrition, which results to severe illness. Malnutrition therefore needs to be taken into account to guarantee proper care for the HIV sero-positive child. There is paucity of data on the magnitude of malnutrition among HIV infected children in Nigeria. This study aims to determine the magnitude of malnutrition among HIV infected children at Federal Teaching Hospital Owerri and to compare the anthropometry of rural and urban inhabitants of children living with HIV.

Methods: A cross-sectional study, in which one hundred and two HIV seropositive children majority of whom are on HAART, aged 1-18 years were recruited consecutively over four months at the pediatric infectious disease clinic of the Federal University Teaching Hospital Owerri. Socio demographic variables were collected. Anthropometric characteristics; weight and height of each participant were measured using a stadiometer (RGZ-160 England). The weight and height were classified as normal, underweight, overweight, obese and stunted, normal and tall stature using CDC classification.

Results: 102 Participants aged 1-18 years included 53 females and 49 males, 22.5% of children were underweight, 34.3% were stunted and 5.8% were obesed. All forms of malnutrition both underweight and overweight were more common in children living in rural areas. 14 children from rural areas were underweight while 10 children from urban areas were underweight. Similarly, 3 children from rural areas were overweight as compared to 2 children from urban areas. Stunting was also more common among rural dwellers.

Conclusion: Malnutrition is common among HIV infected children attending the paediatrics infectious disease clinic in Federal University

Teaching Hospital Owerri. Appropriate nutritional care needs be provided for HIV sero positive children.

"© 2025 by the Author(s). Published by IJAR under CC BY 4.0. Unrestricted use allowed with credit to the author."

Introduction:-

Worldwide, both malnutrition and human immunodeficiency virus (HIV) are highly prevalent, particularly in the sub-Saharan Africa region.¹ Among the estimated 38.4 million people living with HIV globally in 2021, 2.73 million were children aged 0-19 years.² In 2018, approximately 49 and 149 million under-five children were stunted and wasted, respectively, and more than 90% lived in low and middle-income countries.³ The magnitude of stunting and wasting in Sub-Saharan Africa varies in the region by as much as 32% and 10%, respectively.⁴ Nigeria accounted for 59% of all new child HIV infections in west and central Africa in 2016. ⁵ In 2018, the UNAIDS, ⁵ estimated that there were about 37.9 million (32.7 million - 44.0 million) people living with HIV 36.2 million adults and about 1.7 million (1.3 million - 2.2 million) children (<15 years).

Nigeria currently has the largest burden of paediatric HIV infection in the world, with an estimated 321,000 children living with the human immunodeficiency virus (HIV) in Nigeria as at 2010. ⁶ HIV/AIDS, poverty, and food insecurity are the main causes of these high under-nutrition problems.⁷ Studies have shown that stunting, under-weight, and wasting are more prevalent among HIV infected children than uninfected children.^{8–10}

HIV/AIDS is associated with nutritional deficiencies in infected children;¹¹ While under nutrition influences disease progression, increases morbidity and lowers survival of HIV infected persons.¹² Additionally, HIV/AIDS has enormous impact on food security of affected households;^{13,14} especially when the most productive members of the family is affected, the household's economic capacities and the agricultural production are reduced, leading to a situation of food insecurity ¹⁵

In malnourished children, lymphoid tissues are damaged, and CD4 T-cell concentration **is** decreased¹⁶. Deficiencies in vitamins and minerals contribute to oxidative stress, which can accelerate immune cell death ¹⁷ and increase HIV replication¹⁸. On the other hand, HIV infection increases the risk of malnutrition, because of a high proinflammatory cytokine activity which can cause growth impairment among children ¹⁹. HIV-related opportunistic infections such as persistent diarrhoea or oral and oesophageal candidiasis have a negative impact on nutritional status among children ²⁰. HIV infection can also indirectly affect the child's nutritional status, when it has an impact on the child's social environment. Furthermore, poor weaning practices among HIV-infected mothers can also have an impact on the child's nutritional status ²¹. Low weight-for-age has been reported in up to 50 % of untreated HIV-infected children in resource-limited settings²². Among children ^{with} severe malnutrition, mortality risk is three times higher in HIV-infected children than in non-HIV-infected children²³. Therefore, nutritional care should be part of the pediatric HIV care.

The World Health Organization recommends that an asymptomatic HIV-infected child should increase his energy requirements by 10 %, compared to a non-infected child; this is extended to 20 to 30 % during symptomatic HIV infection or episodes of opportunistic infections, and up to 50 to 100 % when a severe malnutrition episode occurs ²⁴. A better understanding of this problem is necessary to improve HIV care in children, especially in sub-Saharan Africa.

Other covariates of child malnutrition have been documented including child level factors such as age and birth weight; maternal level factors such as maternal age and education; household level factors such as food insecurity and social economic status; and community level factors such as sanitation and environmental factors. ²⁵⁻²⁷ Therefore malnutrition is a key problem for HIV sero positive children, as it creates a vicious cycle worsening HIV disease by affecting the immune system negatively.

This study therefore aimed to determine the magnitude of malutrition of HIV infected children at Federal University Teaching hospital Owerri, compare its prevalence among urban and rural dwellers and to further advocate that appropriate nutritional care is provided for children living with HIV.

Methods:-

Study design, period and setting.

A hospital based cross-sectional study. Conducted from April 2018 to July 2018 at the paediatric infectious disease clinic of FUTH Owerri Imo State. Owerri is the capital of Imo State in the South East Geopolitical Zone of Nigeria.²⁸ Imo state has an estimated population of about 3,927,563 with about 403,425 people living in Owerri.²⁹ The hospital is the major tertiary health care facility in the state which serves as a referral centre for those living in Owerri and the surrounding towns as well as neighboring states of Abia, Rivers and Anambra.

Inclusion criteria

All HIV positive children aged 1-18 years whose parents consented to the study.

Exclusion criteria

HIV positive children whose parents declined consent.

Study population

HIV positive children aged 1-18years.

Sampling procedure

One hundred and two HIV positive children that met the inclusion criteria were enrolled consecutively.

Data collection

A structured interviewer administered questionnaire was developed and administered to parents and guardians, to obtain child's demographic data, mode of HIV transmission. Physical examination including anthropometry, (weight [kg], height[cm], mid upper arm circumference [MUAC][cm]) was done and documented.

Weight

Weight of each participant was measured using a stadiometer (RGZ-160 England) after standardization using a known weight. The children wore only inner clothing without shoes and they stood on the weighing scale with their head erect and hands loosely hanging by their side. Reading was taken three times in kilogram and the average recorded. The measurement was taken to the nearest 0.1kg.³⁰

Height

Height was measured using a stadiometer (RGZ-160 England). They stood straight without their shoes on the stadiometer, with back on the board, both legs positioned together bringing the knees and ankles together, head held straight up and arms hanging by their side. The movable part of the head piece brought onto the uppermost point of the head and the readings were taken thrice in centimeters and the average recorded. The measurement was taken to the nearest 0.1 cm^{-30}

MUAC

MUAC was measured using a measuring tape. The left upper arm of the children aged 24 - 60 months were exposed. The circumference of the left upper arm was measured at the mid-point between the tip of the shoulder and the tip of the elbow (acromium and olecranon process respectively). The average of two readings was taken. Measurment was taken to the nearest 0.1 cm.³⁰

Data analysis

The data collected was coded and imputed into Statistical Package for Social Sciences (SPSS) version 20.0. Frequency tables and figures were used to present relevant variables.

Descriptive statistics (mean and standard deviation) was used to summarize quantitative variable (age and social class). Qualitative variables were summarized as proportions. The Chi-square test was used to compare associations between proportions.

Bias

Several strategies were employed to minimize bias in the study. Selection bias was addressed by applying strict inclusion and exclusion criteria, ensuring that the study population was representative of the broader pediatric population seen in the HIV clinic. Information bias was minimized through standardized data collection methods and the cross-verification of data sources. Measurement bias was minimized by taking average of multiple measurements asnd standardization of measuring equipments.

Ethical approval

Ethical approval was obtained from the ethics committee of Federal Medical Center Owerri. Informed consent of the parents and assent of 7 years and above child were gotten for the study.

Results:-

A total of 102 patients were recruited from April 2018 to June 2018. 53 were females while 49 were males. Majority (67) are rural dwellers. Majority (95) are on HAART as shown in table 1 below. **Table 1:** Socio demographic Characteristics of participants

Table 1:- Socio demographic Characteristics of participants.							
Variable	Min	Max	Mean	SD			
Age (years)	1	18	9.72	3.11			
MAC (cm)	12.5	14.0	13.16	0.51			

Variable	Frequency	Percentage		
Gender		<u> </u>		
Male	49	48.1		
Female	53	52.0		
Place of Residence				
Rural	67	65.7		
Urban	35	34.3		
Age Band				
0 -4	5	5.1		
5-9	43	43.4		
10-14	39	39.4		
15 – 19	12	12.1		
ART Use				
On ART	95	93.1		
Not on ART	7	7.0		

To determine the magnitude of malnutrition of HIV infected children at FUTH Owerri using BMI, it was observed that twenty three out of 102 (22.5%) of Children living with HIV seen at FUTH Owerri are underweight as depicted in table II

Table II:- Magnitude of malnutrition in Children Living with HIV.

BMI Classification	Frequency	Percentage
Underweight	23	22.5
Normal	69	66.9
Overweight	4	3.9
Obessed	6	5.8
Total	102	100

All forms of Malnutrition both underweight and overweight are commoner among children living with HIV in rural areas compared to urban dwellers, however this did not reach statistically significant level p-value 0.41 as displayed in Table III

Table III:- Nutritiona	l status of Participants	by Place of Residence.
------------------------	--------------------------	------------------------

Nutritional Status	Place of residence n(%)		χ2	p-value
	Rural	Urban		
Underweight	14 (20.8)	10 (28.6)		
Normal	46 (68.6)	21 (60.0)	2.9	0.41
Overweight	3 (4.4)	2 (5.7)		
Obessed	4 (5.9)	2 (5.7)		
Total	67 (100)	35 (100)		

Thirty two out of 102 (31.4%) of children living with HIV in FUTH Owerri are stunted as shown in table IV

Table IV:- Summary of Ht for Age Characteristics of CALHIV in FMC Owerri.

Height for Age	Frequency	Percentage
Stunted	32	31.4
Normal	66	64.7
Tall Stature	4	3.9

Total	102	100

Stunting is commoner among rural dwelling CALHIV compared to urban counterparts though this is not statistically significant as seen in table V

Height for Age	Place of residence		χ2	p-value
	Rural	Urban		
Stunted	23 (34.3)	9 (25.7)		
Normal	43 (64.2)	23 (65.7)	2.86	0.24
Tall Stature	1 (1.5)	3 (8.6)		
Total	67 (100)	35 (100)		

Table	V. Ca		of IIt for	A and hateriaan	Dunal and	I Inhon du	vallin a C	1 A T TTTT
I able	V:- CO	mparison	OI HUIOI	: Age between	i Kurai and	Urban dv	vening C	ALHIV.

Discussion:-

This study aimed to determine the magnitude of malnutrition amongst HIV infected children aged 1-18 years attending the federal Teaching hospital Owerri Imo state South East Nigeria, and to compare the anthropometry of urban and rural dwelling children living with HIV.

The current study shows that 22.5% of the study population were underweight. Further more, 34.3% of participants were stunted. This is similar to the Ibadan study by Okafor et al. ³¹ Results in this study were also similar to those in the Oyo study by Fagbamigbe et al. ³²

Prevalence of underweight and stunting in this study were lower than the Central and West African study by Jesson ³³ et al and Malawi study by Weigel et al ³⁴ whom observed that almost half and more than half of their study participants respectively were stunted and underweight. The differences in the results may be due to the fact that majority of the study participants in the current study were on HARRT and were not immunosuppresed, as compared to the Malawi study whom the participants were just initiating HARRT as at the time of the study.

This study also depicts that underweight and stunting were more common in rural than urban dwellers. This finding was similar to that by Alyson et al in the USA study. ³⁵ This may be due to the fact that people living in rural communities face significant barriers to accessing prevention and treatment services for HIV. These communities have fewer options for health care. This means rural residents usually need to travel for health care, often with the added burden of transportation.

Health care providers in the rural areas may also lack the knowledge or training to treat HIV and often don't understand the complex medical needs of those living with HIV, which involves treating ongoing issues like malnutrition, depression, pain, and substance use disorder.

Rural communities may not have access to health insurance and financial resources like those in urban areas. Those with HIV need help paying for treatment if they don't have health insurance.

People with the disease face negative beliefs about their condition due to stigma. When you live in a small, closely connected community, it could be difficult maintaining confidentiality while getting the care you need. This may result to refusal to get tested and treated. Some HIV/AIDS groups even avoid advertising their services in rural areas to avoid stigma.³⁵

Limitations of study

This is a single center study. A multi center study may provide a broader view of magnitude of malnutrition in HIV infected children.

Conclusion:-

Underweight and stunting are common in HIV infected children at FUTH Owerri. Stunting and underweight are also more common in rural areas than in urban dwellers. We therefore recommend that HIV infected children be given adequate nutritional care including micronutrient deficiency. Also, quality health care for rural areas regarding HIV diagnosis and treatment is advocated.

Conflict of interest:

Authors declare no conflict of interest.

Funding:

Study received no external funding

Author contributions:

Chinelo Vivian Okeke: conceptualization of study, writing, expertise, review, data collection, data analysis, result interpretation, supervision.

Victor Ikechukwu Ogoke: writing, review, result interpretation, expertise

Maureen Anikpe Okumoko: writing, review, result interpretation.

Ann Oluchi Ikeabbah: writing, review, data analysis

Stella Ijeoma Umeh: writing, review, data analysis

Abbas Mohammed Abdulsalam: conceptualization, writing, expertise, review.

Oluwafunke Morounkeji Olugbade: writing, review, data analysis

References:-

- Gedle D, Gelaw B, Muluye D, Mesele M. Prevalence of malnutrition and its associated factors among adult people living with HIV/AIDS receiving anti-retroviral therapy at Butajira hospital, southern Ethiopia. BMC Nutri. 2015;1(1):1–11. doi: 10.1186/2055-0928-1-5. [CrossRef] [Google Scholar]
- 2. HIV Statistics-Global and Regional Trends- UNICEF DATA https://data,unicef.org/topic/hivaids/global-regional-trends.
- 3. Organization WH. UNICEF/WHO/the World Bank Group joint child malnutrition estimates: levels and trends in child malnutrition: key findings of the. 2020th ed; 2020.
- 4. UNCEF: Children, food and nutrition growing well in a changing world. In.; 2019.
- 5. United States Agency for International Development (UNAIDS) (2017) Ending AIDS: Progress towards the 90-90-90 Targets.
- 6. World Health Organization (WHO) (2012) Global Monitoring Framework and Strategy for the Global Plan towards the Elimination of New HIV Infections among Children by 2015 and Keeping Their Mothers Alive (EMTCT).
- Friedman JF, Kwena AM, Mirel LB, Kariuki SK, Terlouw DJ, Phillips-Howard PA, Hawley WA, Nahlen BL, Shi YP, Ter Kuile FO. Malaria and nutritional status among pre-school children: results from cross-sectional surveys in western Kenya. Am J Trop Med Hyg. 2005;73(4):698–704. doi: 10.4269/ajtmh.2005.73.698. [PubMed] [CrossRef] [Google Scholar]
- Sunguya BF, Poudel KC, Otsuka K, Yasuoka J, Mlunde LB, Urassa DP, Mkopi NP, Jimba M. Undernutrition among HIV-positive children in Dar Es Salaam, Tanzania: antiretroviral therapy alone is not enough. BMC Public Health. 2011;11(1):869. doi: 10.1186/1471-2458-11-869. [PMC free article] [PubMed] [CrossRef] [Google Scholar]
- Anyabolu HC, Adejuyigbe EA, Adeodu OO. Undernutrition and anaemia among HAART-naïve HIV infected children in Ile-Ife, Nigeria: a case-controlled, hospital based study. Pan Afr Med J. 2014;18(1). 10.11604/pamj.2014.18.77.3746. [PMC free article] [PubMed]
- Poda GG, Hsu C-Y, Chao JC. Malnutrition is associated with HIV infection in children less than 5 years in Bobo-Dioulasso City, Burkina Faso: A case–control study. Medicine. 2017;96(21). 10.1097/MD.00000000007019. [PMC free article] [PubMed]
- 11. Wilson D, Pencharz P: Nutritional Care of the chronically ill. Nutrition during infancy: birth to 2 years. 1997, Cincinnati: Digital Educational Publishing Inc, 37-46
- 12. Berhane R, Bagenda D, Marum L, Aceng E, Ndugwa C, Bosch RJ, Olness K: Growth failure as a prognostic indicator of mortality in pediatric HIV infection. Pediatrics. 1997, 100 (1): E7-10.1542/peds.100.1.e7.
- Hunter LM, Twine W, Patterson L: "Locusts are now our beef": adult mortality and household dietary use of local environmental resources in rural South Africa. Scand J Public Health. 2007, 35 (Suppl 69): 165-174. 10.1080/14034950701356385.
- 14. de Waal A, Whiteside A: New variant famine: AIDS and food crisis in southern Africa. Lancet. 2003, 362 (9391): 1234-1237. 10.1016/S0140-6736(03)14548-5.
- 15. Anema A, Vogenthaler N, Frongillo EA, Kadiyala S, Weiser SD. Food insecurity and HIV/AIDS: current knowledge, gaps, and research priorities. Curr HIV/AIDS Rep. 2009;6(4):224–31.

- Cunningham-Rundles S, McNeeley DF, Moon A. Mechanisms of nutrient modulation of the immune response. J Allergy Clin Immunol. 2005;115(6):1119–28.
- 17. Romero-Alvira D, Roche E. The keys of oxidative stress in acquired immune deficiency syndrome apoptosis. Med Hypotheses. 1998;51(2):169–73.
- 18. Allard JP, Aghdassi E, Chau J, Tam C, Kovacs CM, Salit IE, et al. Effects of vitamin E and C supplementation on oxidative stress and viral load in HIV-infected subjects. AIDS. 1998;12(13):1653–9.
- Johann-Liang R, O'Neill L, Cervia J, Haller I, Giunta Y, Licholai T, et al. Energy balance, viral burden, insulinlike growth factor-1, interleukin-6 and growth impairment in children infected with human immunodeficiency virus. AIDS. 2000;14(6):683–90
- 20. Trehan I, O'Hare BA, Phiri A, Heikens GT. Challenges in the Management of HIV-Infected Malnourished Children in Sub-Saharan Africa. AIDS Res Treat. 2012;2012:790786
- 21. Saloojee H, De Maayer T, Garenne ML, Kahn K. What's new? Investigating risk factors for severe childhood malnutrition in a high HIV prevalence South African setting. Scand J Public Health Suppl. 2007;69:96–106
- 22. Anabwani G, Navario P. Nutrition and HIV/AIDS in sub-Saharan Africa: an overview. Nutrition. 2005;21(1):96–9
- Chinkhumba J, Tomkins A, Banda T, Mkangama C, Fergusson P. The impact of HIV on mortality during inpatient rehabilitation of severely malnourished children in Malawi. Trans R Soc Trop Med Hyg. 2008;102(7):639–44
- 24. WHO. Training Course on Child Growth Assessment. Module B: Measuring a child's growth. Geneva, Switzerland: WHO. 2008. Available at:
 - [http://www.who.int/childgrowth/training/module_b_measuring_growth.pdf]
- 25. Hien NN, Kam S: Nutritional status and the characteristics related to malnutrition in children under five years of age in Nghean, Vietnam. J Prev Med Public Health. 2008, 41 (4): 232-240. 10.3961/jpmph.2008.41.4.232.
- 26. Linnemayr S, Alderman H, Ka A: Determinants of malnutrition in Senegal: Individual, household, community variables, and their interaction. Econ hum Biol. 2008, 6 (2): 252-263. 10.1016/j.ehb.2008.04.003.
- 27. Pongou R, Ezzati M, Salomon JA: Household and community socioeconomic and environmental determinants of child nutritional status in Cameroon. BMC Public Health. 2006, 6: 98-10.1186/1471-2458-6-98.
- 28. Legal notice on publication of the details of the breakdown of the national and state provisional total 2006 census. Constitution of Federal Republic of Nigeria 2007;94:175
- 29. National Bureau of statistics. Annual Abstract of Statistics 2011. Federal Republic of Nigeria.[online] Available from istmat.info/files/uploads/53129/annual_abstract_of_statistics_2011.pdf (Accessed 2018 Sept 25).
- Center for disease control and prevention. National Health and Nutrition Examination Survey. Anthropometry procedures, manual 2007. Available from www.cdc.gov/nchs/ data/nhanes/0708/manuel on.pdf. (Accesed 2019 June12).
- Okafor, C., Fadupin, G. and Oladokun, R. (2021) Nutritional Status and Virological Outcomes of Children HIV Positive Attending Anti-Retroviral Clinic at University College Hospital, Ibadan. Food and Nutrition Sciences, 12, 1088-1097. doi: 10.4236/fns.2021.1211080.
- 32. Fagbamigbe, A.F., Adebowale, A.S. and IkeOluwapo, A. (2019) An Assessment of the Nutritional Status of ART Receiving HIV-Orphaned and Vulnerable Children in South-West Nigeria. Heliyon, 5, e02925.
- Jesson, J., Masson, D., Adonon, A. et al. Prevalence of malnutrition among HIV-infected children in Central and West-African HIV-care programmes supported by the Growing Up Programme in 2011: a cross-sectional study. BMC Infect Dis 15, 216 (2015).
- Weigel R, Phiri S, Chiputula F, Gumulira J, Brinkhof M, Gsponer T, et al. Growth response to antiretroviral treatment in HIV-infected children: a cohort study from Lilongwe, Malawi. Tropical Med Int Health. 2010;15(8):934–44.
- 35. Alyson P and Carmelita S. HIV in rural areas. HIV and AIDS. 2022 April 15.