



Journal Homepage: [-www.journalijar.com](http://www.journalijar.com)

## INTERNATIONAL JOURNAL OF ADVANCED RESEARCH (IJAR)

Article DOI:10.21474/IJAR01/21025

DOI URL: <http://dx.doi.org/10.21474/IJAR01/21025>



### RESEARCH ARTICLE

## A CASE REPORT OF INTENSIVE CARE MANAGEMENT OF REFRACTORY MYASTHENIA GRAVIS WITH ACUTE RESPIRATORY FAILURE: DIAGNOSIS, TREATMENT AND MANAGEMENT

Poonam Negi<sup>1</sup>, Dhenesh Rajavel<sup>2</sup>, Senthilnathan<sup>3</sup> and Manjunath G<sup>4</sup>

1. Department of Critical care, Voluntary Health Services, Taramani, Chennai, India.
2. Department of Anaesthesia and critical care, Delta hospitals, Adambakkam, Chennai.
- 3 Department of Neurology, Voluntary Health Services, Taramani, Chennai, India.
4. Department of CriticalCare, St. Marthas Hospital, Bengaluru, Karnatka, India.

### Manuscript info

#### Manuscript History

Received: 27 March 2025

Final Accepted: 30 April 2025

Published: May 2025

### Abstract

Myasthenia gravis is an autoimmune disorder causing muscle weakness and fatigability due to the antibody against acetylcholine receptor. The disease is characterised by frequent relapse and remission and respiratory distress resulting in myasthenic crisis. Management of disease revolves around suppression of symptoms and disease pathology. Treatment options include acetylcholinesterase inhibitors, steroids, immunosuppressants and plasma exchange therapies. We present a case of 24-year-old female with a known case of myasthenia gravis who showed failure to conventional therapy with pyridostigmine, steroids and IVIG. She received 5 cycles of PLEX to overcome the myasthenic crisis followed by rituximab. She was kept on a maintenance dose of steroids and immunosuppressants. Her course in ICU was stormy with respiratory distress requiring intubation and mechanical ventilation and development of VAP. She was extubated after 4 cycles of PLEX and VAP management.

"© 2025 by the Author(s). Published by IJAR under CC BY 4.0. Unrestricted use allowed with credit to the author."

### Introduction:

Myasthenia gravis (MG) is an autoimmune neuromuscular disorder characterized by fluctuating muscle weakness, often involving ocular, bulbar, and respiratory muscles. A myasthenia crisis (MC) is a life-threatening condition involving respiratory failure due to diaphragmatic and intercostal muscle weakness, requiring mechanical ventilation and intensive care support.

It affects a substantial number of individuals, leading to considerable morbidity due to its unpredictable nature and potential for severe complications. The clinical implications are profound, as the fluctuating nature of muscle weakness may lead to respiratory complications, worsening patient care. As the incidence of MG continues to rise, this case study serves as an essential source for medical professionals detailing the intensive care management strategies and multidisciplinary approaches in enhancing patient outcomes. This case study presents the management

of a 24-year-old female with refractory MG in crisis, highlighting the challenges in airway management, plasma exchange therapy (PLEX) and its complications, immunosuppression, and secondary infections in the ICU<sup>1</sup>.

### Case Presentation:

A 24-year-old female, with a known case of non thymomatous myasthenia gravis presented to us with chief complaints of shortness of breath, inability to swallow oral secretions and fever for 1 day.

She was on anticholinesterase inhibitor pyridostigmine 60 mg thrice a day and mycophenolate mofetil 500 mg once daily with a minimal dose of wysolone 5 mg once daily.

Her past history was significant with admission for myasthenia crisis for which she was treated with IVIG (0.4 g / kg / day x 5 days) 2 months back.

She is also a known case of carcinoma ovary for which she underwent Oophorectomy and received chemotherapy in 2010.

On physical examination, patient had a BP of 130/90 mmHg in right upper limb in supine position, HR: 110 bpm, RR: 35/min, SpO<sub>2</sub> – 88% RA, with single breath count of 5. She did not have a sustained neck lift for more than 5 seconds, had a significant proximal muscle weakness of bilateral upper and lower limbs [ 3/5]. Rest all osteotendinous reflexes were normal. Systemic examination of respiratory system and cardiovascular systems were insignificant.

### ICU Management Strategy

In view of respiratory distress, patient was intubated with 7.5 mm ETT under opioid sedation. She was initiated on mechanical ventilation and placed on volume control mode. Laboratory findings revealed Hb of 10.0 g/dL with wbc count of 10,000 and normal renal and liver function test. Her acetylcholine receptor antibody levels were quantified and was found to be more than 8 mmol/L {normal<0.40}. Patient was started on pulse dose of glucocorticoids with injection methylprednisolone 1 gm iv od for 3 days and tapered slowly. A broad-spectrum antibiotic was initiated empirically in view of aspiration pneumonia.

Post pulse dose of steroids, patient did not show any improvement in muscle power of upper and lower limbs with absent sustained neck lift. Neurologist opinion was obtained and suspecting steroids unresponsive myasthenic crisis, it was decided to initiate plasma exchange [PLEX] for 5 cycles, with each cycle on alternate days<sup>2</sup>.

After informed consent, right internal jugular vein was cannulated with 13 cm, 16 fr triple lumen HD catheter. With patients' weight being 60 kg and hematocrit of 31, Total plasma volume (TPV) of 2691 ml was calculated. It was decided to exchange 1-1.5 times of TPV in each cycle over alternate days. On day 3 of ICU admission, patient developed increased ET tube secretions with a radiological opacity seen in left lower lobe on chest x ray. There was an increase in total white cell count to 14,000. Endotracheal tube culture was sent which came out as growth of *Aceinetobacter baumannii* which was carbapenem sensitive. Patient's antibiotics were escalated to meropenem. After the 3<sup>rd</sup> cycle of PLEX, patient started showing improvement in muscle power of both the limbs UL 5/5, LL 3/5, sustained neck lift >5 sec.

Weaning trial was initiated and patient was placed on pressure support ventilation. On the 4<sup>th</sup> cycle of PLEX, after successful weaning trial; she was extubated to NIV. The last cycle of PLEX was continued on the next day. Towards the next few days, patient total count started decreasing, radiological regression of the opacity was noticed, her single breath count improved to 20, she was mobilized to chair with the help of physiotherapist and shifted to ward. She was discharged on maintenance dose of tablet wysolone 5 mg, pyridostigmine bromide 60 mg 6<sup>th</sup> hrly and mycophenolate mofetil 500 mg once daily.

Three weeks after the current admission, patient presented to us again with shortness of breath, unable to swallow secretions and respiratory distress. Her MG-ADL (Activities of Daily Living) score was around 18. In view of respiratory distress, patient was electively intubated and started on PLEX again. The PLEX cycles were continued for 5 cycles, alternating every day with 1.5 TPV to be exchanged in every cycle<sup>3,7,8</sup>.

Considering frequent myasthenic relapse, elective tracheostomy was performed. Patient was initiated on injection Rituximab 1000 mg iv stat for 4 hours; 2<sup>nd</sup> dose repeated after 4 weeks. Post rituximab, patient was discharged with tracheostomy tube in situ. Plan to decannulate tracheostomy was made after 4 weeks based on the disease remission.

## Discussion:

Refractory MG was defined as lack of response to treatment with steroids and at least 2 immunosuppressants with an inability to withdraw treatment without relapse in the last 12 months, or intolerance to treatment resulting in adverse reactions<sup>6</sup>. The incidence of refractory MG is 14 – 20 cases per 100,000 individuals, with increased in prevalence noted among women under 40 and men over 60<sup>10</sup>.

This case poses significant challenge for healthcare providers due to its unpredictable nature and potential for severe exacerbations, leading to critical illness or respiratory failure.

Effective management is imperative, as exacerbations, particularly during crisis, can lead to respiratory failure and necessitate intensive care interventions. This case study illustrates the need for prompt, multidisciplinary strategies to enhance patient outcomes in MG crisis, underscoring its relevance for medical professionals and researchers dedicated to improving care protocols.

## Conclusion:

Non-thymomatous myasthenia gravis refractory to steroids with frequent relapses poses a significant challenge to the practitioner<sup>6</sup>. This case exemplifies how prompt interventions, multidisciplinary care, and vigilant ICU monitoring can significantly improve outcomes in MG crisis<sup>9</sup>.

This case also emphasizes the critical role of effective interventions, such as plasma exchange and newer immunosuppressants like rituximab in managing autoimmune respiratory crisis<sup>3,4,5</sup>.

## References:

1. Sanders DB, Wolfe GI, Benatar M, Evoli A, Gilhus NE, Illa I, et al. International consensus guidance for management of myasthenia gravis: Executive summary. *\*Neurology\** (2016) 87:419–25. doi: 10.1212/WNL.0000000000002790
2. Mandawat A, Kaminski HJ, Cutter G, Katirji B, Alshekhlee A. Comparative analysis of therapeutic options used for myasthenia gravis. *\*Ann. Neurol.\** (2010) 68:797–805. doi: 10.1002/ana.22139
3. NIH Consensus Development. The utility of therapeutic plasmapheresis for neurological disorders. *\*JAMA\** (1986) 256:1333–7. doi: 10.1001/jama.256.10.13334
4. Barth D, Nabavi Nouri M, Ng E, Nwe P, Bril V. Comparison of IVIg and PLEX in patients with myasthenia gravis. *\*Neurology\** (2011) 76(23):2017–23. doi: 10.1212/WNL.0b013e31821e5505. Epub 2011 May 11. PMID: 21562253; PMCID: PMC3109880.
5. Narayanaswami P, Sanders DB, Wolfe G, Benatar M, Cea G, Evoli A, Gilhus NE, Illa I, Kuntz NL, Massey J, Melms A, Murai H, Nicolle M, Palace J, Richman D, Verschuuren J. International Consensus Guidance for Management of Myasthenia Gravis 2020 Update. *\*Neurology\** January 19, 2021 issue 96(3):114–122. doi: 10.1212/WNL.0000000000011124
6. DESAI, KARNA et al. REFRACTORY SEROPOSITIVE NONTHYMOMATOUS MYASTHENIA GRAVIS IN MYASTHENIC CRISIS: A CASE REPORT, *CHEST* Volume 166, Issue 4, A5494
7. Gajdos P, Chevret S, Toyka K. Plasma exchange for myasthenia gravis. *Cochrane Database Syst. Rev* 2002 (4) C0002275
8. Guptill JT, Juel VC, Massey JM, Anderson AC, Chopra M, Yi JS, et al. Effect of therapeutic plasma exchange on immunoglobulins in myasthenia gravis. *Autoimmunity*. (2016) 49:472–9. doi: 10.1080/08916934.2016.1214823
9. Linda M, Wendell C, Joshua Levine M. Myasthenic crisis neuroanesthesianeurocritical care. *Case Stud.* (2011) 1:321–3. doi: 10.1177/1941875210382918
10. Cortés-Vicente E, Álvarez-Velasco R, Pla-Junca F, Rojas-García R, Paradas C, Sevilla T, Casasnovas C, Gómez-Caravaca MT et al. Drug-refractory myasthenia gravis: Clinical characteristics, treatments, and outcome. *Ann ClinTranslNeurol.* 2022 Feb;9(2):122–131. doi: 10.1002/acn3.51492. Epub 2022 Jan 26. PMID:35080153; PMCID: PMC8862423.