

Journal Homepage: -www.journalijar.com

INTERNATIONAL JOURNAL OF ADVANCED RESEARCH (IJAR)

EVIENATIONAL POEMAE OF ABITANCES RESEARCH STARS

Article DOI:10.21474/IJAR01/21040
DOI URL: http://dx.doi.org/10.21474/IJAR01/21040

RESEARCH ARTICLE

A QUASI EXPERIMENTAL STUDY TO ASSESS THE KNOWLEDGE REGARDING PRE-MENSTRUAL SYNDROME AND ITS MANAGEMENT AMONG EARLY ADOLESCENT GIRLS IN SELECTED GOVT. SCHOOL IN JALANDHAR, PUNJAB, INDIA

Priya¹ and Sukhmeet Kaur²

.....

- 1. Assistant Professor, M.H.R. DAV Institute of Nursing Jalandhar, Punjab, India.
- 2. Clinical Instructor, M.H.R. DAV Institute of Nursing Jalandhar, Punjab, India.

Manuscript Info

Manuscript History Received: 27 March 2025 Final Accepted: 30 April 2025 Published:May 2025

Key words:

Pre-menstural syndrome, Knowledge and early adolescent girls

Abstract

A quasi-experimental Study was undertaken to assess the Knowledge regarding Pre-menstrual Syndrome and its management among early adolescent in selected govt. school in Jalandhar, Punjab, India.

Objectives:To assess the level of knowledgeregarding Pre-menstrual Syndrome and its management among early adolescent girls.

To provide knowledgeregarding Pre-menstrual Syndrome with the help of Power point presentation and provide pamphlets regarding its management.

To determine the association between the level of knowledge with selected demographic variables and self structured tools.

Methodology: A quantitative research approach with quasi-experimental research design was used in the study. Research setting was government middle school Jalandhar Punjab India. Total 60 samples were selected with non-Probability purposive sampling technique. Verbal consent were taken from the students. Self-structured tool was used to assess the knowledge regarding premenstrual syndrome among early adolescent girls.

Result and conclusion: The Present study was done on 60 early adolescent girls to assess the Knowledge regarding Pre menstrual syndrome in selected government school, Jalandhar, Punjab India. The study comparison was done based on pre test and posttest level of Knowledge Pre test 7.53 + 2.581. Post test 19.63+ 1.84. Hence, the finding revealed the significant improvement in the knowledge.

Recommendation: Similar studies can be replicated with large samples withgeneralization. A study can be conducted to evaluate the knowledge and the attitude of adolescent girls regarding PMS.

An exploratory study can be done at various settings to identify factors influencing PMS.

[&]quot;© 2025 by the Author(s). Published by IJAR under CC BY 4.0. Unrestricted use allowed with credit to the author."

.....

Introduction:-

PMS can significantly impact a woman's daily life, affecting her relationships, work, and overall well-being. Understanding and managing PMS symptoms can help improve quality of life for those affected⁽¹⁾. Premenstrual syndrome (PMS) is a common condition affecting women of reproductive age Global Prevalence: Estimates range from 30% to 80% of women experiencing some symptoms of PMS, with 20% to 40% experiencing severe symptoms that impact daily life.⁽²⁾ According to a systematic review and meta-analysis, the pooled prevalence of PMS in India is approximately 43% (95% CI: 0.35-0.50). ⁽³⁾ This translates to about 4 out of 10 women experiencing PMS symptoms. Prevalence Rates in Different Age Groups: Adolescents: The prevalence of PMS is higher among adolescents, at around 49.6% (95% CI: 0.40-0.59). Adult Women: The prevalence rates vary across different studies, but overall, it's estimated that around 40-50% of adult women in India experience PMS symptoms. Regional Variations:- Studies have reported varying prevalence rates across different states in India, ranging from 14.3% to 74.4%. Some states with notable prevalence rates include: Tamil Nadu: 74.4% (Kavitha et al., 2015) and 14.3% (Durairaj et al., 2019) Gujarat: 42% (Brahmbhatt et al., 2013) and 18.97% (Kamat et al., 2019) West Bengal: 42.38% (Ray et al., 2010) and 53.96% (Mandal et al., 2015)

Research Problem

A quasi-experimental Study was undertaken to assess the Knowledge regarding Pre-menstrual Syndrome and its management among early adolescent in selected govt. school in Jalandhar, Punjab, India.

Objectives:-

- 1. To assess the level of knowledgeregarding Pre-menstrual Syndrome and its management among early adolescent girls.
- 2. To provide knowledgeregarding Pre-menstrual Syndrome with the help of Power point presentation and provide pamphlets regarding its management.
- 3. To determine the association between the level of knowledge with selected demographic variables and self structured tools.

Research Methodology:-

Research approach:

Quantitative (experimental research approach)

Research design:

Quasi-experimental design

Setting of the study:

Government middle school Punjab India.

Target Population:

Early adolescent age group 11 to 14 years.

Sampling sizes:

60 students

Tool description:

Self-structured tool was formed to assess the knowledge regarding premenstrual syndrome among early adolescent girls.

Tool was develop in the following ways:

Part –I Socio-demographic variables This part consist of items for obtaining personal information from the adolescent girls i.e. Age, Type of family, current academic class, sleeping pattern, food habits, play time, socioeconomic status, Type of school, Medication and duration of menstrual cycle.

Part -II - Self Structured Tool: The self structured tool was divided into three sections in the form of Likert Scale⁽⁵⁾.

Section I: Knowledge about PMS (Premenstrual syndrome)

Section II: Attitude toward PMS (Premenstrual syndrome)
Section III: Experiences with PMS (Premenstrual syndrome)

Each sections contain Five Questions on the basis of Strongly agree, Agree, Neutral, disagree and strongly disagree. The scoring was done based on poor knowledge (15-35), average knowledge (36-50) and adequate knowledge (51-75).

Reliability of tool:

Reliability was commuted and calculated by using Karl Pearson's coefficient of Correlation ⁽⁶⁾ and Spearman's Brown Prophecy ⁽⁷⁾ formula the reliability of the tool was 0.77. Hence, the tool was reliable.

Results:-

The analysis was done using descriptive and inferential statistic

Findings related to socio-demographic variables:

This part consist of items for obtaining personal information from the adolescent girls i.e. Age, Type of family, current academic class, sleeping pattern, food habits, play time, socioeconomic status, Type of school, Medication and duration of menstrual cycle.

Table I:- Table I are showing a frequency and % Percentage distribution of socio-demographic variables. **N=60**

Sr. no.	Socio demographic variables	Frequency(f)	Percentage (%)
1.	Age (in years)		
	10-11	01	1.70
	11-12	08	13.30
	12-13	22	36.60
	13-14	29	48.40
2.	Type of family		10,10
	Nuclear Family	30	50
	Joint Family	30	50
	John Lamily		
3.	Current academic Class		
	5 th	40	66.70
	6 th	20	33.30
	7 th	00	
4.	Sleeping pattern		
	Less than 8 hours	35	58.40
	More than 8 hours	25	41.60
	William O mours	23	11.00
5.	Food Habits		
	Vegetarian	38	63.40
	Non- Vegetarian	20	33.40
	Vegan	00	00
	Eggetarian	02	3.20
6.	Play time		
	Less than 1 hour	05	8.40
	1-3 hours	50	83.40
	More than 3hours	05	8.20
7.	Socio economic status		
	Below poverty line	40	66.70
	Above poverty line	20	33.30
8.	Type of school		
-	Private	00	00
	Semi-government	00	00
	Government	60	100
9.	Medication		100

	Never used	30	50	
	When needed	30	50	
	Always	00	00	
10	Duration of menstrual cycle			
	Less than 28 days	10	16.66	
	28 days to 32 days	40	66.70	
	More than 32 days	10	16.66	

Results pertaining Knowledge among early adolescent girls on pre menstrual syndrome Pre-Test was conducted, than the intervention i.e. structured teaching program and pamphlets were given leading to the posttest

Table II:-Mean and Mean percentage and standard deviation of pre test N=60

Early adolescent girls	Mean	Standard Deviation	Mean Percentage
Knowledge on PMS	7.53	2.581	26.90

Table III:-Mean and Mean percentage and standard deviation of post test. N=60

Early adolescent girls	Mean	Standard Deviation	Mean Percentage
Knowledge on PMS	19.63	11.389	70.12




Fig. 1:- Depicting mean and standard deviation of pre and posttest.

Conclusion:-

The Present study was done on 60 early adolescent girls to assess the Knowledge regarding Pre menstrual syndrome in selected government school, Jalandhar, Punjab India. The study comparison was done on the basis of pre test and post test level of Knowledge Pre test- 7.53 + 2.581. Post test 19.63+ 1.84. Hence the finding revealed the significant improvement in the knowledge.

References:-

1-Premenstrual syndrome (PMS) - Symptoms & causes - Mayo Clinic. (2022, February 25). Mayo Clinic. https://www.mayoclinic.org/diseases-conditions/premenstrual-syndrome/symptoms-causes/syc-20376780 2-Modzelewski, S., Oracz, A., Żukow, X., Iłendo, K., Śledzikowka, Z., & Waszkiewicz, N. (2024). Premenstrual syndrome: new insights into etiology and review of treatment methods. Frontiers in Psychiatry, 15. https://doi.org/10.3389/fpsyt.2024.1363875

3-Dutta, A., & Sharma, A. (2021). Prevalence of premenstrual syndrome and premenstrual dysphoric disorder in India: A systematic review and meta-analysis. Health Promotion Perspectives, 11(2), 161–170. https://doi.org/10.34172/hpp.2021.20

4-Nagarkar, A., & Kulkarni, A. S. (2024). Regional variation in prevalence of frailty in India: Evidence from longitudinal ageing study in India (LASI) wave-1. The Indian Journal of Medical Research, 159, 441–448. https://doi.org/10.25259/ijmr_865_22

5-Wikipedia contributors. (2025, May 16). Likert scale. Wikipedia. https://en.wikipedia.org/wiki/Likert_scale 6-GeeksforGeeks. (2024, May 31). Karl Pearson's Coefficient of Correlation | Methods and Examples. GeeksforGeeks. https://www.geeksforgeeks.org/karl-pearsons-coefficient-of-correlation-methods-and-examples/7-Thompson, N. (2025, April 10). What is the Spearman-Brown formula? Assessment Systems. https://assess.com/spearman-brown-prediction-formula/