

 <p>ISSN NO. 2320-5407</p>	<p>Journal Homepage: www.journalijar.com</p> <h2 style="text-align: center;">INTERNATIONAL JOURNAL OF ADVANCED RESEARCH (IJAR)</h2> <p style="text-align: center;">Article DOI:10.21474/IJAR01/21161 DOI URL: http://dx.doi.org/10.21474/IJAR01/21161</p>	
---	--	---

RESEARCH ARTICLE

CU-SIL DENTURES FOR PARTIAL EDENTULISM: PRESERVING FUNCTION AND TEETH - A CLINICAL REPORT SERIES

Harishini S., Prasad Aravind, Jailaan Murshitha, Peter S. and Shabana

Manuscript Info

Manuscript History

Received: 12 April 2025

Final Accepted: 15 May 2025

Published: June 2025

Abstract

Edentulism is a serious health issue that impairs daily activities, social interactions, and oral function in addition to degrading overall quality of life in an individual. As a result, every preventive precaution should be done to protect any teeth that may still be present, and efforts should be focused on getting a person's oral cavity back to its optimal condition. Preserving the remaining natural teeth is dentistry's top priority at the moment. A patient's psychological well-being is enhanced and the integrity of the alveolar ridge is preserved when there are a few teeth in the oral cavity. Generally, Overdentures or transitional dentures or immediate dentures following complete extractions are the treatment for the remaining natural teeth for an individual. However, Cu-Sil dentures are the most straightforward partially edentulous treatment option accessible in the transitional phase for individuals who are afraid of full extractions.

"© 2025 by the Author(s). Published by IJAR under CC BY 4.0. Unrestricted use allowed with credit to the author."

Introduction:-

MM DeVan has rightly said that "Perpetual preservation of what is remaining is more important than the meticulous reconstruction of what is lost". (1) The primary goal of contemporary dentistry is to preserve teeth in order to maintain the integrity of the alveolar ridge and the periodontium's proprioceptive capabilities. (2) Alveolar ridge height can be preserved even in cases where an arch has one healthy tooth. Studies by Va Crum and Rooney have demonstrated that, in comparison to edentulous patients, there is comparatively much less alveolar bone resorption when some teeth are present. (3) Additional benefit of maintaining natural teeth means keeping your proprioceptive sense intact of periodontium and this offers the patient psychological advantages. The remaining natural teeth are treated with overdentures, transitional dentures, or immediate dentures. However, Cu-Sil dentures are the most straightforward kind of partially edentulous treatment accessible during the transitional period for patients who are afraid of full extractions. (4) Rehabilitation of patients with Cu-Sil denture for the maxillary partially edentulous arch and a removable partial denture for the mandibular arch is covered in this clinical report.

Clinical Reports:

Case 1

A 56-year old male patient reported to our department with a chief complaint of difficulty in chewing, also complaints of poor esthetic appearance and wanted replacement of missing teeth. On intra-oral examination the patient had lost almost all the teeth in maxilla with presence of only 11 (Fig. 1) and Kennedy's class II modification 2 edentulous space in mandibular arch (Fig. 2). On examination of individual teeth 11 had gingival recession and the

mandibular teeth were supra-erupted with a full metal crown in relation to 45 and amalgam restoration in 47. A brief dental history stated that the missing teeth were extracted due to caries. It was planned to fabricate a transitional denture (CU-SIL denture) for maxillary arch and a removable partial denture in mandibular arch as the patient was not willing for extraction of his remaining teeth.



Fig. 1: -Pre-operative maxillary arch



Fig. 2:- Pre-operative mandibular arch

Treatment Procedure:

In the first appointment, diagnostic impression of maxilla and mandible were made using irreversible hydrocolloid (ZhermackTropicalgin alginate impression material). Followed by fabrication of custom tray with full coverage spacer design on the diagnostic cast on maxilla using autopolymerizing acrylic resin was done (DPI RR Cold cure, DPI India). In the second appointment, border molding using green stick compound (DPI pinnacle tracing sticks, DPI India) and secondary impression was made with light body impression material (HUGE PERFIT Elastomeric Impression Material, Light Body- Type 3). Master cast was prepared (Fig. 3) and occlusal rims fabricated on which bite registration was recorded on the third appointment.



Fig. 3:- Maxillary master cat.



Fig. 4:- Maxillary final prosthesis.



Fig. 5:- Maxillary and Mandibular final prosthesis; A- Left view, B- Front view, C- Right view.

The wax try-in was done in both maxillary and mandibular arch, evaluation of occlusion, dental esthetics, facial esthetics were examined and proceed with acrylization. The denture was finished, polished and insertion was done in maxillary and mandibular arch (Fig. 5). Denture was fabricated with opening to allow the remaining natural teeth to pass through and protrude while providing a seal around them. The patient was recalled on a periodic review (1 week, 3 weeks, 1 month, 3rd month & 6th month) which showed that the patient was comfortable in using the denture.

Case 2

A 50-year old female patient reported to our department with chief complaint of difficulty in chewing and also complaints of poor esthetic appearance and wanted replacement of her missing teeth. On intra-oral examination the patient had Kennedy's class I modification 1 edentulous space in maxilla with presence of only 13, 21 (Fig. 6) and Kennedy's class I edentulous space in mandibular arch (Fig. 7). On examination of individual teeth 13 & 21 had gingival recession. It was planned to fabricate a transitional denture (CU-SIL denture) for maxillary arch and a removable partial denture in mandibular arch as the patient was not willing for extraction of her remaining teeth.



Fig. 6:-Pre-operative maxillary arch.



Fig. 7:- Pre-operative front view of maxilla & mandible at occlusion.

Treatment Procedure:

In the first appointment, diagnostic impression of maxilla and mandible were made using irreversible hydrocolloid (Zhermack Tropicalgin alginate impression material). Followed by fabrication of custom tray with full coverage spacer design on the diagnostic cast on maxilla using autopolymerizing acrylic resin was done (DPI RR Cold cure, DPI India). In the second appointment, border molding using green stick compound (DPI pinnacle tracing sticks, DPI India) and secondary impression was made with light body impression material (Fig. 8) (HUGE PERFIT Elastomeric Impression Material, Light Body- Type 3). Master cast was prepared and occlusal rims fabricated on which bite registration was recorded on the third appointment.



Fig. 8:- Secondary impression of maxillary arch



Fig. 9:- Post-operative image with denture in maxillary and mandibular arch with CUSIL denture.

The wax try-in was done in both maxillary and mandibular arch, evaluation of occlusion, dental esthetics, facial esthetics were examined and proceed with acrylization. The denture was finished, polished and insertion was done in maxillary and mandibular arch. (Fig. 9) Denture was fabricated with opening to allow the remaining natural teeth to pass through and protrude while providing a seal around them. The patient was recalled on a periodic review (1 week, 3 weeks, 1month, 3rd month& 6th month) which showed that the patient was comfortable in using the denture.

Discussion:-

The case report describes a treatment modality for partially edentulous patient with few remaining teeth. It describes the use of an innovative technique for fabrication of transitional partial dentures based on the design for Cu-sil dentures. The vertical height of bone and proprio-sensitivity is preserved. There is no requirement of a special armamentarium and main advantage is that if a tooth is lost in future the existing denture can be modified accordingly. (2) They should be avoided in patients with bruxism, severe undercut areas and patient with high smile line. (5) soft liners are used as a reliner in cu-sil denture. Due to its viscoelastic properties, they provide cushion like effect which can distribute forces more evenly by absorbing energy. (6) Thus, the use of soft liner cushions and splints the remaining teeth which provide added stability. Moreover, it is also easy to manage a smooth transition to edentulousness. As teeth are lost, the existing prosthesis can be modified to replace them. It is thereby the simplest, gentlest way to postpone or eliminate total loss of mobile, isolated, elongated or periodontally involved teeth. (1) However, such transitional partial dentures can be used in selected cases only. Its use in patients with too many teeth and/or those exhibiting unfavourable undercuts would hinder with its fabrication and placement. Creating too many holes to accommodate natural teeth would compromise the strength of the denture, especially the mandibular one. Severe undercuts like those surrounding some natural teeth would also warrant the use of a thin denture flange and a resultant weak denture, more prone to fracture. Last, but not the least, if a patient shows a lot of their teeth and gums when smiling, and still has some natural front upper teeth, it's harder to make the denture look fully natural and esthetically perfect. Lip fullness due to the underlying flange can also result in such patients. (7)

Conclusion:-

Cu-Sil dentures offer a conservative, patient-friendly approach to managing partial edentulism, especially in individuals unwilling or unsuitable for full extractions. By preserving the remaining natural teeth, they maintain proprioceptive feedback, contribute to better alveolar ridge preservation, and provide psychological benefits to the patient. This clinical report series highlights the practical utility and adaptability of Cu-Sil dentures as an interim solution, enabling comfortable function and esthetics while delaying or avoiding complete edentulism. However, careful case selection, regular follow-up, and meticulous hygiene maintenance are vital for long-term success.

References:-

1. D. KP, Shetty M, Alva H, D. AP. PROVISIONAL RESTORATIONS IN PROSTHODONTIC REHABILITATIONS - CONCEPTS, MATERIALS AND TECHNIQUES. J Health Allied Sci NU. 2012 Jun;02(02):72–7.
2. IJAR, Arora A, Upadhyaya V, Malik D, Rohilla. K. CU-SIL DENTURE: CONSERVING THE REMAINING. Int J Adv Res. 2017 Jul 31;5(7):2086–91.
3. Crum RJ, Hospital V, Center V. Alveolar bone loss in overdentures: A S-year study. J Prosthet Dent.
4. Jain AR. Cu-sil denture for patients with few remaining teeth - A case report.
5. Jain JK, Prabhu CRA, Zahrane MA, Esawy MSA, Ajagannanavar SL, Pal KS. Cu-sil dentures – a novel approach to conserve few remaining teeth: Case reports. Case Rep.
6. Odelola HA, Koza J. Characterization of Nigerian strains of West Nile virus by plaque formation. Acta Virol. 1975 Nov;19(6):489–92.
7. Eswaran DMA. Cusil Denture -The Last Resort before Complete Denture. 2022;3(12).