

# RESEARCH ARTICLE

# EXPRESSION OF VEGF IN BREAST CANCER.

# Shamim Shera<sup>1</sup>, Syed Imtiyaz Hussain<sup>2</sup>, Shuaeb Bhat<sup>2</sup>, Sumat-ul-Khurshid<sup>3</sup>, Suhail Farooq<sup>2</sup>, Majid Ahmad Khan<sup>4</sup>, Salma Gull<sup>4</sup>, Tajali Nazir Shora<sup>4</sup> and Showkat Ahmad Mir<sup>2</sup>.

- 1. Senior Consultant, Department of Health, Government of Jammu & Kashmir, Anantnag, Jammu & Kashmir.
- 2. Assistant Professor, Department of Pathology, Government Medical College, Anantnag, Jammu & Kashmir, India.
- Assistant Professor, Department of Pathology, Government Medical College, Doda, Jammu & Kashmir, India. 3.
- Senior Resident, Department of Pathology, Government Medical College, Anantnag, Jammu & Kashmir, India. 4.

.....

# Manuscript Info

#### Abstract

• • • • • • • • • • • • • • • • • • • •	•••••••••••••••••••••••••••••••••••••••		
Manuscript History	Aim: Breast Cancer is the most frequent cancer in India. Tumor		
Received: 14 August 2019	angiogenesis has been considered as a crucial step in the cancer		
Final Accepted: 16 September 2019	development and progression Vascular endothelial growth factor		
Published: October 2019	(VEGF) is one important prognostic marker in patients with breast		
Vanwarda	cancer .The aim of the study is to see the expression of VEGF in breast		
Key words:- VEGF, Immunohistochemistry.	cancers by immunohistochemistry.		
	Materials and methods: Formalin fixed paraffin embedded sections of		
	100 cases of malignant breast lesions were taken up for the study and		
	subjected to immunohistochemistry using VEGF. <b>Results:</b> The intensity of VEGF immunostaining in malignant breast		
	lesions was evaluated and scoring was graded as $0,1+,2+,3+$ and $4+$ .		
	Statistical analysis was performed with Chi-Square test and significant		
	differences were noted between these 3 groups (n value $< 0.05$ )		
	<b>Conclusion:</b> VEGE expression correlated well with the grade and		
	stages of types indicating that VECE positive types are biologically		
	stages of tumor indicating that VEOF positive tumors are biologically		
	aggressive and are associated with poor prognosis.		
	Copy Right, IJAR, 2019,. All rights reserved.		
Published: October 2019 <i>Key words:-</i> VEGF , Immunohistochemistry.	<ul> <li>(VEGF) is one important prognession. Vascular endothenar growth factor (VEGF) is one important prognostic marker in patients with breas cancer .The aim of the study is to see the expression of VEGF in breas cancers by immunohistochemistry.</li> <li>Materials and methods: Formalin fixed paraffin embedded sections on 100 cases of malignant breast lesions were taken up for the study and subjected to immunohistochemistry using VEGF.</li> <li>Results: The intensity of VEGF immunostaining in malignant breas lesions was evaluated and scoring was graded as 0,1+,2+,3+ and 4+ Statistical analysis was performed with Chi-Square test and significant differences were noted between these 3 groups (p value&lt; 0.05).</li> <li>Conclusion: VEGF expression correlated well with the grade and stages of tumor indicating that VEGF positive tumors are biologically aggressive and are associated with poor prognosis.</li> </ul>		

Introduction:-

Breast cancer is the most frequent cancer in India and mortality rates associated with it is higher in India. Molecular mechanism are involved in its pathogenesis with many genetic alterations and oncogene protein products having a role to play which interfere with the mechanism of proliferation and differentiation of tumor growth. The prognostic biomarkers include raised levels of expression of Proliferation markers like Ki67, expression of Estrogen Receptor (ER) and Progesterone Receptor (PR), amplification and overexpression of HER2, cyclin D1 etc.

Recent studies have found tumor angiogenesis as a critical step in cancer development and progression. Among these, vascular endothelial growth factor (VEGF) has emegred as a prognostic marker with several type of cancer including breast cancer.<sup>1</sup>

VEGF is produced and secreted by a number of normal cells and is a poly-functional molecule that has been implicated in vasculogenesis, endothelial cell proliferation and migration, vascular permeability and stromal degradation by the activation of some proteolytic enzymes involved in tumor invasiveness and angiogenesis.<sup>2</sup>VEGF

Corresponding Author:- Dr. Showkat Ahmad Mir. Address: - Department of Pathology, Government Medical College, Anantnag, Jammu & Kashmir, India.

is required for the initial stages of breast cancer tumorogenesis, and this initial effect is related to the formation of neovascular stroma. Keeping all these alterations in mind, in this present study we have studied the VEGF expression in malignant breast lesions.

# **Materials and Methods:-**

The study was conducted at a Tertiary Care Hospital with 100 cases of malignant breast lesions taken up for study. All female patients irrespective of their age and other physical conditions during the period of June 2010 to January 2017 were taken up for the study. Females of age ranging from 20 to 70 years were taken up for the study and all of them underwent modified radical mastectomy (MRM). The diagnosis was reconfirmed on Hematoxylin and Eosin (H&E) stained sections and the appropriate blocks were subjected to IHC using VEGF antibody.

# Scoring system:

Positive immunoreaction showed a dark brown precipitate (cytoplasmic for VEGF). The intensity of the staining was assessed by a scoring system laid by Sophia K. et al<sup>2</sup>.

Score	Results	Interpretation			
0	Negative	None or <5% cells positive			
1+	Weak or Mild staining	Weak or mild staining; 5-10% of tumor cells are positive			
2+	Moderate staining	<25% of tumor cells are positive			
3+	Strong staining	Strong staining; 25-50% of tumor cells are positive			
4+	Highly strong staining	Highly strong staining; >50% of tumor cells are positive			
Table 1. Searing for VECE					

## Scoring according to Sophia K. et al, at objective 40x: (Table 1)

Table 1:-Scoring for VEGF

#### Statistical analyses

Statistical analyses of all results were done by using Chi square test at level of significance  $p \le 0.05$  was done.

## **Results:-**

The malignant lesions in the study were seen in all histological grades of the tumor with majority of the cases having a higher grade. These also presented with different stages with predominant cases having a higher stage with a presentation of Stage III being more common followed by Stage IV. None of the cases presented in Stage 0 or Stage I in our study. The axillary lymph nodes were involved in 70% of our cases (Table 2).

	Grade			Stage					Axillary Lymph Nodes	
Features	Ι	II	III	0	Ι	II	III	IV	Positive	Negative
No. of cases	10	40	50	0	0	20	60	20	70	30

**Table 2:-**The characteristic features of all malignant breast lesions

Out of the 100 cases of malignant breast lesions, expression of VEGF was noted 75% of cases in the current study (Figure 1).

#### Intensity of staining of VEGF in malignant breast lesions:

Among the 75 cases, 25 cases showed a weak (1+) positivity for VEGF wheras 35 cases showed a moderate positivity (2+), 15 cases showed a strong (3+) positivity and none of the cases showed a highly strong (4+) positivity (p = <0.05) (Table 3).

Intensity of staining of VEGF	No. of cases	Percentage
1+	25	33
2+	35	47
3+	15	20
4+	0	0



Table3:-Percentage Of Staining Intensity Expression In Malignant Vegf Breast Lesions.

**Figure 1:-**VEGF in IDC Breast. a) IDC breast with a large firm grey white area. b) H & E of IDC breast. (10x) c & d) Positive Immunostaining of VEGF in IDC breast (10x).

#### **VEGF** overexpression and grade of tumor

VEGF immunohistochemical analysis in relation to grade of tumor revealed that none of grade I was positive, 30 (75%) out of 40 cases of grade II were positive, 45 out of 50 (90%) cases of grade III were positive for VEGF. The detection rate of VEGF correlated well with the grade of tumor with higher grade tumor showing positivity in a higher proportion whereas the tumor with lower grade I, VEGF expression was not detected p<0.05 (Table 4).

Grade of Tumor	Positive Staining	Negative Staining	Total
Ι	0	10	10
П	30	10	40
III	45	5	50

Table 4:-Expression of VEGF in relation to grade of the tumor

#### **VEGF** Overexpression and stage of tumor

VEGF immunohistochemical expression was reported in all 20 cases of stage IV with 45 out of 60 cases in stage III showing positivity and among the 20 cases in stage II, 10 cases were positive for VEGF. There was significant positive correlation between VEGF overexpression and the stage of tumor (p value <0.05), and a higher proportion of cases were found in stage III and IV (Table 5).

Stage of Tumor	Positive Staining	Negative Staining	Total
Ι	0	0	0
II	10	10	20
III	45	15	60
IV	20	0	20

Table 5:-Expression of VEGF in relation to the stage of the tumor

#### VEGF immunohistochemical expression and axillary lymph node involvement

65 out of 70 cases of node-positive breast cancer found to have VEGF overexpression (92%) while only ten out of 30 cases of node-negative breast cancer showed VEGF overexpression (33%) with significant difference between these two groups. (p value < 0.05)

## **Discussion:-**

#### Expression of VEGF in malignant breast lesions

Overexpression of VEGF in infiltrating ductal carcinoma was found in 75% of cases. This finding is higher than those of Melanie Schmidt *et al* (2008) who reported that 60% were VEGF positive<sup>9</sup>. It was higher than a study conducted by ES Al Harris *et al* who reported. 61.5% were VEGF positive<sup>4</sup>. It was lower than those of Anca Maria Cimpean *et al* (2008) who reported that 87.1% of primary breast carcinoma was VEGF positive<sup>5</sup>. In this study VEGF overexpression was detected in 75% of breast cancer patients.

#### VEGF overexpression and grade of tumor

VEGF immunohistochemical analysis in relation to grade of tumor revealed that none of grade I was positive, 75% of grade II were positive, 90% of grade III were positive for VEGF. There was highly significant positive correlation between VEGF overexpression and grade of breast cancer (p<0.05). This finding correlated well with studies conducted by Linderholm B *et al.*,(2000), Gottfried *et al.*,(2004), Shankar R *et al.*,2006) and AL-Harris E *et al*(2007)<sup>4,6-7</sup>

#### **VEGF** overexpression and stage of tumor

VEGF immunohistochemical expression was reported in 10 out of 20 cases of stage II, in 45 out of 60 cases of stage III, and in all 20 cases of stage IV. There was significant positive correlation between VEGF overexpression and the stage of tumor (p value <0.05), and a higher proportion of cases were found in stage III and IV. These findings correlated well with Collagy G *et al.*, (2000), Bolat F *et al.*,(2006), AL-Harris E *et al*(2007) and Xu W *et al.*, (2007).<sup>4,9-11</sup>

VEGF is expressed more in those with advanced stage which reflects the aggressive behavior of the tumor.

#### VEGF immunohistochemical expression and axillary lymph node involvement

VEGF overexpression is higher in node positive breast cancer than in node negative breast cancer with significant difference between these two groups (p value <0.05).

This finding agreed with that reported by Yi WJ et al., (2003), Gottfried et al., (2004), Wang X et al., (2006)<sup>8, 1 2-13</sup>

#### **Conclusion:-**

The pathogenesis of breast cancer is a multi-stage process involving progressive accumulation of genetic alterations. In this study, VEGF overexpression was significant in all grades and stages of breast cancer (p<0.05). VEGF overexpression correlated well with the grade and stage of tumor indicating that VEGF positive tumors are biologically aggressive and are associated with poor prognosis.

#### **References:-**

- 1. Giampietro Gasparini Emanuela Bonoldi Carlo Gatti Orazio Vinante Masakazu ToiTakeshi Tominaga Massimo Gion Ruggero Dittadi Paolo Verderio Patrizia BoracchiMitsuya Hanatani Isamu Matsubara Hideo Suzuki. Prognostic significance of vascular endothelial growth factor protein in node-negative breast carcinoma. Journal of the National Cancer Institute 1997; 89: 39-47.
- 2. Sophia K, Apple J, Randolph H, et al. Immunohistochemical evaluation of K-ras, p53 and HER-2/neu expression in hyperplastic, dysplastic and carcinomatous lesions of the pancreas: evidence for multistep carcinogenesis. Human Pathology 1999;30:123-130.
- 3. Melanie Schmidt *et al*(2008), Expression of VEGF-R1 in Breast cancer is associated with VEGF expression and Node negative tumor stage, Anticancer Research 28:1719-1724.
- 4. AL-Harris ES. Overexpression of VEGF in Correlation to Ki-67 grade and stage of breast cancer. Saudi Medical Journal 2008.Vol.29(8):page 477-482.
- 5. Anca Maria Cimpean *et al* (2008), Correlation of VEGF with recurrences, survival, and first relapse site in primary breast cancer after adjuvant treatment. J Of Clin Oncol 2008;18(7):1423-1431.
- 6. Linderholm B, Lindg B and Tavelin B. p53 and vascular endothelial growth factor (VEGF) expression predicts outcome in 833 patients with primary breast carcinoma. Int J Cancer2000;89:51-62.

- 7. Shankar RS, Tiwary K, Khanna R, et al. Tumor Angiogenesis: Determined By VEGF Expression, MAGS Scoring, Doppler Study, In Carcinoma Breast. The Int J Surgery. 2006; 8:No.1.
- 8. Gottfried E. Konecny, Glosia YM, Michael U, et al. Association between HER-2/neu and VEGF expression predicts clinical outcome in primary breast cancer patients. Clin Can Res 2004;10:1706-1716.
- 9. Collagy G, Dimitriadis E, Harmey J, et al. Immunohistochemical measurement of tumor VEGF in breast cancer ,a more reliable predictor of tumor stage than Microvessel density or serum VEGF. Appl Immunohistochemical Mol Morphol 2000 Jun;8(2):104-113.
- 10. Bolat F, Kayaselcuk F, Nursal TZ, et al. Microvessel density, VEGF expression, and tumor associated macrophages in breast tumors. J Exp Clin Cancer Res 2006 Sep ; 25(3):365-437.
- 11. Xu W, Wang G, Zou Y, et al. VEGF expression in IDC of breast. Chinese J Of Cancer Res 2007;19(1):56-59.
- 12. Yi WJ, Tang ZH, Yang ZL, et al. Difference in expression of VEGF, bFGF and their receptors between the young and postmenopausal women with breast cancer. Zhongua Zhong Liu Za Zhi 2003 March;25(2):141-145.
- 13. Wang XB, Yang QX and Pei XJ. Expression of angiogenesis related factors in invasive breast cancer and its clinical significance. Nan Fang Yi Ke Da Xue Xue Bao.2006 Jun;26(6):860-863.