

Journal homepage: http://www.journalijar.com

INTERNATIONAL JOURNAL OF ADVANCED RESEARCH

RESEARCH ARTICLE

Bidirectional double muscle transfer to restore shoulder abduction and external rotation in late obstetric brachial plexus injuries

MOHAMMAD-REDA AHMAD M.D

The Department of general surgery, Plastic and reconstructive surgery unit, faculty of medicine, Zagazig University

Manuscript Info	Abstract
Manuscript History:	Background: Medial rotation contracture represents the most persistent and
Received: 11 January 2015 Final Accepted: 23 February 2015 Published Online: March 2015	frequent late deformity of the shoulder, in obstetric brachial plexus injuries (OBPIs). Many techniques have been tried to solve this problem.Aim: This study aimsto enhance shoulder abduction and outward rotation by performing an ordinary single direction muscle transfer (SDMT) or
Key words:	bidirectional double muscle transfer (BDMT) around the shoulder. Methods: 40 patients: 21 male and 19 female, their age ranged from 2 to10
Bidirectional, Brachial plexus, External rotation, Shoulder	years complaining of weak abductors and external rotation of the shoulder due to OBPIs. They were divided into two groups according to the
*Corresponding Author	(23 patients) operated with SDMT.
MOHAMMAD-REDA AHMAD M.D	 Results: There was significant postoperative improvement in shoulder abduction, external rotation and flexion in both procedure "BDMT and SDMT". In addition, the young age gives better results as there are negative correlations between patient age and degree of shoulder functions improvement in both groups.Moreover, BDMT procedure had significant higher degree of improvement in shoulder abduction and external rotation than SDMT procedure. Conclusion: BDMT is recommended as an optimal option to improve both abduction and external rotation of shoulder in late OBPIs.
	Copy Right, IJAR, 2015,. All rights reserved

.....

INTRODUCTION

Obstetric brachial plexus injuries (OBPIs) is a mixed nerve and musculoskeletal complication that happens during the process of difficult delivery ^[1]. A considerable percent of cases tends to acquirelifelong limb deficits^[2]. The fast limb growth with marked asymmetry in the pathology of nerve involvement within the brachial plexus, may lead to a lot of sequalae as muscle contracture, weakness, and imbalances. These deformities become more evident with failure of growth of the denervated muscles, or at least developed at a lower rate than the antagonist innervated muscles. Teathering effect may develop gradually, that limits the passive range of motion around the shoulder joint. And muscle imbalances and contractures can lead to a lot of bony deformities ^[3]. In addition, bony deformities can also result from decreased innervation from the brachial plexus to shoulder joint bones ^[4-5].

All the previous prerequisites can lead to progressive and persistent glenohumeral (GH) joint dysplasia and shoulder joint sublaxation and dislocation ^[6].

Muscle balance between agonists and antagonists around the shoulder joint is the main aim of the treatment. This aim can be achieved firstly by release of the teathering contractures of the internal rotators. Many techniques had been published for this release, as posterior subscapularis muscle slide, or through anterior approach lengthening of the tendon of subscapularis. Through this anterior approach the tendon of pectoralis major can be also lengthen, and any ligamentous or capsular contractures can be dealt with ^[7-8].

If the active external rotation and abduction were not recovered well, muscle transfer can be done in supple and congruent GH joint. If there is severe glenohumeral dysplasia with internal rotation contraction especially in older children, then external humeral osteotomy is preferred ^{[9-10].}

In this research, the bidirectional double muscle transfer is proposed to improve shoulder functions after late OPPIs and compared with the single muscle transfer.

PATIENTS AND METHODS

Patient data analysis:

From January 2011 to June 2014 forty patients {21 male (52.5%) 19 female (47.5%)} were operated. They were divided into two groups according to the performed operation (table 1). Ethical committee permission was granted for this study(approved by the Institutional Review Board and ethics committee of faculty of medicine- Zagazig University) and written consent of patients was also taken.

Group 1 (17 patients) operated with bidirectional double muscle transfer (BDMT): their age ranged from 2 to 10 years mean \pm SD (4.18 \pm 2.24). Affected side was the right side in 10 cases (58.82 %) and left side in 7 (41.18%). The affected roots were C5-6 in 1 case (5.89 %), C5-6-7 in 11 cases (64.70 %) and total in 5 cases (29.41%). The follow up period: ranged from 6 to 36 months mean \pm SD (18.76 \pm 7.35). Before operation, degree of shoulder abduction ranged from 30 to 75 degrees, shoulder external rotation ranged from zero to 20 and shoulder flexion ranged from 0-30.

Group 2 (23 patients) operated with Single muscle transfer (SDMT): their age ranged from 2 to 9 years mean \pm SD (4.48 \pm 1.93). Affected side was the right side in 8 cases (34.78 %) and left side in 15 (65.22%). The affected roots were C5-6 in 3 cases (13.04 %) and C5-6-7 in 20 cases (86.96 %). The follow up period: ranged from 6 to 38 months mean \pm SD (19.74 \pm 9.25). Before operation, degree of shoulder abduction ranged from 40 to 90 degrees, shoulder external rotation ranged from zero to 30 and shoulder flexion ranged from 10-25.

In this study, two methods of transfer around the shoulder joint were done. **Operative technique of the single direction muscle transfer (SDMT)**:

In this technique I used a procedure similar to that of Gilbert^[11] and Ahmad^[12].

Patient after being anaesthetized, was lied in lateral decubitus while the affected side being up. At the start of this surgery, assessment of the passive range of movement (PROM) around the shoulder joint is so essential. Any contractures, hypertrophy of the internal rotators is to be considered. I used posterior approach at the lateral edge of the scapula in a zigzag incision. Through this approach, posterior subscapularis muscle slide and release was achieved. Then after, the conjoint tendon was dissected from surrounding structures, and the part related to latissimus dorsi (LD) muscle in the conjoint tendon was separated from that of terries major muscle (tmm). Dissection and mobilization of the LD muscle to enable successful transfer to a new insertion site was done. In all previous steps (posterior subscapularis muscle release, conjoint tendon dissection, and mobilization of the upper part of LD muscle), the pedicles of the LD muscle, and tmm, should be saved and protected. After that, detaching the tendon of LD muscle from the medial side of the humerus with great protection and care to neurovascular structures of the upper arm was done. After that, the arm adjusted in 90° abduction, 90° external rotation, then the posterior fibers of the deltoid was elevated to make a tunnel for new insertion of LD tendon. At this step, care must be experienced to avoid injury of the fibers of the axillary nerve. I used heavy polyprolene suture to fix the tendon of LD into the point at the greater tuberosity of the humerus at the site of insertion of the infraspinatous. Little sutures was taken between infraspiatous muscle and transferred LD muscle belly, to make some synergism in function and direction of pull. Closure of the wound, (while the arm was in 90° abduction, 90° external rotation, and the elbow at 90° flexion) was achieved. In the same position, shoulder spica cast was applied for 6 weeks. After removal of the cast airplane custom made splint was applied with gradual weaning for 3 months, after that physiotherapy was scheduled.

Operative technique of the bidirectional double muscle transfer (BDMT):

In this group, I used the same position, incision, and subscapularis release of previous SDMT. After release of the part of LD muscle of the conjoint tendon from the tmm part (Figure 1A and 1B).

Statistical analysis:

Data were presented as mean \pm standard deviation (SD). Statistical analysis was performed using the 20 version of SPSS statistical software for windows (SPSS Inc. Chicago, IL, USA). Paired student t test used in the comparison between preoperative and postoperative evaluation and unpaired student t test used in the comparison between 2

groups. Person correlation used to relate patient's age and postoperative improvement. P values less than 0.05 were considered significant

RESULTS

Table 2 shows a comparison between shoulder abduction improvements after the two operative procedures. There was significant postoperative shoulder abduction improvement in both procedure "BDMT and SDMT" when compared with the degree of abduction before operation (P = 0.000). In addition there is negative correlation between patient age and degree of shoulder abduction improvement in both groups (r = -572 and -575, P = 0.016 and 0.004 respectively). Moreover, BDMT procedure had significant higher degree of improvement in shoulder abduction than SDMT procedure (P = 0.000) with percentage of change 149.45% in comparison to 80.48% in SDMT.

Table 3 shows a comparison between shoulder external rotation improvements after the two operative procedures. There was significant postoperative shoulder external rotation improvement in both procedure "BDMT and SDMT" when compared with the degree of external rotation before operation (P = 0.000). In addition there is negative correlation between patient age and degree of shoulder external rotation improvement in both groups (r = -720 and -578, P = 0.001 and 0.004 respectively). Furthermore, BDMT procedure had significant higher degree of improvement in shoulder external rotation than SDMT procedure (P = 0.009) with percentage of change 1285.54% in comparison to 1095.89% in SDMT.

Table 4 shows a comparison between shoulder flexion improvements after the two operative procedures. There was significant postoperative shoulder flexion improvement in both procedure "BDMT and SDMT" when compared with the degree of flexion before operation (P = 0.000). In addition there is negative correlation between patient age and degree of shoulder flexion improvement in both groups (r = -523 and -448, P = 0.031 and 0.032 respectively). However, there was non-significant statistical difference between the degree of improvement in shoulder flexion in both procedures (P = 0.061) with percentage of change 602.64% in comparison to 470.73% in SDMT.



Figure 1A: diagram for the dissection of the tendons of latissimus dorsi and terris major muscles and their pedicles.

Figure 1 B: the dissection of the tendons of latissimus dorsi and terris major muscles and their pedicles

The tendon of tmm was fixed to the inferior surface of the head of the humerus below the site of lesser tuberosity in full abduction. Care to avoid injury of the neurovascular bundle of the upper limb is essential. After that, the tendon of the LD muscle was fixed like the previous transfer to the upper most part of the greater tuberosity of the humerus, while the arm was at 90° abduction and 90° external rotation. I used the same fixation position, period of cast fixation, same proid of custom made brace, and the same physiotherapy schedule likeSDMT(Figure 2A and 2B).



Figure 2 A: Diagram for the direction and site of insertion of latissmus dorsi and terries major muscles.

Figure 2 B: the direction and site of insertion of latissmus dorsi and terries major muscles.

Table 2:A comparison between shoulder abduction improvements in the two procedures.

Shoulder Abduction	Group 1: H	BDMT (n=17)	Group 2: Sl	DMT (n= 23)
	before	After	before	After
range	30-75	95-160	40-90	90-130
Mean	51.76	129.12	60.22	108.69
SD	11.45	20.33	12.83	11.10
% of change		149.45 %		80.48 %
P value of tt	0	.000	0.0	000
P value VS BDMT				0.000
r (with age)		572*		575**
P value		0.016		0.004

Shoulder External	Group 1:	BDMT (n=17)	Group 2: SDMT (n= 23)				
rotation	before	After	before	After			
range	0-20	65-90	0-30	60-90			
Mean	5.88	81.47	6.09	72.83			
SD	8.70	8.43	9.88	10.75			
% of change		1285.54		1095.89			
P value of tt		0.000	0.000				
P value VS BDMT				0.009			
r (with age)		720 **		578 **			
P value		0.001		0.004			

Table 3: A comparison between shoulder external rotationimprovements in the two procedures.

Table 4: A comparison between shoulder flexion improvements in the two procedures.

Shoulder flexion	Group 1: B	SDMT (n=17)	Group 2:	: SDMT (n= 23)		
	before	After	before	After		
Range	0-30	50-90	10-25	60-90		
Mean	10.59	74.41	11.96	68.26		
SD	7.48	11.16	5.38	8.99		
% of change		602.64%		470.73%		
P value of tt	0.	000		0.000		
P VS BDMT				0.061 (NS)		
r (with age)		523*		448*		
P value		0.031		0.032		



Figure 3: A case of bidirectional muscle transfer; 3A preoperative,3B postoperative abduction and 3C postoperative external rotation.



4A 4B 4B 4B 4B 4B 4B

Figure 4: A case of bidirectional muscle transfer; 4A: preoperative, 4B: X ray showing hypoplasia of the humeral head with deformities of the glenoid cavity. 4C: postoperative abduction and 4D: postoperative external rotation.



Figure 5: A case of single direction muscle transfer; 5A preoperative,5B postoperative abduction and 5C postoperative external rotation.

Table 1: Patient data of bidirectional double muscle transfer and single direction muscle transfer in late
OBPIs(BD = bidirectional double . S = single. F U/ M = follow-up per months. Pre. Op.= previous
operation. Ab = shoulder abduction. ER= shoulder external rotation. FL = shoulder flexion.

Ν	A	S	Pre.	Sid	F	Preoperative			6 month post			12 n	nonth p	ost	≥ 18	Affec		
	ge (y	e x	Op.	e	0/ M	А	Е	F	Ab	E	FL	Ab	ER	F	Ab	E E	F	root
)					b	R	L		R				L		R	L	
1B	2	m	No	Rt	12	4 0	0	10	150	80	70	150	95	70				C5-6- 7
2B	2	f	1ry ex	Rt	24	5 0	10	20	150	85	90	145	90	65	145	90	65	C5-6- 7
3B	2	m	no	Rt	24	7	0	10	110	85	70	150	90	60	150	85	60	C5-6- 7
4B	2	m	no	Rt	12	5 0	20	10	150	90	100	150	80	75				C5-6
5B	2	m	1ry ex	Rt	18	5 0	20	0	160	90	90	160	60	40	160	60	40	Total
6B	4	m	no	Rt	18	3 0	0	0	150	90	75	90	75	60	150	85	65	Total
7B	3	m	No	Lt	24	6 0	0	20	140	90	80	130	85	80	140	85	80	C5-6- 7
8B	3	f	no	Lt	36	4 0	0	10	140	85	80	140	90	65	140	90	65	C5-6- 7
9B	4	m	no	Lt	30	5 0	0	10	120	70	75	120	65	65	120	65	65	Total
10B	4	m	no	Lt	12	5 0	20	0	120	75	75	110	55	50				Total
11B	6	m	No	Rt	22	4 0	0	10	120	85	80	100	55	55	100	55	55	Total
12B	8	f	no	Rt	17	6 0	10	30	140	90	90	95	55	60				C5-6- 7
13B	4	f	No	Lt	6	5 0	0	10	130	85	80							C5-6- 7
14B	5	f	no	Lt	16	6 0	20	10	120	80	70	100	65	60				C5-6- 7
15B	5	f	no	Rt	13	6 0	0	10	100	70	60	100	60	60				C5-6- 7
16B	5	f	No	Lt	18	7 0	0	10	100	70	55	100	70	55	100	65	55	C5-6- 7
17B	10	М	No	Rt	17	4 5	0	10	95	65	50	90	65	50				C5-6- 7
1 S	2	m	no	Lt	12	6 0	0	10	120	80	60	160	90	65				C5-6- 7
28	2	f	no	Lt	14	7 0	20	10	110	80	70	140	90	70				C5-6- 7
3S	2. 5	f	no	Lt	38	5 0	0	10	100	75	80	150	90	70	145	90	70	C5-6- 7
4S	2. 5	f	no	Lt	12	4 0	0	20	100	90	90	150	85	70				C5-6- 7
55	3	f	1ry ex	Lt	6	6 0	0	10	110	85	70	140	85	60	140	85	60	C5-6- 7
6S	2	m	no	Rt	36	6 0	20	10	90	75	60	120	70	60	120	65	60	C5-6- 7 227
7S	3	m	no	Rt	18	9	20	25	120	90	80	145	90	70	145	90	70	C5-6-

						Δ												7
8S	4	m	no	Lt	16	6	0	10	90	75	65	125	80	65				/ C5-6-
						0												7
9S	6	m	no	Lt	24	4	10	10	100	80	75	130	80	60	130	80	60	C5-6-
100				.	•	0	•	10	100	0.0		105		60	105	0.0		7
108	4	t	no	Lt	28	0	20	10	120	80	80	125	80	60	125	80	60	C5-6- 7
11 S	4	f	No	Rt	11	6	0	10	120	80	70	120	70	60				C5-6-
						0												7
12S	4	m	No	Lt	20	7	0	10	130	85	80	120	75	65	120	75	56	C5-6-
120	4			T.	1.4	0	20	10	105	00	75	120	70	(0)				/
155	4	m	по	Lt	14	8 0	30	10	125	80	75	130	70	00				C5-6- 7
14S	4.	f	no	Lt	10	8	0	10	120	65	65	120	65	60				C5-6-
	5					0	_											7
15S	9	f	No	Rt	22	6 0	0	20	110	65	65	110	70	70	110	70	70	C5-6- 7
16S	5	f	No	Rt	36	5	0	20	110	70	60	110	70	65	110	70	65	C5-6-
						0												7
17S	8	m	No	Rt	36	6 0	0	20	115	60	60	110	60	60	115	60	60	C5-6
18S	5	m	no	Lt	12	6	20	10	100	65	60	100	65	60				C5-6-
						0												7
19S	8	f	No	Rt	19	5 0	0	10	100	60	60	100	60	60	100	60	60	C5-6
20S	5	f	no	Lt	18	4	0	10	100	60	65	100	60	60	100	60	60	C5-6-
						5												7
21S	5	f	No	Lt	18	5	0	10	100	60	60	100	60	65	100	60	65	C5-6
220	~			D	1.4	0	0		100	60	(0)	100	65	60				05.6
228	5	m	no	Rt	14	5 0	0	0	100	60	60	100	65	60				C5-6- 7
23S	5.	m	no	Lt	20	7	0	10	110	55	60	100	60	60	100	60	60	C5-6-
	5					0												7

DISCUSSION

Progressive and lifelong deformities of the shoulder represent the most common complications of OBPIs are ^[13-15].

Many authors considered muscle release procedure is indicated in internal rotation contracture before the age of two years, and the muscle transfer to achieve good external rotation and abduction in older children ^[16].

It is logic to use the procedure of muscle release and transfer before evident dysplasia of GH joint that usually happened at age of 3 or 4 year^{[17].}

Gilbert et al. ^[18] and many authors suggested that the release of the internal rotation contracture is indicated if passive external rotation is less than 20°. I used the same concept in this work, especially while assessing the PROM of the shoulder joint at the start of the surgery.

With the same concept, **Raimondiet al.** ^[19]were doing the release of subscapularis, and then they waited for the recovery of active external rotators. They preferred to combine both release and transfer in the surgery in children more than 4 year old.

Another concept presented by**Muhlig et al**^[16]is remarkable. They described a policy for management of internal rotation contracture deformity. They suggested to do muscle release when passive external rotation is less than 30°. And in cases without shoulder joint sublaxation or dislocation, they were doing posterior approach muscle slide operation. In other situation, when there was dislocation or sublaxation of the shoulder, then anterior subscapular tendon lengthening was preferred. They added that, in cases showed no recovery of infraspinatous muscle at the age of 2 years old, then LD muscle transfer was added to the release. I did muscle transfer and release in all cases at the same time, to avoid reoperation with suspected fibrosis and adhesions in the field if the rotator cuff muscles were

not regained function after the release surgery. The other reason is to avoid internal rotation contracture recurrence. Also is to provide the cases with rapid, sufficient, and long lasting improvement. Lastly is to limit the costs of 2 surgeries in a country with financial problems.

Hoffer et al. ^[20] used tmm in the technique to improve active outward rotation of the shoulder. They listed the indications for this when the flexors and abductors of the shoulder are weak against gravity, and when tmm shows cocontraction during active abduction of the shoulder. The third indication is to augement the muscle volume of the cranial part of trapezius muscle transfer.

Bahm and Ocampo-Pavez^[21] supposed that the monopolar transfer of the tmm around its vascular pedicle by keeping the insertion of the muscle intact is a perfect transfer to improve shoulder abduction, especially if there is tmm cocontraction during active shoulder abduction.

In group I, I used BDMT by transferring the tendon of tmm to a site proposed to aid in abduction and stabilization of the GH joint. From my opinion my proposed transfer is easier, feasible, and safe for the pedicles of the LD muscle, and tmm muscle. The significant improvement in abduction after BDMT in comparison to SDMT makes me to say that this technique is a safe, easy, and effective as regard achieving perfect abduction and external rotation function restoration.

Gilbert ^[11]after his work on 44 cases stated that LD muscle transfer is sufficient for shoulder functions restoration (abduction and external rotation), if the active abduction is more than 90° . In other words, if active external rotation was between 45° - 90° the LD muscle transfer was not sufficient. So he advised to add trapezius transfer for shoulder in this situation.

Chen et al ^[22] agreed with previous idea, and they added a modification in the transfer. After insertion of the tendon of LD muscle to the insertion of the infraspinatous muscle, they did tenotomy of the tendon of tmm and reattached it to the muscle belly of the LD muscle in its new position to augment its function.

Ahmad et al.^[23]asserted that the concomitant transfer of trapezius muscle with LD muscle transfer given marked improvement in both shoulder abduction and external rotation, rather than LD muscle transfer only.

Al-Qattan^[24] has a different concept. He argued that there is no differences in the results of LD muscle transfer in correlation with preoperative abduction degrees after his work on 12 cases with deficient shoulder abduction and external rotation.

The derotation osteotomy of the humerus is a traditional method to improve cases with late resistant internal rotation contracture, But this procedure doesn't care about the GH joint dysplasia or shoulder joint dynamics, rather than its high failure rate^[25-26].

Another surgical idea by opening the GH joint anterior capsule, then tilting the humeral head laterally within the deformed glenoid cavity was done by some surgeons. But this surgical technique carries the risk of severe external rotation contracture, which is might be disabling more than internal rotation contracture itself^[3, 27, 28].

Finally, Bidirectional double muscle transfer procedure had significant higher degree of improvement in shoulder abduction and external rotation than single muscle transfer procedure. The young age gives better results. Therefore, this technique is recommended as an optimal procedure to improve both abduction and external rotation of shoulder joint.

REFERENCES

- 1- Pollack RN, Buchman AS, Yaffe H, Divon MY (2000) Obstetrical brachial palsy: pathogenesis, risk factors, and prevention. *Clin Obstet Gynecol.*, 43:236–246.
- 2- Hoeksma AF, Ter Steeg AM, Dijkstra P, Nelissen RG, Beelen A, de Jong BA (2003) Shoulder contracture and osseous deformity in obstetrical brachial plexus injuries. J Bone Joint Surg (Am), 85(A):316–322.
- 3- Nath RK (2007) Obstetric brachial plexus injuries-Erb's palsy. In Nath RK. Ed. the *Nath method of diagnosis* and treatment. Virtual Bookworm.com Publishing, College Station, 79.
- **4-** Garcia-Castellano JM, Diaz-Herrera P, Morcuende JA (2000)Is bone a target-tissue for the nervous system? New advances on the understanding of their interactions. Iowa *Orthop J*. 20:49–58.

5- Terzis JK, Kokkalis ZT (2010) Bone discrepancy as a powerful indicator for early surgery in obstetric brachial plexus palsy. *Hand* (NY) 5: 386–396.

- 6- Birch R, Bonney G, Wynn Parry CB (1998) Birth lesions of the brachial plexus. In: Birch R, Bonney G, Wynn Parry CB eds. Surgical disorders of the peripheral nerves. Churchill Livingstone, New York, 209–233.
- 7- Chuang DC, Ma HS, Wei FC. A (1998) new strategy of muscle transposition for treatment of shoulder deformity caused by obstetric brachial plexus palsy. *Plast Reconstr Surg.* 101:686-694.
- 8- Waters PM, Allan E, Peljovich E (1999) Shoulder reconstruction in patients with chronic brachial plexus birth palsy. *Clin Orthop.* 364:144-152.

- **9- Birch R (2001)** Medial rotation contracture, posterior dislocation of the shoulder. In: Gilbert A. ed. *Brachial plexus injuries*. Hampshire (UK), Taylor & Francis. 249-259.
- Al-Qattan MM (2003)Classification of secondary shoulder deformities in obstetric brachial plexus palsy. J Hand Surg [Br] 28:483-486.
- **11- Gilbert A** (**1993**)Obstetrical brachial plexus palsy. In: Tubiana R. Ed. *The Hand*. Philadelphia, WB Saunders. 4: 575–601.
- 12- Ahmad MR (2011)Operations around shoulder. In: Ahmad MR. ed. *Management of late obstetric brachial plexus*. IAP Lambert academic publishing, Beau Bassin, Mauritius 80-81.
- 13- Nath RK, Humphries AD (2009) Computed tomography of the shoulders in patients with obstetric brachial plexus injuries: a retrospective study. *Ann Surg Innov Res.* 2008; 2:4-9.
- 14- Nath RK, Amrani A, Melcher SE, Eichhorn MG (2009) Triangle tilt surgery in an older pediatric patient with obstetric brachial plexus injury. *ePlasty*9:e26.
- 15- Abdelgawad A, Naga O (2014)Birth injuries and orthopedic manifestations in newborns. In: Abdelgawad A and Naga O, Eds. *Pediatric Orthopedics: Handbook for primary care physicians*. Springer New York; 75-83.
- 16- Muhling RS, Blaauw G, Slooff ACJ, Kortleve JW, Tonino AJ (2001) Conservative treatment of obsterical brachial plexus palsy and rehabilitation. In: Gilbert A. *ed. Brachial plexus injuries*. London, Martin Dunitz Ltd 173-187.
- 17- Covey DC, Riordian DC, Milstead ME, Albright JA (1992) Modification of the L'Episcopo procedure for brachial plexus birth palsies. *J Bone Joint Surg* [Br] 74:897-901.
- 18- Gilbert A, Brockman R, Carlioz H (1991) Surgical treatment of brachial plexus birth palsy. *Clin Orthop*. 264:39-47.
- **19- Raimondi PL, Muse A, Saporiti E** (2001)Palliative surgery: shoulder paralysis. In: Gilbert A. ed. *Brachial plexus injuries*. London, Martin Dunitz Ltd225-238.
- **20- Hoffer M, Wickenden R, Roper B (1978)** Brachial plexus birth palsies. Results of tendon transfers to the rotator cuff. *J Bone Joint Surg.* 60(A):691-695.
- 21- Bahm J and Ocampo-Pavez C (2009)Monopolar teres major muscle transposition to improve shoulder abduction and flexion in children with sequelae of obstetric brachial plexus palsy. J Brachial Plexus and Peripheral Nerve Injury 4:20.
- 22- Chen L, Gu Y, Hu S (2002) Applying transfer of trapezius and/or latissimus dorsi with teres major for recontruction of abduction and external rotation of the shoulder in obstetrical brachial plexus palsy. J Recons Microsurg. 18:275-280.
- 23- Ahmad MR, El-Sadek AN, Abdeldaym A (2011)Compined latissimus dorsi and trapezius transfer to improve zero shoulder after brachial plexus palsy. Egypt J. plastic reconstr. Surg. 35(1);83-92.
- 24- Al-Qattan MM (2003)Latissimus dorsi transfer for external rotation weakness of the shoulder inobstetric brachial plexus palsy. *J Hand Surg [Br]*28:487-490.
- 25- Al-Qattan MM (2002)Rotation osteotomy of the humerus for Erb's palsy in children with humeral head deformity. J Hand Surg Am. 27:479–483.
- 26- Waters PM, Bae DS (2006) The effect of derotational humeral osteotomy on global shoulder function in brachial plexus birth palsy. *J Bone Joint Surg Am.* 88:1035–1042.
- 27- Abzug JM, Chafetz RS, Gaughan JP, Ashworth S, Kozin SH (2010) Shoulder function after medial approach and derotational humeral osteotomy in patients with brachial plexus birth palsy. *J Pediatr Orthop.*. 30(5): 469 74.

28- Abdelgawad A. and Miguel A. Pirela-Cruz (2014) Humeral Rotational Osteotomy for Shoulder Deformity in Obstetric Brachial Plexus Palsy: Which Direction Should I Rotate? *The Open Orthopaedics Journal* 8, 130 – 134.